

A Revision of the Stylasteridae (Cnidaria, Hydrozoa, Filifera) from Alaska and Adjacent Waters

Stephen D. Cairns^{1,†}, Alberto Lindner^{2,‡}

1 Department of Invertebrate Zoology, National Museum of Natural History, Smithsonian Institution, Washington DC 20560, USA **2** Departamento de Ecologia e Zoologia, Universidade Federal de Santa Catarina, Florianópolis, SC 88040-970, Brazil

† urn:lsid:zoobank.org:author:337544DA-F365-4DBA-816E-0EA934CD1292

‡ urn:lsid:zoobank.org:author:150D52B1-3586-4667-AC07-E295C757FB36

Corresponding author: Stephen D. Cairns (cairnss@si.edu)

Academic editor: B. Hoeksema | Received 12 August 2011 | Accepted 10 November 2011 | Published 22 December 2011

urn:lsid:zoobank.org:pub:BDBD8901-0EEC-4134-8000-AB84006E5E57

Citation: Cairns SD, Lindner A (2011) A Revision of the Stylasteridae (Cnidaria, Hydrozoa, Filifera) from Alaska and Adjacent Waters. ZooKeys 158: 1–88. doi: 10.3897/zookeys.158.1910

Abstract

The stylasterid fauna of Alaska is revised, consisting of the description or redescription and illustration of 21 species, one additional subspecies, and a geographically adjacent species: *Stylaster venustus*. Six new species and one new subspecies are described: *Errinopora fisheri*, *E. undulata*, *E. disticha*, *E. dichotoma*, *Stylaster crassiseptum*, *S. repandus*, and *Stylaster parageus columbiensis*. Four subspecies are raised to species rank: *Stylaster leptostylus*, *S. trachystomus*, *S. parageus*, and *Distichopora japonica*, and five species and one subspecies were synonymized. A dichotomous key to the *Errinopora* species and tabular keys to the *Errinopora* and Alaskan *Stylaster* species are provided. The focus of the study was on the stylasterids from Alaska, primarily those from the diverse Aleutian Islands, but also including records from British Columbia. This is the first revisionary work on this fauna since the seminal report by Fisher in 1938.

Keywords

Stylasteridae, Hydrozoa, Cnidaria, Aleutian Islands, Alaska, British Columbia, *Stylaster*, *Errinopora*

Introduction

The history of the published information about Alaskan stylasterids is not long. The first stylasterids to be reported from off Alaska were beach-worn specimens or coralla obtained from very shallow water from the Aleutian Islands and the Alaskan Penin-

sula (Dall 1884): *Stylaster verrillii* (Dall, 1884) and *Stylantheca papillosa* (Dall, 1884). Dall's report also included a description of *Stylaster moseleyi* (Dall, 1884), which is now considered to be a junior synonym of *S. verrillii*. Broch (1936) later reported *Stylaster boreopacificus typica* from Yukatat Bay, which is herein reidentified as *Stylaster parageus parageus*.

The major contributor to this fauna, however, was W. K. Fisher (1938), who, based on specimens from 19 stations from the Pacific Expeditions of the US Fish Commission Steamer *Albatross*, described 12 sp. n. and four new subspecies from off Alaska, mainly the Aleutian Islands. Herein, five of those species and 1 subspecies are synonymized, and three of his subspecies are raised to species status. Fisher was a careful observer, with a good eye for variation, but he did not have access to scanning electron microscopy and a much larger collection than his is now available, and thus we have made some changes to his taxonomy. Most of these specimens are deposited at the NMNH, and a representative set exists at the CAS.

Although no specimens were reported from the Aleutians, Naumov (1960) reported many of Fisher's species from the Kurile Islands and the Sea of Okhotsk; these have not been verified. Eguchi (1968) described *Allopora abei* Eguchi, 1968, from an unknown location in the Aleutian Islands, this species presumed to be a junior synonym of *Stylaster brochi* (Fisher, 1938).

Twenty years ago, Cairns (1991) described the deep-water, lamellate sp. n. and genus *Cyclohelia lamellata* from off Pribilof Islands. Cairns and Macintyre (1992) used seven Alaskan species for calcium carbonate polymorph determinations, some of which have been re-identified herein. Heifetz (2002) and Heifetz et al. (2005) listed the Alaskan species in an effort to analyze distribution and abundance to help in fishery management efforts, although their identifications were not accurate. Wing and Barnard (2004) produced a field guide to the Alaskan corals, which included a key and brief discussions of 37 stylasterid taxa, however their taxonomy was not current and is corrected herein. As part of his dissertation, Lindner (2005) revised the genus *Errinopora*, including a key to all species and the description of four sp. n., which are officially described herein. Stone and Shotwell (2007) uncritically listed all species known from the Alaskan region, and Brooke and Stone (2007) discussed reproductive aspects of nine Alaskan species. Finally, Lindner et al. (2008) used molecular sequences of three genes (nuclear rDNA 18S, Calmodulin, and mitochondrial rDNA 16S) of 11 Alaskan stylasterid species in a phylogenetic study of the family.

Material and methods

Abbreviations:

AB	Auke Bay Marine Lab (NOAA), Juneau, Alaska
AL	Alberto Lindner
<i>Alb</i>	USFWS <i>Albatross</i>
CAS	California Academy of Sciences, San Francisco

coll.	private collector
FOMC	Fiber Optic Monitoring Cruise of 2008
GM	Geologisk Museum ved København Universitet, Copenhagen, Denmark
H:D	Height to Diameter ratio of a gastrostyle
indet.	gender indeterminate
MCZ	Museum of Comparative Zoology at Harvard, Cambridge
MF	<i>Miller Freeman</i>
NMNH	National Museum of Natural History, Smithsonian Institution, Washington, DC
NOAA	United States National Oceanic and Atmospheric Administration
RACE	Resource Assessment and Conservation Engineering
USNM	United States National Museum (now the NMNH)
WoRMS	World Register of Marine Species

Specimens were obtained from collections deposited at the following institutions: CAS; MCZ; GM; and the NMNH. Specimens were also collected by one of us (AL) during NOAA's RACE bottom trawling survey off the Gulf of Alaska and the Aleutian Islands in 2001 and 2002. Additional specimens were provided by Mr. Arthur Schultz and by NOAA's Alaska Fisheries Science Center. We examined type specimens of all species reported herein.

Specimens were studied using scanning electron microscopy using the methodology as described by Cairns (1983). Genomic DNA and 37 new mitochondrial rDNA 16S sequences for the six *Errinopora* species from Alaska (GenBank accession numbers: JN572403-JN572439) were obtained as described in Lindner et al. (2008), using primers as described in Cunningham and Buss (1993). Sequences were aligned using ClustalX (Thompson et al. 1997) and phylogenetic analyses were performed in PAUP* 4.0 (Swofford 1999). Morphological terms used herein are defined in Cairns (2011), and the term pseudocyclosystem is defined as a cyclosystem-like structure (composed of a gastropore surrounded by dactylopores) that may be found at basal branches of some stylasterid species, and in which the dactylopores are usually not fused with the central gastropore and may not have the slits (dactylotome) of the dactylopore spines directed towards the central gastropore. In contrast to cyclosystemate stylasterids (e.g., *Stylaster*, *Conopora*, *Crypthelia*), pseudocyclosystemate species (e.g., *Errinopora*) have this polyp arrangement at basal portions of branches and colonies, and a different polyp arrangement at branch tips.

The specimens on which this study was based resulted from collections from both the United States government and university vessels as well as from privately owned fishing vessels, the latter often having many with colourful names. The US research vessels owned by the US Government are the *Albatross* and the *Miller Freeman*, whereas fishing vessels temporarily chartered by the US Government (for NOAA's RACE surveys) included: *Dominator*, *Sea Storm*, *Pacific Knight*, and the *Vesteraalen*. Additional private fishing vessels included: *Alaska Beauty*, *Alaska Sea*, *Alaska Trojan*, *Alaskan Leader*, *Ballyhoo*, *Delta Alfa*, *North Pacific*, *Ocean Olympic*, *Patricia Lee*, and *Shishaldin*. Additionally, the manned submersible *Delta* and the Remotely Operated Vehicles *Jason II*

and *Ropos* were also used to collect specimens as part of NOAA expeditions. The types of all species known from have been examined with the only exception of *Allopora abei*. All of these types are deposited at the NMNH and a subset at the CAS. Unless otherwise stated, the condition of specimens examined is assumed to be in the dry state.

Taxonomy

Family Stylasteridae Gray, 1847

Genus *Distichopora* Lamarck, 1816

<http://species-id.net/wiki/Distichopora>

Diagnosis. Colonies usually flabellate, with blunt branch tips. Coenosteum reticulate-granular and of many colors. Gastropores aligned along branch edge or, more rarely, meandering in lines on colony faces, the gastropore rows flanked (usually on both sides) by rows of horseshoe-shaped dactylopore spines. Gastro- and dactylopore tubes axial, extending along branch axis. Dactylostyles absent. Gastrostyles elongate (needle-shaped), often stabilized by transverse tabulae; a diffuse ring palisade sometimes present. Ampullae usually superficial and clustered.

Discussion. Equally diverse in tropical and temperate waters as well as in shallow and deep environments, *Distichopora* is known from 26 Recent species and two fossil species (see Appeltans et al. 2001: WoRMS data base: www.marinespecies.org). The genus was revised by Boschma (1959) and Cairns (1983b).

Type species. *Millepora violacea* Pallas, 1766, by monotypy.

Distribution. Eocene to Recent: cosmopolitan except for off Antarctica, 1–806 m.

Distichopora borealis Fisher, 1938

http://species-id.net/wiki/Distichopora_borealis

Figs 1A–C, 2A–K

Distichopora borealis Fisher, 1938: 543–545, pl. 70, fig. 3, pl. 71–73.—Broch 1942: 20.—Boschma 1957: 41 (listed).—Naumov 1960: 562, text-fig. 407.—Wing and Barnard 2004: 10, 27, fig. 26.—Heifetz et al. 2005: 133, 137 (listed).—Jamieson et al. 2007: 224 (listed).—Brooke and Stone 2007: 529, fig. 2I.—Stone and Shotton 2007: 107 (listed).—Lindner et al. 2008: 3, and supplemental Table S1: 3 (phylogeny and DNA sequences).

Distichopora sp. Heifetz, 2002: 22 (listed).—Heifetz et al. 2005: 133 (listed).

Type material. Holotype, *Alb*-3480, a female colony (dry) of 9 cm length (USNM 43274, Fig. 1B). Paratypes, *Alb*-3480, include 3 male, 4 female, 1 indeterminate dry colonies, and SEM stubs 1489–1494 (USNM 76810) and 2 dry female colonies from




Figure 1. Color figures of skeleton of various Alaskan stylasterids: **A–C** *Distichopora borealis* **D–I** *Cyclobelia lamellata* **A** arborescent colony, USNM 1122878 **B** holotype, USNM 43274 **C** largest known colony, USNM 1122542 **D** holotype, USNM 85077 **E** digitate form, USNM 1122492 **F** intermediate form between lamellar and digitate, USNM 96236 **G** juvenile corallum **H–I** side and apical views of fully developed lamellar colony.

Alb-4781 (USNM 76375). **Type locality.** *Alb*-3480: 52°06'00"N, 171°45'00"W (Amukta Pass, Aleutian Islands), 518 m.

Material examined. *Alaskan Leader* 40, 52°09'18"N, 175°40'42"W, 174 m, 8 Jun 2002, 1 female and 1 male, USNM 1122542; *Alaskan Leader* 54–14, 51°44.4'N, 178°16.7'W, 567–680 m, 11 Jun 2002, 6 indet., AB02–29; *Alaska Trojan*, 51°24'47"N, 178°50'02"W, 640 m, 2 Feb 2000, 1 female, USNM1122447; *Ballyhoo*, 54°49'52"N, 178°43'14"E, 236, 7 June 2000, 1 female, USNM 1122565; *Delta* 5600, 52°33'47"N, 179°26'45"W, 225 m, 15 Jul 2002, 1 male, USNM 1122543; *Dominator* 971–181, 51°27'43"N, 178°35'20"E, 384 m, 27 Jul 1997, 1 female and 1 male, USNM 1123357; *Dominator* 971–219, 53°00'03"N, 172°18.86'E, 223 m, 4 Aug 1997, 1 female, 2 male, USNM 1123358; *Jason*2–2103–7-4, 51°47.898'N, 179°57.169'E, 1267 m, 1 female, AB08–0036; *MF* 70, 52°03'24"N, 179°25'06"E, 174 m, 1 male, USNM 77047; *Ocean Olympic*, 52°04.78'N, 177°13.39'E, 256 m, 1 male, AB00–57; *Ocean Olympic*, 52°08.67'N, 176°35.46'E, 292 m, 1 female, AB00–21; *Ommaney* 8, 56°11'42"N, 135°06'31"W, 53 m, 17 Aug 2006, 1 female in alcohol, USNM 1086324; *Patricia Lee*, 51°53.44'N, 179°47.7'E, 298 m, 1 indet., AB00–41; *Sea Storm* 36, 53°05'45"N, 171°41'56"E, 458 m, 19 Jun 2002, 1 female and 1 male in alcohol, USNM 1076507 and 1122770; *Sea Strom* 92, 51°33'34"N, 177°36'59"W, 367 m, 4 Jul 2002, 7 males, USNM 1137601, 1123541, 1122874–76; *Sea Storm* 93, 51°50'59"N, 178°26'02"E, 390 m, 5 Jul 2002, 2 male, USNM 1122877 and 1123287; *Sea Storm* 105, 52°08'59"N, 176°02'17"E, 201 m, 8 Jul 2002, 2 female in alcohol, USNM 1122884; *Sea Storm* 106, 52°10'46"N, 175°10'34"E, 165 m, 8 Jul 2002, 1 female and 1 male in alcohol, USNM 1122885 and 1123288; *Sea Storm* 107, 52°10'28"N, 175°14'14"E, 214 m, 8 Jul 2002, 4 female and 2 males in alcohol, USNM 1122866–70; *Sea Storm* 108, 52°11'32"N, 175°17'E, 208 m, 8 Jul 2002, 1 female and 3 males, USNM 1122862–65; *Sea Storm* 109, 52°17'16"N, 175°20'56"E, 238 m, 8 Jul 2002, 6 female, 8 male, 3 indet. in alcohol, USNM 1122886 and 1123289; *Sea Storm* 122, 52°02'49"N, 175°16'54"E, 143 m, 13 Jul 2002, 2 female and 7 males, USNM 1122858–61; *Sea Storm* 132, 52°12'02"N, 176°05'56"E, 150 m, 15 Jul 2002, 1 female in alcohol, USNM 1122854; *Sea Storm* 133, 52°13'40"N, 176°02'13"E, 148 m, 15 Jul 2002, 1 male, 2 females in alcohol, USNM 1122855–57; *Sea Storm* 149, 52°30'15"N, 173°33'03"W, 239 m, 21 Jul 2002, 1 male in alcohol, USNM 1122883; *Sea Storm* 151, 52°33'40"N, 173°33'03"W, 203 m, 21 Jul 2002, 1 female and 2 male, USNM 1122878–80; *Sea Storm* 155, 52°38'43"N, 173°34'31"W, 393–401 m, 22 Jul 2002, 1 male and 1 indet., USNM 1122881–82; *Shishaldin*, 53°56'24"N, 179°49'31"E, 318 m, 14 Mar 2000, 1 female and 1 male, USNM 1122549; *Shishaldin*, 54°07'N, 179°45'E, 366 m, 20 Feb 2000, 1 female and 1 male, USNM 1122544–455; *Shishaldin*, 54°23'29"N, 179°19'46"E, 413 m, 12 Feb 2000, 1 male, USNM 1122548; *Vesteraalen* 941–165, 51°34'N, 178°19'E, 470 m, 18 Jul 1994, 1 female and 1 male, USNM 96247; Slear, coll., 51°59'52"N, 176°47'05"E, 241 m, 13 Nov 2000, 1 male, USNM 1122458; University of Washington, 51°32'N, 179°15'W, 278–289 m, 1 Sep 1968, 1 female, USNM 76377; 54°20'15"N, 133°03'19"W, 457–466 m, 1 Sep 2002,




Figure 2. Paratypes of *Distichopora borealis*, USNM 76810: **A** stereo view of carinate branch face **B–C** pore row, B being a stereo view **D** branch face with longitudinal ridges **E–F** coenosteal texture **G** gastrostere **H** dactylospine **I** stereo view of two gastrostyles **J** stereo view of pore row and female ampullae **K** stereo view of branch face and male ampullae.

1 female, USNM 1123535; 51°13'43"N, 179°49'31"E, 465–529 m, 13 Jun 2000, 1 male, USNM 1137421.

Description. Colonies usually uniplanar (Fig. 1B), but occasionally multiplanar or arborescent (Fig. 1A, C). Largest specimen examined (USNM 1122542, Fig. 1C) 11 cm tall and 20 cm broad, having a basal diameter of 3 cm, colonies broader than tall not uncommon. Base of colony broadly encrusting. Branching irregularly dichotomous, occasionally anastomotic. Large basal branches circular to elliptical in cross section, but terminal branches have flattened faces, thus rectangular in cross section. Branch tips blunt and rounded. Coenosteum white to light orange, the latter usually having branches with a white core. Coenosteum reticulate-granular in texture, the strips being 0.10–0.14 mm in width, each strip covered with tall (22 μ m tall, 9 μ m in diameter), slender granules arranged 8–10 across the width of a strip. Faces of distal branches often bear longitudinal ridges (Fig. 2A, D, K), some ridges up to 4 mm in length, sometimes ending distally in what appears to be a dactylopore opening (Fig. 2D).

Pore rows well defined (Fig. 2B), 1.0–1.3 mm in width, and restricted to lateral branch edges, although isolated rows and some irregularities may occur proximally on large colonies. Gastropores linearly arranged in a shallow sulcus, closely spaced (approximately 12–16 per cm), and circular in shape, measuring 0.33–0.41 mm in diameter. Gastropore tubes long and slightly curved, a diffuse ring palisade present near gastrostyle tip, the elements globular and about 30 μ m in diameter; tabulae absent. Gastrostyles long, slender, and fragile, up to 1 mm in length and 0.09 mm in diameter, having high H:W ratios up to 8–10. Gastrostyles longitudinally ridged and bear numerous long slender spines up to 0.07 mm in length and 10–12 μ m in diameter. U-shaped dactylopore spines arranged on both sides of gastropore row, but often more common on one side than the other, e.g., 18–20 dactylopore spines on one side opposite 28–40 spines on other side. Dactylopore spines up to 0.4 mm tall and 0.25–0.30 mm in width, with a dactylotome width of 0.07–0.08 mm, always directed toward or slightly obliquely to the adjacent gastropore. Dactylostyles absent.

Female ampullae (Fig. 2J) prominent, superficial, hemispherical mounds clustered on branch faces, 1.0–1.5 mm in diameter. Ampullae usually covered by low ridges, either radiating from the central apex or arranged in parallel, longitudinal rows aligned with branch axis. Lateral efferent pores 0.25–0.30 mm in diameter. Male ampullae (Fig. 2K) also superficial mounds, but mostly embedded in coenosteum (internal), also occurring in clusters on branch faces. Male ampullae irregular in shape, often with an apical tip, and smaller than female ampullae (0.35–0.45 mm in diameter).

Remarks. Among the 26 Recent species of *Distichopora* (see Cairns et al. 1999; Cairns 2005; and Appeltans et al. 2011: WoRMS database – www.marinespecies.org), the most similar is *Distichopora borealis japonica* Broch, 1942, described from Sagami Bay, Japan (110 m). Although similar in gross morphology, subspecies *japonica* differs in coenosteal texture (porcellaneous), and in ampulla shape, each female ampulla having more than one efferent pore and the male ampullae clustered densely in a continuous mat. Also, the dactylotomes of the dactylopore spines are oriented upward to outward (away from the gastropores), not

inward toward the gastropores. These differences would argue for an elevation to the species level of this putative subspecies. The observations are based on examination of a fragment of a syntype of *D. borealis japonica* (USNM 76868) and additional Japanese material deposited at the NMNH (USNM 44198, 44199, 44202, and 44217). Of the 111 specimens examined, 45 are female, 53 male, and 13 indeterminate in gender.

Distribution. Common in the Aleutian Islands from the Near Islands to Amukta Pass, including Bowers Bank; off Cape Ommaney, Alexander Archipelago, and Dixon Entrance, Queen Charlotte Islands; 53–1267 m, the shallowest record being from Cape Ommaney.

Genus *Cyclohelia* Cairns, 1991

<http://species-id.net/wiki/Cyclohelia>

Diagnosis. Colonies lamellar, often attaining a large size. Coenosteum reticulate-granular, reddish-orange. Gastro- and dactylopores uniformly distributed on corallum faces, of the axial type. Dactylopore spines circular to elliptical, enclosed by a thin wall for entire perimeter. Dactylostyles absent. Gastrostyles elongate but tabulae absent; a diffuse ring palisade present. Ampullae internal.

Discussion. The genus is monotypic. According to the molecular analysis of Lindner et al. (2008) it groups within the *Distichopora* clade as a sister species of *D. borealis*. The close relationship to *Distichopora* was also suggested by Cairns (1991) in the original description of *Cyclohelia*, both genera having axial gastro- and dactylopores, elongate gastrostyles, and in lacking dactylostyles. *Cyclohelia* differs only in having non-linear gastro- and dactylopores, internal ampullae, and differently shaped dactylopore spines. The colony morphology of *Distichopora anceps* Cairns, 1978 is remarkably similar to that of typical *Cyclohelia*, i.e., lamellar with a cleft dividing the colony into two adjacent undulating fronds. However, large specimens of *Cyclohelia* tend to form almost entirely lamellate colonies, whereas large specimens of *D. anceps* remain clearly lamellate-branching (Cairns 2005), rendering the colonies of *D. anceps* as a “transition” in morphology between the two genera.

Type Species. *Cyclohelia lamellata* Cairns, 1991, by original designation.

Distribution. Known only from the Aleutian Islands and Pribilof Islands, 27–567 m.

***Cyclohelia lamellata* Cairns, 1991**

http://species-id.net/wiki/Cyclohelia_lamellata

Figs 1D–I, 3A–C

Cyclohelia lamellata Cairns, 1991: 384–386, figs 1a–g, 2a–g.—Heifetz 2002: 22 (listed).—Wing and Barnard 2004: 10, 27, fig. 25, not Appendix fig. 1 (= *Errinopora undulata*).—Stone and Shotwell 2007: 107 (listed).—Brooke and Stone 2007: 529–530, figs 2K, 3E.—Jameison et al. 2007: 224 (listed).—Lindner et al. 2008: 3, and supplemental Table S1: 3 (phylogeny and DNA sequences).

Cyclohelia lamellate.—Heifetz et al. 2005: 133, 137 (listed).

Type material. The holotype, a dry female colony (USNM 85077, Fig. 1D). **Type locality.** Off Pribilof Islands, 550 m.

Material examined. *Alaskan Leader* 35, 53°01'48"N, 170°06'12"W, 172–178 m, 4 Jun 2002, 1 female, 1 male, USNM 1122491; *Alaskan Leader*, 54–14, 51°44.4'N, 178°16.7'W, 567–680 m, 11 Jun 2002, 1 female, AB02–0029; *Alaska Beauty*, 51°42.43'N, 176°56.05'E, 351 m, 3 Feb 2000, 1 female, AB00–15; *Delta* 5999–8E-5, 52°21.042'N, 179°30.483'W, 115 m, 4 Jul 2003, 3 fragments, AB10–0001; *Delta* 6000–10D-2, 51°50.8545'N, 179°49.6488'E, 150 m, 5 Jul 2003, 1 indet., AB; *Delta* 6001–10C-1, 51°52.396'N, 179°46.191'E, 230 m, 5 Jul 2003, 1 female, AB; *Delta* 6213–12B-1, 2 and 4, 6 males, AB; *Delta*, 51°50'55"N, 179°48'35"W, 125 m, 17 Jul 2002, 1 female in alcohol (digitate form), USNM 1122494; *North Pacific*, 52°04.28'N, 176°05.35'E, 366 m, 1 indet., AB00–28; *Ocean Olympic*, 52°11.89'N, 176°17.3'E, 366 m, 20 Oct 2000, 1 female (digitate form), USNM 1122513; *Pacific Knight* 941–121, 51°38'00"N, 178°19'00"W, 0–373 m, 5 Jul 1994, 1 female and 1 male, USNM 96235 and 1122900; *Pacific Knight* 941–179, 52°00'N, 176°44'E, 0–90 m, 23 Jul 1994, 1 female (intermediate form), USNM 96236; *Patricia Lee*, 51°18'42"N, 179°30'04"E, 329 m, 20 Oct 2000, 1 male, USNM 1122488; *Patricia Lee* 51°53.44'N, 179°47.7'E, 298 m, 3 male, SEM stub 1488, AB00–41; *Sea Storm* 107, 52°10'28"N, 175°14'14"E, 214 m, 8 Jul 2002, 1 female in alcohol, USNM 1123297; *Sea Storm* 108, 52°11'32"N, 175°17'E, 208 m, 8 Jul 2002, 10 female, 6 male, 8 indet. (digitate form) in alcohol and dry, USNM 1122951, -54, -55, -57, -58, -59, 61, 1123290, 91; *Sea Storm* 111, 52°16'20"N, 175°59'13"E, 137 m, 9 Jul 2002, 1 male in alcohol, USNM 1122949; *Sea Storm* 112, 52°15'12"N, 176°10'42"E, 137 m, 9 Jul 2002, 1 indet., USNM 1122946; *Sea Storm* 116, 52°04'10"N, 177°14'25"E, 87–94 m, 11 Jul 2002, 1 male in alcohol (digitate form), USNM 1122942; *Sea Storm* 122, 52°02'49"N, 179°25'18"E, 143 m, 13 Jul 2002, 8 female, 4 male, 5 indet., USNM 1122899, 1122948, 1123294–96, -98; *Sea Storm* 123, 52°10'53"N, 179°37'02"E, 124 m, 13 Jul 2002, 2 male, USNM 1122952; *Sea Storm* 125, 52°12'53"N, 179°56'49"W, 89–96 m, 13 Jul 2002, 1 male, USNM 112305; *Vesteraalen* 3, 52°38'13"N, 169°47'14"W, 75–83 m, 22 May 2001, 1 male in alcohol, USNM 1076490; *Vesteraalen* 941–36, 52°56'N, 169°31'W, 0–227 m, 10 Jun 1994, 1 female, USNM 96237; *Vesteraalen* 941–46, 53°04'N, 170°09'W, 0–174 m, 12 Jun 1994, 1 female, USNM 96238; *Vesteraalen* 941–153, 52°10'N, 179°43'E, 0–94 m, 11 Jul 1994, 1 male, USNM 96239; "Vessel" 35, 51°45'48"N, 178°09'47"W, 200–400 m, 4 Jun 2000, 2 female and 1 male, USNM 1122493; Renfro, coll., 51°56'35"N, 179°17'58"E, 236 m, 1 female, USNM 1122456; coll. Slear, 51°59'52"N, 176°47'05"E, 241 m, 13 Nov 2000, 1 male (digitate form), USNM 1122457; Petrel Bank, 1 male, 1 indet., USNM 1123540; 51°52.11'N, 179°49.51'W, 27 m, 17 Jul 2002, 1 female (digitate form), USNM 1122492.




Figure 3. *Cyclohelia lamellata*, USNM 1122488: **A** stereo view of male ampullae in cross section **B–C** male efferent pores.

Remarks. This species is not redescribed herein as it was adequately described originally (Cairns, 1991) and, being the only species in the genus, conforms to the genus diagnosis above. But, the species was previously known from only two specimens, the type and a specimen sequenced by Lindner et al. (2008), and thus more can be added to its description, including details of the male ampullae, based on the 81 additional specimens reported herein.

Colonies are usually firmly attached to a hard substrate by a robust cylindrical base and stem (Fig. 1H) up to 2.5 cm in diameter, which, at the height of about 2 cm, bifurcates into two lamellae or sheets, each sheet increasing its surface area by folding and undulating its surfaces into a complex three dimensional structure. Large colonies may attain a height of about 8 cm and a width of 10 cm, colonies wider than tall being the norm. However, a subset of colonies from seven stations (see Material Examined), termed the “digitate form”, differ from the typical colony morphology in having dissected plates, resulting in numerous flattened lobes and clavate branches (Fig. 1E). Furthermore, one colony (Fig. 1F) is intermediate between these two extremes, suggesting that these shapes are simply variations of the same species.

Male ampullae (Fig. 3A) entirely internal, and elliptical in shape, the greater axis of the ellipse perpendicular to the branch surface, unlike that of the female ampullae. Male ampullae are up to 0.6 mm in greater diameter and 0.45 mm in lesser diameter, occurring in great concentrations and invariably present on almost any broken surface. Each ampulla communicates to the surface by a circular efferent duct, the duct about 0.07–0.09 mm in diameter, which is surrounded by small inward projecting granules, giving the pore a star-shaped appearance (Fig. 3B–C). Of the 81 specimens examined, 34 are female, 29 male, and 18 indeterminate in gender.

Distribution. In addition to the type locality off the Pribilof Islands, this species is common throughout the Aleutian Islands from the Rat Islands to the Islands of Four Mountains at 27–567 m, although it was more commonly collected from the western

part of this range at depths of 100–300 m. The digitate form is known only from the western Aleutian Islands from 27–366 m.

Genus *Errinopora* Fisher, 1931

<http://species-id.net/wiki/Errinopora>

Errinopora Fisher, 1931: 397; 1938: 536.—Boschma 1956: F102; 1957: 57.—Cairns 1983a: 123; 1983b: 462. —Lindner 2005: 79–88.

Protoerrina Broch, 1935: 59; 1936: 99–100.

Diagnosis. (emended from Cairns 1983b and Lindner 2005): Colonies uniplanar to slightly bushy; branches round, elliptical, or lamellar in cross section, often robust with blunt tips. Coenosteal texture reticulate-spinose (with wide slits resulting in a spongy texture) or reticulate-granular; exterior surface of dactylopore spines usually inconspicuously longitudinally ridged; coenosteum orange, pink, and white. One species, *Errinopora cestoporina*, bears numerous perforated mounds on surface. Dactylopores dimorphic, the most common, termed the primary dactylopore spine, is U-shaped and usually robust (thick-walled), occurring randomly, in pseudocyclo systems, or often laterally fusing to form rows or taller terraces that flank rows of gastropores. When dactylopore spines flank both sides of a gastropore row and their dactylotomes are directed toward the gastropores it is termed bilateral or distichoporine; if only one row of spines flank a row of gastropores, then unilateral. If isolated, dactylotomes usually abcauline in orientation. Much smaller flush dactylopores, termed secondary dactylopores, which lack dactylostyles, commonly scattered over coenosteum of many species. Dactylostyles usually well developed, easily seen from external view. Secondary dactylopores much smaller, flush with coenosteum, and lack styles. Gastropores also dimorphic, the primary gastropores being circular in outline, flush with coenosteum (having no lip), and arranged in irregular vertical rows, short horizontal rows, or randomly. Tabulae and ring palisades absent. Gastrostyles lanceolate, covered with longitudinal or oblique, spiny ridges. Smaller secondary gastropores much smaller, having only a small gastrostyle or none at all. Female ampullae superficial hemispheres, often without an obvious efferent pore. Male ampullae usually smaller hemispheres and spongy.

Discussion. The ten species in this genus are differentiated and compared in both a dichotomous key (see below) and tabular key (Table 1); six of them occur exclusively in the Aleutian Islands. Another species was tentatively assigned to this genus by Cairns (1983b), *Errinopora lobata* (Nielsen, 1919), a Paleocene fossil from Denmark. This species was re-examined by Bernecker and Weidlich (1990, 2005), based on subsequently collected non-type material from the Fasse Formation in Denmark. They noted that whereas the dactylopore spines were typical of *Errina* or *Errinopora*, the spines did not contain dactylostyles, and thus resembled *Errina* more than *Errinopora*. We have examined the holotype of *Labiopora lobata* Nielsen, 1919 (GM1782), which

is a uniplanar colony 10.2 cm tall and 8.0 cm wide, with branches about 0.5 cm in diameter embedded within a *Dendrophyllia* matrix. The colony has pores, or broken bulges, of three sizes: small and elongate (75–110 μm in width), medium and round to somewhat quadratic (0.3–0.53 mm wide), and large, round or somewhat triangular (up to 1.7 mm). The former are quite shallow and probably the result of a reticulate coenosteal surface, whereas the medium-sized pores are deep and possibly represent the gastropores, one of which has a cyclo-system-like structure 1.9 mm in diameter. The largest pores appear to be ruptured ampullae. None of them, however, resemble dactylopore spines like those reported by Nielsen (1919) or Bernecker and Weidlich (1990, 2005), suggesting that the species reported by the latter authors is neither *Errina* nor *Errinopora*. Likewise, the lack of dactylopore spines in the type of *L. lobata* precludes it from being *Errinopora*, and thus we currently suggest an *incertae sedis* placement of this species until further analysis.

Species within *Errinopora* are unique with the Stylasteridae in having both dimorphic gastro- and dactylopores, a condition first noted by Fisher (1938) for three Alaskan species but interpreted exclusively as secondary dactylopores. Careful examination of longitudinal sections of several of these pores reveal that they contain not dactylostyles, but rather small gastrostyles. This implies that most species of *Errinopora* (all but *E. fisheri* and *E. cestoporina* Cairns, 1983, see Table 1) have two types of gastrozooids (feeding polyps), a unique case for stylasterids but previously reported for the hydractiniids *Stylactaria conchicola* (see Namikawa et al. 1992). Secondary gastropores differ from primary gastropores in being narrower and deeper, and in having only a minute or no gastrostyle. All but one species (*E. fisheri*, see Table 1) also have dimorphic dactylopores: a large type surrounded by a prominent horseshoe-shaped spine, and smaller, flush pores only 40–110 μm in diameter, which do not have dactylostyles and are termed secondary dactylopores.

In comparison to other stylasterid genera, *Errinopora* is most similar to *Gyropora* Boschma, 1960, whose only species, *Gyropora africana* Boschma, 1960, has dactylopore spines and gastropores coordinated in pore rows, as in some species of *Errinopora*. *Errinopora* is also similar to *Errina* Gray, 1835 and *Errinopsis* Broch, 1935, whose species may also have thick-walled dactylopore spines. None of these genera, however, include species with dactylostyles, as in *Errinopora*. Among genera with species having dactylostyles, only species of *Errinopora*, *Inferiolabiata* Broch, 1951, and *Paraerrina* Broch, 1942 lack a coordination of gastropores and dactylopores in well-developed cyclo-systems (whereas species of *Stenohelia* Kent, 1870, *Stylantheca* Fisher, 1931, *Stylaster* Gray, 1831 and *Calyptopora* Boschma, 1968 do have both dactylostyles and well-developed cyclo-systems). *Inferiolabiata* differs from *Errinopora* in many characters, in particular by having thin-walled dactylopore spines (instead of thick-walled) that are markedly truncated at the tip (instead of rounded), whereas *Paraerrina* Broch, 1942 differs in having delicate dactylostyles (instead of robust) and either flush or only slightly raised dactylopore spines—instead of tall and robust (Cairns 1984, 1991). *Errinopora* is also one of the few stylasterid genera with species having calcitic, rather than aragonitic, colonies (Thompson and

Table 1. Tabular Key of the Ten Species of *Errinopora* (pscs = pseudocyclcosystems)

	<i>iE. fisher</i>	<i>aE. cestoporin</i>	<i>aE. porifer</i>	<i>aE. undulat</i>	<i>aE. distich</i>	<i>iE. pourtalesi</i>	<i>aE. stylifer</i>	<i>aE. zarhynch</i>	<i>aE. nanac</i>	<i>aE. dichotom</i>
Dactylopor Spines										
Arrangement	Tall terraces and pscs	Tall terraces and pscs	Isolated or short unilateral rows	Pscs and short unilateral rows	Distichopore rows; pscs at base	Distichopore rows, unilateral rows, isolated, pscs	Distichopore rows, pscs at base	Short unilateral rows, isolated	Short unilateral rows, pscs, isolated	Short unilateral rows, isolated
Compound Dactylopor	Absent	Absent	Absent	Absent	Present	Present	Absent	Present	Absent	Present
Spines										
Maximum Height	1.5 mm	1.3 mm	0.46 mm	0.25 mm	0.5–0.9 mm	1.1–1.2 mm	0.5–0.9 mm	to 3 mm	0.4 mm	1.2 mm
Wall Thickness	Not individualized	Thick	Thin	Thin	Thick	Thick	Thick	Thick	Thick	Thick
Dactylopor	Elliptical pores	Slit	Apical pores	Slit	Slit	Slit	Slit	Slit	Slit	Slit
Shape										
External Surface of Dactylopor	Inconspicuously ridged	Reticulate-Granular	Not ridged	Inconspicuously ridged	Inconspicuously ridged	Inconspicuously ridged	Inconspicuously ridged	Inconspicuously ridged	Prominently ridged	Inconspicuously ridged
Spine										
Dactylostyle	Narrow	Robust	Moderate width	Robust	Robust	Robust	Robust	Narrow	Moderate width	Robust
Secondary Dactylopor	Absent	Present	Present	Present	Present	Present	Present	Present	Present	Present
Corallum										
Shape	Uniplanar	Bushy	Uniplanar	Lamellate	Uniplanar	Bushy	Uniplanar	Uniplanar	Uniplanar branching or plate-like	Three dimensional
Branch Size	Delicate	Delicate	Delicate	Sheets thin	Robust	Medium	Medium	Large	Delicate	Robust
Branch Cross Section	Circular to elliptical	Circular	Circular to slightly elliptical	Lamellar	Flattened	Circular	Circular to slightly elliptical	Circular to elliptical	Circular to lamellar	Circular to slightly elliptical

	<i>iE. fisher</i>	<i>aE. cestoporin</i>	<i>aE. porifer</i>	<i>aE. undulat</i>	<i>aE. distich</i>	<i>iE. pourtalesi</i>	<i>aE. stylifer</i>	<i>aE. zarhynch</i>	<i>aE. nannee</i>	<i>aE. dichotom</i>
Dactyloporo Spines										
Color	Orange	White	White, light orange	Orange	Light orange	Orange	Orange, yellow	Orange	Orange, pink	Orange
Other Characters										
Gastropore Diameter	0.3–0.5 mm	0.30–0.55 mm	0.23–0.27 mm	0.30–0.45 mm	0.3–0.7 mm	0.20–0.38 mm	0.4 mm	0.2–1.1 mm	about 0.2 mm	0.3–0.5 mm
Secondary Gastropores	Absent	Absent	Present	Present	Present	Present	Present	Present	Present	Present
Coenosteal Texture	Reticulate-Spinose	Reticulate-Granular	Reticulate-Granular	Reticulate-Spinose	Reticulate-Spinose	Reticulate-Granular	Reticulate-Granular	Reticulate-Spinose	Reticulate-Spinose	Reticulate-Spinose
Unique Features	Gastropore tube constricted; ring palisade present; gastrostyles squar	Conical ampullae; coenenchymal papillae; gastrostyle pedicellate	Dactyloporo spines conical		Gastrostyles flattened in cross section	Encrusting base		Gastropore tubes commodious		U-shaped branching axils
Distribution and Depth Range	Aleutian Islands, 455 m	Tierra del Fuego, 359–384 m	Okhotsk Sea, 190–250 m	Aleutian Islands; 350–640 m	Aleutian Islands, 178–536 m	Central California, 49–183 m	Off Japan, Okhotsk Sea, 84–379 m	Aleutian Islands, 207–658 m	Aleutian Islands, 40–517 m	Aleutian Islands, 178–217 m

Chow 1955, Lowenstam 1964, Cairns and Macintyre 1992). The only other stylasterid genera with species known to have mostly calcitic colonies are *Errinopsis*, *Errina*, and one species of *Stylaster*—*Stylaster verrillii* (Dall, 1884) (see Cairns and Macintyre 1992). Within *Errinopora*, only *E. cestoporina*, known solely from the Subantarctic Region, is known to have coralla formed by precipitation of aragonite. This result confirms the more general observation of the prevalence of calcitic stylasterids in the North Pacific (Cairns and Macintyre, 1992), possibly related to the shallower depth of the Aragonite Saturation Horizon (ASH) in the Region (Guinotte et al. 2006).

In an attempt to investigate phylogenetic relationships, 37 partial mitochondrial rDNA 16S sequences were obtained for the six species of *Errinopora* from the Aleutian Islands, including the holotypes of *E. dichotoma*, *E. disticha*, *E. fisheri* and *E. undulata*. Based on Lindner et al. (2008), *Lepidotheca chauliostylus* Cairns, 1991 and *Inferiolabiata lowei* (Cairns, 1983) were used as the most closely related outgroups. The monophyly of *Errinopora* is well supported with 100% bootstrap support, but both maximum likelihood and parsimony analyses do not provide strong support for a particular relationship within the genus, resulting in a basal polytomy (results not shown). We thus aligned and analyzed solely the 37 *Errinopora* sequences but, similarly, no support was obtained for particular phylogenetic relationships within the genus (Fig. 4). Despite low bootstrap support, three of the six most parsimonious trees recover all morphologically-determined *E. nanneca* Fisher, 1938 specimens as a monophyletic clade, except specimen USNM1123524, which clusters together with *E. undulata*. The difficulties in estimating phylogenetic relationships for the Alaskan species of *Errinopora* using rDNA 16S may stem from relatively recent divergence times. The fossil-calibrated chronogram estimated by Lindner et al. (2008) included *E. nanneca* (specimen USNM1027820) and *E. zarhyncha* Fisher, 1938 (specimen USNM1071915), and shows that these species diverged only about 4 million years ago. Moreover, the same study shows that *Cyclohelia lamellata* Cairns, 1991 and *Distichopora borealis* Fisher, 1938, two sympatric Alaskan stylasterids that are also clearly distinguishable with marked morphological differences (see above), may have diverged as recently as 1 million years ago. These results indicate that part of stylasterid species diversity in Alaska may have diverged only within the past 1–4 million years and, at least for *Errinopora*, the results presented herein show that despite the marked morphological differences, some species are not recovered as reciprocally monophyletic lineages (Baum and Shaw 1995, Avise 2000) using mitochondrial rDNA 16S.

Type species. *Errina pourtalesii* Dall, 1884, by original designation.

Distribution. North Pacific: Aleutian Islands, Kurile Islands, Sea of Okhotsk, Sea of Japan, off California. Subantarctic: off Tierra del Fuego, 40–658 m.

Key to the Recent species of *Errinopora* (bold face = occurs off Alaska)

- 1 Dactylopore spines arranged in pseudocyclo systems near base but otherwise predominantly laterally fused into tall terraces flanking the lower side (uni-

- lateral) of one or a series of gastropores, their dactylotomes facing the gastropores **2**
- 1' Dactylopore spines isolated, and/or arranged in pseudocyclo systems, short unilateral rows, or bilateral (distichoporine) rows..... **3**
- 2 Individuality of fused dactylopore spines lost, the terraces having a continuous edge and the dactylotomes being elliptical pores; ampullae hemispherical; coenosteum orange; no coenosteal papillae; known only from Alaskan ***E. fisheri* sp. n.**
- 2' Fused dactylopore spines retain individuality, terraces having an irregular edge and dactylotomes being traditionally slit like; ampullae conical; coenosteum white; perforated papillae common on coenosteum; known only from the Subantarctic *E. cestoporina* Cairns, 1983
- 3 Dactylopore spines predominantly conical, having a circular dactylotome *E. porifera* Naumov, 1960
- 3' Dactylopore spines horseshoe-shaped, having a slit-like dactylotome **4**
- 4 Colonies lamellate (sheet-like) **5**
- 4' Colonies branching..... **6**
- 5 Colony a continuous, sinuous, thin sheet; pseudocyclo systems predominant form of dactylopore spine arrangement; dactylopore spines very short (0.25 mm) and thin walled ***E. undulata* sp. n.**
- 5' Colony a series of dissected blades of a larger sheet; pseudocyclo systems occur only at colony base, transverse rows of dactylopore spines being more common; dactylopore spines short (0.40 mm) and thick walled ***E. nanneca* Fisher, 1938 (lamellate form)**
- 6 When arranged in rows, dactylopore spines predominantly in bilateral (distichoporine) rows, the dactylotomes of both rows facing the flanked gastropore row (spines may also be isolated, in pseudocyclo systems, and/or arranged in unilateral rows) **7**
- 6' When arranged in rows, dactylopores spines predominantly in unilateral rows flanking a single row of gastropores, the dactylotomes of only the proximal row facing the gastropores (spines may also be isolated and/or arranged in pseudocyclo systems especially basally)..... **9**
- 7 Coenosteum reticulate-spinose; branches and gastrostyles flattened; known only from the Aleutian Islands ***E. disticha* sp. n.**
- 7' Coenosteum reticulate-granular; branches and gastrostyles circular in cross section; known from off California and off Japan and Russia **8**
- 8 Colonies bushy, with an encrusting base; ampullae small (0.5–0.9 mm in diameter); compound dactylopore spines present; known only from off California..... *E. pourtalesii* (Dall, 1884)
- 8' Colonies uniplanar without an encrusting base; ampullae large (up to 1.6 mm in diameter); compound dactylopore spines lacking; known only from off Japan and Russia..... *E. stylifera* (Broch, 1935)

- 9 Dactylopore spines tall (up to 3 mm), and often compound; dactylostyle narrow in respect to dactylotome; gastropore tube short and commodious, much larger than gastrostyle***E. zarhyncha* Fisher, 1938**
- 9' Dactylopore spines smaller (0.4–1.2 mm), rarely compound; dactylostyle moderate to large with respect to dactylotome; gastropore tube longer and less commodious, providing a close fit for the gastrostyle **10**
- 10 Dactylopore spines short (up to 0.4 mm); gastropores small (0.2 mm in diameter); colonies uniplanar, with angled branch axils.....
.....***E. nanneca* Fisher, 1938 (branched form)**




Figure 4. Midpoint rooting phylogeny of the Alaskan species of *Errinopora* estimated using mitochondrial rDNA 16S sequences. Numbers at nodes are the result of 100 bootstrap pseudoreplicates whenever >70. Shown is one of six most parsimonious trees (length: 38; CI: 0.868). Numbers preceded by ‘USNM’ represent specimen catalog numbers at the United States National Museum, Smithsonian Institution. Numbers preceded by ‘EU’ or ‘JN’ represent GenBank accession numbers. Asterisks (*) indicate the two specimens and rDNA 16S sequences of *Errinopora nanneca* and *Errinopora zarhyncha* previously published by Lindner et al. (2008).

- 10' Dactylopore spines taller (up to 1.2 mm); gastropores larger (0.3–0.5 mm in diameter); colonies 3-dimensional, with U-shaped branch axils
.....*E. dichotoma* sp. n.

***Errinopora fisheri* Lindner & Cairns, sp. n.**

urn:lsid:zoobank.org:act:9D4515E9-8067-4E6B-8C59-46B775683599

[http://species-id.net/wiki/Errinopora fisheri](http://species-id.net/wiki/Errinopora_fisheri)

Figs 5A, 6A–K

Errinopora fisheri Lindner, 2005: 100–102, figs 4.6A, 4.14 (unpublished name).

Type material. Holotype: *Pacific Knight* 204, a dry female colony, USNM 1123526 (Fig. 5A). Paratype: *Sea Storm* 36, 53°05'45"N, 171°41'56"E, 458 m, 1 branch, USNM 1157073. **Type locality.** 53°06'N, 171°42'E (off Attu Island, Aleutian Islands), 455 m, 31 Jul 1994.

Etymology. Named in honor of Walter K. Fisher, who wrote the original revision of the Aleutian Island stylasterids (Fisher 1938) and described the genus *Errinopora* (Fisher, 1931).

Material examined. Holotype.

Description. Holotype (Fig. 5A) uniplanar with a secondary flabellum at mid-height, 9 cm in height and 6.5 cm in width, with a basal branch diameter of 10 × 7.5 mm. Branches circular to slightly elliptical in cross section and irregularly dichotomous, small-diameter branches often diverging from larger main branches; branch anastomosis absent. Coenosteum reticulate-granular in texture, the strips being 80–90 µm in width and covered with small granules 5–15 µm in diameter; strips bordered by slits 40–45 µm wide. Coenosteum light orange.

Dactylopore spines favor one face of corallum over the other, arranged in abcauline, crescent-shaped terraces flanking one or two gastropores (Fig. 6A–D). Terraces sometimes almost completely surround a gastropore (Fig. 6C–D), resembling a cyclo-system of 1.5 mm diameter with an adcauline diastema, much as in *Stylaster*, but isolated dactylopore spines also fairly common, especially near base of colony. Dactylopore terraces quite thin (0.35 mm) and up to 1.5 mm in height, and, although laterally fused, the individuality of each dactylopore spine is subsumed into the more continuous terrace structure; compound dactylopore spines absent; exterior surface of spines longitudinally ridged. Terraces often horizontally oriented on branches, and slightly flared around gastropore(s). Dactylo-tomes not slit-shaped as in most other *Errinopora*, but instead small elliptical pores about 0.27 mm long and 0.15 mm in width occurring only near tips of spine structure; dactylo-tomes always directed toward a gastropore. Dactylostyles small, about 50 µm in width, having stubby elements up to 25 µm in length and 10–15 µm in diameter. Secondary dactylo-pores absent.

Gastropores circular, flush with coenosteum, 0.3–0.5 mm in diameter, being fairly consistent in size; gastropores isolated, not arranged in rows; secondary gastropores




Figure 5. Color figures of skeleton of various Alaskan stylasterids: **A** *Errinopora fisheri* **B, G–I** *E. nanneca* **C, F** *E. undulata*, **D** *E. disticha* **E** *E. zarhyncha*. **A** holotype, USNM 1123526 **B** holotype, USNM 42875 **C** holotype, USNM 112327 **D** holotype, USNM 1123524 **E** holotype, USNM 42874 **F** lateral view of a large paratype, USNM 1123528 **G** lamellate colony with digitate distal branches, USNM 1123462 **H** lamellate colony, USNM 44070 **I** intermediate form between lamellate and digitate, USNM 1123510.




Figure 6. Holotype of *Errinopora fisheri*, USNM 1123526: **A–D** branch tips with dactyloporous spines arranged in crescents (female ampulla shown in figure D) **E, H–I** lateral view of gastrostyles **F–G** coenosteal texture **J** detail of gastrostyle spines **K** dactylostyle.

absent. Gastropore tube cylindrical, but with a slight medial constriction; often a ring palisade is present, the elements 50 μm tall and 15–25 μm in diameter. Gastrostyle squat, an illustrated one being 0.33 mm tall and 0.29 mm in diameter (Fig. 6I). Gastrostyles covered with longitudinal anastomosing ridges as consistent with the genus. Female ampullae (Fig. 6D) hemispherical but often irregular in shape due to protrud-

ing dactylopore spines. Female ampullae 1.0–1.2 mm in diameter; efferent pores not observed. Male ampullae unknown.

Remarks. Even though represented by only one specimen, *Errinopora fisheri* can be distinguished from other Alaskan congeners by the distinctively terraced arrangement of its dactylopore spines, its small elliptically-shaped dactylotomes, its squat gastrostyles, and its constricted gastropore tube with a ring palisade (see Dichotomous Key and Table 1). Most of these features, however, are shared with *E. cestoporina*, a species known only from Burdwood Bank and off Tierra del Fuego at 359–384 m. *E. cestoporina* differs in having differently shaped ampullae, a different corallum color (white), small perforated mounds covering the coenosteum, and a differently shaped gastrostyle.

Distribution. Known only from Aleutian Islands: off Attu Island; 455–458 m.

***Errinopora undulata* Lindner & Cairns, sp. n.**

urn:lsid:zoobank.org:act:CCDBFA1A-6AA3-4C95-A85A-216A3A760B4C

http://species-id.net/wiki/Errinopora_undulata

Figs 5C, F, 7A–J

Errinopora undulata Lindner, 2005: 120–122, figs 4.6B, 4.19 (unpublished name).

Type material. Holotype: Vessel and collector unknown, Amukta Pass, 350–500 m, 1 dry female colony 12 cm tall, USNM 1123527 (Fig. 5C). Paratypes: Vessel and collector unknown, Amukta Pass, 356–640 m, 1996, 1 dry female colony, USNM 1123528; south of Semisopchnoi Island, 366 m, 1 dry female colony, USNM 88371. **Type locality.** Aleutian Islands: Amukta Pass, 350–500 m.

Etymology. The specific name *undulata* (from the Latin *undulatus*, meaning wavy) refers to the sinusoidal margin of the lamellate colonies.

Material examined. Types.

Description. Colonies lamellate, but wavy in construction resulting in a continuous, thin (1.7–2.0 mm thick), sinusoidal distal edge. Largest colony (USNM 1123528, Fig. 5F) 13 cm tall and 15 cm wide, with a massive basal branch 4.5 × 2.5 cm in diameter; holotype (Fig. 5C) smaller and less intact, measuring 12 cm tall and 11 cm wide, with a basal branch 3.5 × 2.8 cm in diameter. Parasitic spionid worm tubes found in only one colony (USNM 1123528). Coenosteum quite porous, consisting of a reticulum of narrow spiny strips separated by even wider slits, the spines being 10–15 µm in diameter. Coenosteum orange.

Dactylopore spines occur equally on both branch faces, but less abundant toward corallum base. Most dactylopores clustered around isolated gastropores in pseudocyclo systems, 3–5 spines surrounding one pore (Fig. 7B). Pseudocyclo systems most common toward base of colony but also occur frequently on more distal parts of corallum where they are interspersed with short, transverse rows of dactylopore spines that border the distal margins of one or more gastropores,




Figure 7. Female paratype of *Errinopora undulata*, USNM 1123528: **A–B, D** plate surface showing female ampullae, dactylospines, and gastropores **C** cross section of plate showing several gastrostyles **E–F** coenosteal texture **G–H** lateral view of a gastrostyle and detail of spination **I–J** lateral and apical view of a robust dactylostyle.

their dactylotomes facing upward (abcauline) toward the gastropore; compound dactylospines absent. Dactylospines fairly short (rarely taller than 0.25 mm), 0.40–0.45 mm in width, and thin walled, the majority of width being the dactylotome; dactylospine walls longitudinally ridged. Dactylostyles robust (Fig. 7I–J), up to 0.12 mm in width, consisting of cylindrical elements up to 72 μ m in length and 15 μ m in diameter. Secondary flush dactylospores common, about 75 μ m in diameter.

Gastropores circular, flush with coenosteum, and 0.3–0.45 mm in diameter; secondary gastropores about 0.19 mm in diameter. Gastropore tubes short and lack a ring palisade. Gastrostyles lanceolate, the figured style (Fig. 7G) 0.54 mm in height and 0.29 mm in diameter, covered with longitudinal anastomosing spiny ridges.

Female ampullae (Fig. 7A–B) irregularly-shaped, flattened hemispheres up to 1.6 mm in diameter. Efferent pores not observed. Male ampullae unknown.

Remarks. *Errinopora undulata* is quite similar to the lamellate form of *E. nanneca*, but differs in a number of points, one being its colony form, which is a continuous wavy sheet of corallum, whereas that of *E. nanneca* is more like a series of smaller dissected flat blades of a larger plate. *E. undulata* also has shorter (0.25 vs 0.4 mm) and thin-walled (vs thick-walled) dactylopore spines, and larger gastropores (0.45 mm vs 0.20 mm in diameter) (see Dichotomous Key and Table 1). Of the 247 known stylasterid species (Appeltans et al. 2011: WoRMS database: www.marinespecies.org) only five have adopted a lamellate corallum shape, four of these occurring in the Aleutian Islands (*Cyclohelia lamellata*, *Stylaster repandus*, *E. nanneca*, and *E. undulata*), the fourth being the Hawaiian *Distichopora anceps* Cairns, 1978.

Distribution. Known only from the Aleutian Islands: Amukta Pass and south of Semisopochnoi Island; 350–640 m (unconfirmed).

***Errinopora disticha* Lindner & Cairns, sp. n.**

urn:lsid:zoobank.org:act:3CA47961-430E-4FDF-BB00-15C6DDB19829

http://species-id.net/wiki/Errinopora_disticha

Figs 5D, 8A–J

Errinopora disticha Lindner, 2005: 97–99, figs 4.2A–B, 4.7B, 4.13 (unpublished name).

Type material. Holotype: a dry female colony 10 cm in height, coll. Renfro, USNM 1123524 (Fig. 5D). Paratypes: also from type locality, 1 female, USNM 1123521; *Alaskan Leader* 35, 53°01'48"N, 170°06.62'W, 172–178 m, 4 Jun 2002, 2 male, USNM 1123525; *Alb-3480*, 52°06'N, 171°45'W, 518 m, 8 Jul 1893, 2 males in alcohol, USNM 43768; *Alb-3480*, see above, 1 male, USNM 1148212 (ex USNM 52247, paratype of *E. zarhyncha*); 52°03'26"N, 179°12.34'E, 475 m, 27 Apr 2000, 1 male, USNM 1123523; 51°52'46"N, 179°17'24"E, 536 m, 16 Jul 2000, 2 female, USNM 1123522. **Type locality.** 51°53'12"N, 179°17'40"E (west of Semisopochnoi I., Petrel Bank), 530 m.

Etymology. The specific name *disticha* (from the Greek *distichos*, meaning “of two rows”) refers to the short distichoporine pore rows of this species.

Material examined. Types.

Description. Colonies uniplanar to multiplanar, consisting of cylindrical to highly compressed branches, the greater axis of compressed branches oriented in plane of colony flabellum and reaching up to two times length of lesser axis. Branching dichotomous and non-anastomotic, terminating in blunt tips and forming U-shaped axils be-




Figure 8. Holotype of *Errinopora disticha*, USNM 1123524: **A–B** distichoporine pore rows and female ampullae (**A** a stereo view) **C** gastropore and adjacent dactylospine with dactylostyle **D–E** coenosteal texture **F–G** dactylostyle **H–I** lateral view of gastrostyle **J** detail of gastrostyle spines.

tween branching. Holotype (Fig. 5D) 10 cm tall and equally wide, with a basal branch diameter of 22×20 mm; largest specimen (USNM 1123523) 13.5 cm in height. Parasitic spionid polychaete tubes found in only one corallum (USMM 1123525). Coenosteum typical for the genus: reticulate spinose, the narrow strips separated by broad porous slits, resulting in a very porous surface. Coenosteum light orange; branch cores white or lighter orange.

Dactylospine spines mostly arranged in long meandering distichoporine (bilateral) rows of up to 18 laterally fused spines (Fig. 8A–B). Two rows or terraces of dac-

tylopore spines usually flank each gastropore row, the dactylotomes facing the gastropore; however, the number of dactylopore spines is often unequal on either side of a pore row and are occasionally absent from one side. Toward corallum base, pore rows decrease in length, often resulting in a single gastropore surrounded by 3–6 dactylopore spines approximating a pseudocyclosystem. Dactylopore spines fairly short (0.5–0.9 mm in height) and thick-walled, the dactylotome consisting of one-third to half width of spine; compound dactylopore spines common. Dactylostyles robust, up to 75 μm in width, consisting of cylindrical elements up to 53 μm in length and 5–10 μm in diameter; exterior surface longitudinally ridged. Secondary dactylopores, which lack styles, very common on coenosteum, especially back sides of dactylopore spines, these pores being circular, flush with the surface, and measure about 75–110 μm in diameter.

Gastropores circular, flush with coenosteum, and variable in size, ranging from 0.3–0.7 mm in diameter, both size classes often adjacent to one another. Gastropores usually closely spaced and unilinearly arranged in a shallow sulcus created by adjacent dactylopore spine terraces. Gastropore tubes cylindrical, fairly shallow, and lack a ring palisade; secondary gastropores 0.19–0.30 mm in diameter. Gastrostyles lanceolate but somewhat stout, the figured symmetrical style (Fig. 8H–I) being 0.49 mm in height and 0.36 mm in diameter, however, many gastrostyles are asymmetrical, being elliptical to flattened in cross section. Gastrostyles bear longitudinal, spiny, anastomosing ridges, and fill most of gastropore cavity.

Female ampullae (Fig. 8A–B) hemispherical to somewhat flattened (1.3–1.5 mm in diameter), their basal perimeter often slightly undercut, and often with an irregular or ridged surface. Discrete efferent pores never observed. Male ampullae small, porous, hemispherical blisters 0.4–0.6 mm in diameter, often occurring in great concentrations on coenosteum.

Remarks. *E. disticha* is the only Alaskan *Errinopora* species to have its dactylopore spines arranged bilaterally in distichoporine rows flanking a row of gastropores, however two other species have a distichoporine arrangement: *E. pourtalesii* (California) and *E. stylifera* (Japan to Okhotsk Sea) (see Dichotomous Key and Table 1). *E. disticha* differs from those two species in having reticulate-spinose coenosteum (vs reticulate-granular), and flattened gastrostyles and branches. Of the 10 specimens examined, 4 are female and 6 male.

Distribution. Known only from the Aleutian Islands: off Petrel Bank, Amukta Pass, and off Four Kings Islands; 178–536 m.

Errinopora zarhyncha Fisher, 1938

http://species-id.net/wiki/Errinopora_zarhyncha

Figs 5E, 9A–L

Errinopora zarhyncha Fisher, 1938: 539–541, pl. 68, pl. 69, fig. 1.—Boschma 1957: 58.—Thomson and Chow 1955: 23, 30 (mineralogy).—Not Naumov 1960:

559.—Lowenstam 1964: 377 (mineralogy).—Lowe 1967: 82.—Cairns 1983b: 462.—Cairns and Macintyre 1992: 100–101 (mineralogy).—Cairns et al. 1999: 43 (listed).—Heifetz 2002: 22 (listed).—Wing and Barnard 2004: 27, 57, fig. 27.—Lindner 2005: 123–127, figs 4.2D, 4.10, 4.20 (redescription, key).—Heifetz et al. 2005: 133, 137 (listed).—Stone and Shotwell 2007: 108 (listed).—Jamieson et al. 2007: 224 (listed).—Lindner et al. 2008: 3, and supplemental Table S1: 3 (phylogeny and DNA sequences).

Errinopora zarhincha: Lowenstam, 1964: 382–383 (misspelling).

Type material. Holotype, *Alb*-3480, a dry male colony 14 cm in height, USNM 42874 (Fig. 5E). Paratypes: *Alb*-3480, 3 dry female branches, USNM 52247; *Alb*-3480, 1 branch, CAS 28186. **Type locality.** *Alb*-3480: 52°06'N, 171°45'W (Amukta Pass, Aleutian Islands), 518 m.

Material examined. *Shishaldin*, 53°56'24"N, 179°49'31"E, 318 m, 14 May 2000, 1 male, USNM 1123447; *Shishaldin*, 54°54'05"N, 178°37.27'E, 410 m, 11 Feb 2000, 3 female branches, and SEM stubs 1542–43, USNM 1123448; *Vesteraalen* 941–59, 52°02'00"N, 172°12'W, 658 m, 14 Jun 1994, 3 branches of a female colony, both dry and in alcohol, USNM 96234; *Vesteraalen* 941–62, 51°58'N, 172°40'W, 207 m, 15 Jun 1994, 1 female, USNM 96233; *Vesteraalen*, 51°13'17"N, 179°05'57"E, 488 m, 4 Jun 2000, 1 male, USNM 1071915; 51°13'43"N, 179°49'31"E, 465–529 m, 13 Jun 2000, large male colony, USNM 1123446; Types.

Description. Colonies uniplanar or multiplanar, robust, with fairly close dichotomous branching, leaving little space between branches; branch anastomosis occurs but uncommon. Largest colony 26 cm in height and 27 cm in width, with a massive basal branch 4.3 cm in diameter (USNM 1123446). Branches circular to slightly flattened in cross section, attenuating to thick (7–15 mm in diameter), blunt tips. Parasitic spionid polychaete worms form tubes along branch axes in two colonies (USNM 96233, 96234), the tubes Fig. 8 in cross section. Coenosteum quite porous in texture (Fig. 9E–H), consisting of a reticulum of thin (25 µm), spinose ridges, separated by wide slits or series of pores. This texture also present on ampullae and dactylopore spines, in the latter the coenosteal strips forming longitudinal ridges. Coenosteum orange.

Dactylopore spines occur on all branch surfaces and are quite variable in orientation, sometimes forming rows of 10–13 spines laterally fused into a short tier on one side of a pore row (unilaterally), the dactylotomes usually facing upward (abcauline), but dactylopore spines also oriented with their dactylotomes facing downward (adcauline) or laterally, and occasionally are isolated; compound dactylopore spines common. Dactylopore spines quite large and thick-walled, up to 3 mm in height and 1.5 mm in width, the dactylotome occupying one-fourth to one-third width of spine; exterior surface longitudinally ridged. Small, circular, flush secondary dactylopores, measuring only 0.08–0.11 mm in diameter, occur on walls of dactylopore spines, often several on each spine. Whereas dactylopore spines are quite large, dactylostyles are small and thin, only about 30–35 µm in width, bearing short spines up to 23 µm in length and 14 µm in diameter.




Figure 9. *Errinopora zarhyncha*, **A–I, K–L** male from USNM 1071915, **J** female from USNM 1123448: **A–C** branch surface showing tall dactylospine spines, gastropores, and ruptured male ampullae in figure C **D** stereo view of gastro- and dactylostyles and dactylospine spines **E–H** coenosteal texture **I** lateral view of a gastrostyle **J** female ampulla **K** dactylotome of a dactylospine spine **L** dactylostyle.

Gastropores circular and quite variable in size, up to 1.1 mm in diameter, often arranged linearly in valleys created by adjacent rows of dactylospines. Large gastropores often sit directly adjacent to much smaller ones; secondary gastropores about 0.38 mm in diameter. Gastropore fairly shallow, lacking a ring palisade, affording an easy view

of base of pore, as the gastrostyle occupies only a small part of gastropore cavity. Gastrostyles lanceolate and slender, up to 0.9 mm in height, and bear longitudinal anastomosing ridges, the gastrostyle size commensurate with diameter of gastropore tube.

Female ampullae inconspicuous (Fig. 9J) and not common, rarely seen in cross section branch fracture. Female ampullae hemispherical, often overshadowed by tall dactylopore spines or becoming incorporated into dactylopore spines. Efferent pores elusive; when detectable they are lateral in position, and about 0.7 mm in diameter, but more often a spent female ampulla lacks its upper half or is simply a crater in the coenosteum, a function of its thin, porous coenosteal cover. Male ampullae equally inconspicuous (Fig. 9C), roughly hemispherical, highly porous, and only about 0.6–0.7 mm in diameter.

Remarks. *Errinopora zarhyncha* is one of three species in the genus having a predominantly unilateral arrangement of dactylopore spines in which only one row of laterally fused spines (usually the proximal row) have their dactylotomes facing a gastropore or gastropore row, the other species being *E. nanneca* and *E. dichotoma* (see Dichotomous Key and Table 1). *Errinopora zarhyncha* differs from the other two species in having very tall dactylopore spines, relatively small dactylostyles, large robust colonies, and gastropore tubes that are much larger than their gastrostyles (see Dichotomous Key and Table 1). Naumov (1960) reported this species from the Kurile Islands, but based on his description in which he reports gastropore diameters of only 0.20–0.25 mm and dactylopore spines only 0.6 mm in height, we conclude that he is referring to a different, and as yet unknown species. Of the 8 lots of specimens examined, 4 are female, and 4 male. Coralla were determined to be calcitic by Thompson and Chow (1955) and Cairns and Macintyre (1992).

Distribution. Endemic to Aleutian Islands in a somewhat disjunct distribution: Amchitka Pass, Bowers Bank, and off Seguan Island; 207–658 m.

***Errinopora nanneca* Fisher, 1938**

http://species-id.net/wiki/Errinopora_nanneca

Figs 5B,G–I, 10A–K

Errinopora nanneca Fisher, 1938: 538–539, pl. 66, fig. 1, pl. 67, pl. 69, figs 2, 2a.—Boschma 1957: 57.—Naumov 1960: 559, fig. 404.—Cairns 1983b : 462.—Cairns et al. 1999: 43 (listed).—Cairns and Macintyre 1992: 100–101 (mineralogy).—Heifetz 2002: 22 (listed).—Lindner 2005: 87 (redescription, key), 103–106, figs 4.2E, 4.8, 4.15.—Heifetz et al. 2005: 133, 137 (listed).—Wing and Barnard 2005: 27 (listed).—Brooke and Stone 2007: 528, figs 2G, 3C.—Stone and Shotwell 2007: 107 (listed).—Jamieson et al. 2007: 224 (listed).—Lindner et al. 2008: 3, and supplemental Table S1: 3 (phylogeny and DNA sequences).

Errinopora nannacea: Cairns 1983a: 127 (incorrect spelling).

Errinopora nummea: Lowe 1967: 83 (incorrect spelling).

Errinopora pourtalesi: Wing and Barnard 2004: 58, fig. 28 (USNM 1123450).




Figure 10. *Errinopora nanneca*, **A–C** female holotype, **D–K** USNM 96252: **A–B** dactylospine spines on branch tips **C** stereo view of clustered female ampullae **D–G** lateral views of gastrostyles **H–I** coenosteal texture **J–K** dactylostyle.

Type material. Holotype: *Alb*-3599, a dry female colony 13 cm in height, USNM 42875 (Fig. 5B). Paratypes: *Alb*-3599, 6 colonies, including that figured by Fisher (1938: plate 67), 4 female, 1 male, 1 indet., all dry, USNM 52263. All other specimens mentioned by Fisher from *Alb*-3599 and 4777 are expressed excluded from type status according to ICZN (1999) article 72.4.6. **Type locality.** *Alb*-3599: 52°05'N, 177°40'E (off Kiska Island, Rat Islands, Aleutian Islands), 101 m, 9 Jun 1894.

Material examined. *Alb*-3480, 52°06'N, 171°45'W, 517 m, 8 Jul 1891, 1 indet., part of type series of *E. zarhyncha*, ex. USNM 52247; *Alb*-3599, type locality, 8 female,

8 male, USNM 52248; *Alb-4777*, 52°11'N, 179°49'E, 95 m, 5 Jun 1906, 11 female, 7 male, USNM 44070, 52249, 60299, and 62715; *Alaskan Leader* 35, 53°01'48"N, 170°06'12"W, 172–178 m, 4 Jun 2002, 1 female, 1 male, USNM 1123451 and 1123532; *Alaskan Leader* 54, 51°45'48"N, 179°06'08"E, 90–467 m, 11 Jun 2002, 1 female, USNM 1123378; *Delta*, 51°13'30"N, 179°53'17" E, 40 m, 16 Jul 2002, 1 indet. in alcohol, USNM 1123377; *Delta* 5599, 52°33.47'N, 179°26.44'E, 225 m, 15 Jul 2002, 1 male, USNM 1123452; *Delta* 5605, 51°50'55"N, 179°48'13"W, 125 m, 17 Jul 2002, 1 male, 4 indet., USNM 1123450 and 1123462; *Delta* 5607, 51°23'57"N, 179°48'35"W, 182 m, 16 Jul 2002, 1 female, USNM 1123459; *Delta* 5620, 51°57'40"N, 176°00'42"E, 150 m, 24 Jul 2002, 2 female with fragments in alcohol, USNM 1027820 and 1123510; *Delta* 5622, 51°57.5'N, 176°50.1'W, 140 m, 24 Jul 2002, 2 female, 1 male, and fragments in alcohol, USNM 1123468, 1123472, and 1123472; *Delta* 5625, 51°57.7'N, 176°50.2'W, 130 m, 25 Jul 2002, 1 female, USNM 1123473; *MF* 833–47, 51°55'36"N, 176°52'48"W, 201 m, 5 Aug 1983, 1 female and 1 male, USNM 77049; *MF* 833–49, 52°00.2'N, 176°21.4'W, 115 m, 5 Aug 1983, 1 male in alcohol, USNM 77031; *MF* 833–56, 52°02'06"N, 176°22'36"W, 249 m, 6 Aug 1983, 1 female, USNM 77044; *MF* 833–69, 52°02.8'N, 179°27.2'E, 63 m, 25 Aug 1983, 1 indet., in alcohol, USNM 77030; *Pacific Knight* 941–61, 52°17'N, 173°06'W, 0–143 m, 17 Jun 1994, 1 female, USNM 96254; *Sea Storm* 86, 51°36'21"N, 176°17'39"W, 345 m, 3 Jul 2002, 1 female in alcohol, USNM 1123475; *Sea Storm* 92, 51°33'34"N, 177°40'W, 367 m, 4 Jul 2002, 3 female, 3 male, USNM 1123529, 1123469–70, and 1123376; *Sea Storm* 100, 51°42'59"N, 175°47'07"E, 86 m, 7 Jul 2002, 1 indet., in alcohol, USNM 1123520; *Sea Storm* 105, 52°08'59"N, 175°06'47"E, 201 m, 8 Jul 2002, 1 indet. in alcohol, USNM 1123519; *Sea Storm* 108, 52°11'32"N, 175°17'E, 208 m, 8 Jul 2002, 1 male in alcohol, USNM 1123517; *Sea Storm* 111, 52°16'20"N, 175°59'13"E, 137 m, 9 Jul 2002, 1 indet. in alcohol, USNM 1123518; *Sea Storm* 112, 52°15'12"N, 176°00'42"E, 137 m, 9 Jul 2002, 1 indet. in alcohol, USNM 1123512; *Sea Storm* 114, 52°02'29"N, 177°39'E, 130 m, 10 Jul 2002, 1 male, USNM 1123515; *Sea Storm* 115, 52°04'16"N, 177°19'04"E, 165 m, 10 Jul 2002, 2 indet. in alcohol, USNM 1123474 and 1123513; *Sea Storm* 116, 52°04'10"N, 177°14.4'E, 87–94 m, 11 Jul 2002, 5 male, 1 indet., USNM 1123516; *Sea Storm* 118, 52°00'14"N, 177°19'04"E, 104–111 m, 11 Jul 2002, 2 female, 2 male, USNM 1123375 and 1123471; *Sea Storm* 122, 52°02'49"N, 179°25'18"E, 143 m, 13 Jul 2002, 1 indet., USNM 1123511; *Sea Storm* 126, 52°13'22"N, 179°49'26"W, 115 m, 13 Jul 2002, 1 female in alcohol, USNM 1123466; *Sea Storm* 128, 52°03'22"N, 179°48.13'W, 242 m, 14 Jul 2002, 1 female in alcohol, USNM 1123374; *Sea Storm* 157, 52°12'35"N, 172°12'20"W, 348 m, 23 Jul 2002, 1 female, USNM 1123467; *Vesteraalen* 941–50, 52°34'N, 170°40'W, 0–88 m, 13 Jun 1994, 3 female, 3 male, USNM 96252 and 1138188; *Vesteraalen* 941–59, 52°12'N, 172°12'W, 0–360 m, 14 Jun 1994, 3 male, USNM 96251 and 1123514; *Vesteraalen* 941–61, 52°05'N, 172°26'W, 0–156 m, 15 Jun 1994, 5 female, USNM 96253; *Vesteraalen* 941–151, 52°11'N, 179°44'E, 0–151 m, 10 Jul 1994, 1 indet., USNM 96257; *Vesteraalen* 941–153, 52°10'N, 179°43'E, 0–94 m, 10 Jul 1994, 1 female, 4 indet., USNM 96531;

Vesteraalen 941–154, 52°04'N, 179°47'E, 0–254 m, 11 Jul 1994, 1 male, USNM 96258; *Vesteraalen* 941–163, 51°37'N, 178°25'E, 0–155 m, 18 Jul 1994, 1 female, 1 male, USNM 96256; *Vesteraalen* 941–185, 52°04'N, 176°30'E, 0–91 m, 24 Jul 1994, 1 male, USNM 96255; *Vesteraalen* 3, 52°37'48"N, 169°47'16"W, 80 m, 21 May 2001, 1 female in alcohol, USNM 1123368; Gulf of Alaska, 219 m, 1 indet., USNM 77414; Renfro, coll., 52°02'17"N, 179°25'21"E, 236 m, 5 Apr 2000, 1 female, USNM 1123455; 52°18'N, 179°48'35"W, 490 m, 28 May 2000, 1 indet., USNM 1123457.

Description. Colonies quite variable in shape. Most colonies examined uniplanar, consisting of irregularly dichotomous, non-anastomotic branching (e.g., the holotype, Fig. 5B). The opposite extreme is multilobate and multiplanar colonies (Fig. 5H), composed of thin blades of corallum set in a three dimensional arrangement. Virtually all intergrades in colony shape were observed, including some having both large flattened lobes and slender branches (Fig. 5G). Tallest colony examined (USNM 96252) 21 cm in height with a basal branch diameter of 3.5 cm, but a broken colony having a basal diameter of 4 cm (USNM 1123516) implies an even larger size. Although distal branches often circular in cross section, more often they are somewhat compressed in branching plane. Parasitic spionid polychaete worms often form tubes along axis of branches, Fig. 8 in cross section. Coenosteum reticulate-spinose, the narrow strips only about 60 μm wide and bordered by slits of equal width, each strip bearing irregularly unilinearly arranged spines, altogether producing a porous or rough coenosteal texture. Coenosteum light orange to light pink.

Dactylopore spines isolated or arranged in transverse to oblique rows on distal branches, their dactylotomes facing upward (abcauline), their edges often fusing with edges of adjacent dactylopore spines. On more proximal branches and the basal branch, dactylopore spines fewer in number, and arranged in pseudocyclostyles, short rows, isolated, or as circles around small islands of 2–5 gastropores. Dactylopore spines strongly favor one face of corallum, and are much less common on opposite face. Dactylopore spines relatively small, only about 0.4 mm in maximum height and 0.30–0.35 mm in width, the dactylotome occupying middle third. Small (40–115 μm in diameter) secondary dactylopores flush with coenosteum common. Outer surface of dactylopore spines prominently ridged; inner surface bears a moderately robust dactylostyle (40–50 μm in width, Fig. 10J–K) composed of elements up to 18 μm tall and 7 μm in diameter.

Gastropores circular and flush with coenosteum, 0.15–0.44 mm in diameter, the average about 0.20 mm. Gastropore tube cylindrical, without a ring palisade. Gastrostyles lanceolate, up to 0.55 mm in height, bearing spinose longitudinal, sometimes anastomosing, ridges that themselves bear small spines up to 32 μm long and 8 μm in diameter. Smaller secondary gastropores, lacking gastrostyles, also present, these 0.11–0.19 mm in diameter.

Female ampullae (Fig. 10C) large hemispheres 1.1–1.8 mm in diameter, occurring fairly densely and equally on both corallum faces. Dactylopore spines often occur on female ampullae. Efferent pores rarely observed, but are lateral and up to 0.5 mm

in diameter. Male ampullae smaller mounds 0.4–0.7 mm in diameter, clustered, and somewhat irregular in shape. Both types of ampullae porous, like the coenosteum.

Remarks. *Errinopora nanneca* is one of three species in the genus having a predominantly unilateral arrangement of dactylopore spines in which only one row of laterally fused spines (usually proximal to gastropores) have their dactyloctomes facing a gastropore or gastropore row, the other species being *E. zarhyncha* and *E. dichotoma* (see Dichotomous Key and Table 1). *E. nanneca* is distinguished from *E. zarhyncha* in the account of the latter species, but differs from *E. dichotoma* in having shorter dactylopore spines and smaller gastropores, uniplanar colonies, and prominently ridged dactylopore spines. Of the 126 specimens examined, 58 are female, 42 male, and 26 indeterminate in gender. The corallum was found to be 100% calcitic according to Cairns and Macintyre (1992).

Distribution. Aleutian Islands from eastern Rat Islands to Islands of Four Mountains, including Petrel Bank; 40–517 m.

***Errinopora dichotoma* Lindner & Cairns, sp. n.**

urn:lsid:zoobank.org:act:7200AE40-A33D-4D86-A96C-1B6FC259A159

http://species-id.net/wiki/Errinopora_dichotoma

Figs 11D, 12A–M

Errinopora dichotoma Lindner, 2005: 93–96, figs 4.2H, 4.7A, 4.12 (unpublished name).

Type material. Holotype: *Dominator* 971–73, 1 alcohol-preserved male colony 11.5 cm in height, USNM 1123508 (Fig. 11D). Paratypes: *Alaskan Leader*, 53°01'42"N, 170°05.99'W, 200–400 m, 4 Jun 2000, 1 male, USNM 1123507; *Alaskan Leader* 35, 53°01'48"N, 170°06'12"W, 172–178 m, 4 Jun 2002, 1 female, 1 male, USNM 1137600; *Dominator* 971–73, topotypic, 1 dry branch, male, USNM 1148143. **Type locality.** *Dominator* 971–73, 52°33'10"N, 179°25'18"E (off northwestern Petrel Bank), 217 m, 26 Jun 1997.

Etymology. The specific name *dichotoma*, meaning “in two parts” (from the Greek *dicha*, meaning “in two” and *tomos*, meaning “part” or “slice”), referring to the dichotomous branching mode of this species.

Material examined. Types.

Description. Colonies three dimensional, robust, and sparsely branched, branching equally and widely dichotomously, resulting in broad U-shaped axils; branch anastomosis absent. Largest colony (holotype, Fig. 11D) 11.5 cm in height, with a basal branch diameter of 16 × 20 mm. Branches circular to elliptical in cross section, attenuating to thick (7–8 mm in diameter), blunt tips; tips of most branches missing (broken) from type material. As in most species of *Errinopora*, coenosteum quite porous (Fig. 12F–I), composed of a reticulum of thin, spinose strips separated by wide slits or series of pores, the spines being up to 55 μm in height and 15–24 μm in diameter. Coenosteum orange, branch cores being white.




Figure 11. Color figures of skeleton of various Alaskan stylasterids: **A, E, G** *Stylaster brochi* **B, F, H** *S. verrillii* **C** *S. stejnegeri*, **D** *Errinopora dichotoma*. **A** holotype, USNM 43264 **B** arborescent colony, USNM 1027819 **C** holotype, USNM 43271 **D** holotype, USNM 1123508 **E** large delicate sponge encrusted colony, USNM 1123304 **F** holotype of *A. moseleyi*, USNM 6851 **G** massive colony, USNM 1123006 **H** syntype, USNM 4193.




Figure 12. Holotype of *Errinopora dichotoma*, USNM 1123508: **A** branch cross section **B** colony surface showing gastro- and dactylopores **C** dactylopore with dactylostyle (upper) and secondary gastropore (lower) **D** gastropore **E, J** gastrostyles **F-I** coenosteal texture **K** gastrostyle spines **L-M** dactylostyle.

Dactylopore spines occur on all branch surfaces and quite variable in orientation (Fig. 12B), sometimes forming transverse or longitudinal rows of up to 7 laterally fused spines, spines sharing the same laterally fused row sometimes oriented in opposite directions, their dactylotomes facing in opposite directions. Isolated dactylopore spines also present. Dactylopore spines of moderate size and thick-walled, up to 1.2 mm in height and 1.1 mm in width, the dactylotome occupying one-fourth to one-third width of spine. Small (80 μ m diameter), circular secondary dactylopores common.

Dactylostyles robust (Fig. 12C, L–M), each about 0.125 mm in width, composed of elements up to 40 μm in height and 10–12 μm in diameter.

Gastropores circular, sometimes arranged in rows, otherwise randomly arranged, 0.3–0.5 mm in diameter; pores lack a ring palisade. Secondary gastropores (Fig. 12C, lower) 0.22–0.30 mm in diameter. Gastrostyle occupies most of gastropore cavity. Gastrostyles lanceolate, up to 0.5 mm in height, bearing longitudinal, anastomosing, spinose ridges.

Intact female ampullae never observed, but coenosteal depressions 1.0–1.1 mm in diameter are common in one specimen, these presumed to be spent female ampullae. Male ampullae small, porous hemispheres 0.5–0.6 mm in diameter.

Remarks. *Errinopora dichotoma* is one of three species in the genus having a predominantly unilateral arrangement of dactyloporous spines in which only one row of laterally fused spines (usually proximal to gastropores) have their dactyloporous faces facing a gastropore or gastropore row, the other species being *E. nanneca* and *E. zarhyncha*. *E. dichotoma* is compared to those species in their respective accounts and in the Dichotomous Key and Table 1. Of the five specimens examined, 1 was presumed to be female and 4 male.

Distribution. Endemic to Aleutian Islands: off Petrel Bank, off Islands of Four Mountains; 178–217 m (Cairns, 1992; Appeltans et al. 2011).

Genus *Stylaster* Gray, 1831

<http://species-id.net/wiki/Stylaster>

Diagnosis. Colonies branching in a flabellum or a bush shape (in only one case lamellate). Coenosteum usually reticulate-granular but may be linear-imbricate; coenosteum of many colors and hardness. Gastro- and dactyloporous arranged in conventional cycloporous arrangement, with only one gastrostyle per cycloporous; pores of the peripheral type; supernumerary dactyloporous often present. Cycloporous arranged on branch edges and/or on corallum faces, or uniformly on all branch surfaces, but not unifacially. Gastropore tube single-chambered, but may be constricted by a ring palisade. Dactylostyles present. Ampullae usually superficial.

Discussion. The genus and its various groupings are discussed by Cairns (1983b) and a key to all stylasterid genera, including the groups of *Stylaster*, is provided by Cairns (1992). Currently there are 80 recognized Recent species and 7 fossil species (Appeltans et al. 2011: WoRMS data base: www.marinespecies.org). It is by far the most species-rich and diverse genus of the stylasterids.

Type species. *Madrepora roseus* Pallas, 1766, by subsequent designation (Milne Edwards and Haime, 1850: xxii), a member of Group B.

Distribution. Oligocene to Recent: cosmopolitan from depths of 0–2010 m (Cairns, 1992; Appeltans et al. 2011).

***Stylaster* (Group A)**

Diagnosis. Species of *Stylaster* in which coralla form branching colonies, and in which cyclo systems are uniformly arranged on all sides of cylindrical, blunt branches.

Discussion. Previously called *Allopora*, these species were synonymized with *Stylaster* (see Cairns, 1983b) because no definitive break could be established between species having uniformly arranged and sympodially arranged cyclo systems.

***Stylaster brochi* (Fisher, 1938)**

http://species-id.net/wiki/Stylaster_brochi

Figs 11A, E, G, 13A–L

Allopora brochi Fisher, 1938: 517–518, pl. 42, figs 3–3d, pl. 44, pl. 45, fig 1.—Nau-mov 1960: 569–570, text-fig. 413.—Boschma 1957: 19.

?*Allopora abei* Eguchi, 1968: 34, pl. 34, figs 1–3.—Mori 1980: 17 (listed).

Stylaster brochi: Cairns 1983b: 429.—Heifetz 2002: 22 (listed).—Wing and Barnard 2004: 11, 27.—Heifetz et al. 2005: 133, 137 (listed).—Stone and Shorwell 2007: 108 (listed).—Brooke and Stone 2007: 521, figs 2A, 3A.—Jameison et al. 2007: 224 (listed).

?*Stylaster abei*.—Cairns 1983b: 430.

Stylaster polyorchis.—Lindner et al. 2008: 3, supplemental Table S1: 2.

Stylaster campylecus.—Lindner et al. 2008, 3, supplemental Table S1: 2.

Type material. *Allopora brochi*: Alb-4777: The dried holotype is deposited at the USNM (43264, also SEM stub 1497), measuring 9 cm in height (Fig. 11A); a para-type is also deposited at the CAS (28183). **Type locality.** *Albatross* 4777: 52°11'N, 179°49'E (Petrel Bank, Aleutian Islands), 79–95 m.

Allopora abei: Holotype, dry, Dept. of Mineral Sciences, Tohoku University (1001).

Type locality. Off the Aleutian Islands, exact locality and depth unknown.

Material examined. Holotype, USNM 43264; Alb-3599, 52°05'N, 177°40'E, 101 m, 9 Jun 1894, 1 female, USNM 76814 (part of type lot of *S. campylecus*); Alb-4779, 52°11'N, 179°57'W, 99–102 m, 5 Jun 1905, 1 female, USNM 44069; *Alaskan Leader* 35, 53°01'42"N, 170°05'59"W, 200–400 m, 4 Jun 2000, 1 male, USNM 1122516; *Delta* 5607, 51°23'02"N, 179°01'38"W, 126 m, 18 Jul 2002, 1 female, USNM 1122486; *Delta* 5620, 51°57'40"N, 176°50'01"W, 150 m, 24 Jul 2002, 5 female, 5 males, 5 indet., some in alcohol, USNM 1112304, 1123302, 1123304, 1123345–1123350; *Delta* 5622, 51°57'37"N, 176°49'58"W, 140 m, 24 Jun 2002, 2 indet., USNM 1123303; *Delta* 5624, 51°47'40"N, 176°49'58"W, 140 m, 25 Jun 2002, 1 male, USNM 1123300; *Delta* 5625, 51°57'42"N, 176°50'01"W, 156 m, 25 Jun 2002, 1 female, 1 indet., USNM 1027821; *Delta* 6000–10D-12, 51°50.828'N, 179°49.674'E, 112 m, 5 Jul 2003, 1 female, AB10–0002; *Dominator* 971–73, 52°33'10"N, 172°20'58"W, 217 m, 6 Jun 1997, 1 male in alcohol, USNM




Figure 13. *Stylander brochi*, **A, E, G–H, K** male from USNM 96264 **B–D, F, J** female syntype, USNM 43264 **I, L** male from USNM 1123049: **A** stereo view of cyclostyles on branch tip **B** stereo view of cyclostyles and supernumerary dactylopores **C** branch tip **D** a cyclostyle **E** coenosteal texture **F** dactylostyle **G** stereo view of longitudinal section through a cyclostyle **H** a gastrostyle surrounded by its ring palisade **I, L** an elongate gastrostyle **J** longitudinal section of a cyclostyle showing curvature of gastropore tube, sectioned female ampulla, and two gastrostyles **K** cross section through several male ampullae and a gastropore tube.

1123366; *Dominator* 971–135, 51°37'22"N, 178°35'E, 163 m, 14 Jul 1997, 1 female, USNM 1123364; *Let's Go*, 51°57'N, 178°11'E, 0–247 m, 30 Aug 1986, 3 males, USNM 96259; *MF* 801–70, 52°03'24"N, 179°25'06"E, 174 m, 29 Jul 1980, 1 female, USNM ; *MF* 833–47, 51°55'36"N, 176°52'48"W, 201 m, 5 Aug 1983, 1 male, USNM 77048; *Ocean Olympic*, 52°04'31"N, 177°12'28"E, 293 m, 4 Mar 2000, 1 male, USNM 1122540; *Pacific Knight* 941–36, 53°02'N, 170°13'W, 0–178 m, 11 Jun 1994, 8 male, USNM 96265, 96525, and 96542; *Pacific Knight* 941–37, 52°17'N, 170°40'W, 0–155 m, 12 Jun 1994, 1 male, USNM 96269; *Pacific Knight* 941–40, 52°41'N, 170°49'W, 0–126 m, 12 Jun 1994, 1 male, 1 indet., USNM 96270 and 96534; *Pacific Knight* 941–42, 52°55'N, 170°24'W, 0–225 m, 13 Jun 1994, 1 male, USNM 96522; *Pacific Knight* 941–49, 52°46'N, 171°45'W, 0–340 m, 14 Jun 1994, 1 female, USNM 96521; *Pacific Knight* 941–121, 51°38'N, 178°19'W, 0–375 m, 5 Jul 1994, 3 female, USNM 96262; *Pacific Knight* 941–204, 53°06'N, 171°42'E, 9–455 m, 31 Jul 1994, 1 female, USNM 96523; *Sea Storm* 90, 51°36'30"N, 177°10'48"W, 217 m, 4 Jul 2002, 1 male in alcohol, USNM 1123009; *Sea Storm* 91, 51°40'51"N, 177°10'49"W, 82 m, 4 Jul 2002, 1 male, USNM 1123021; *Sea Storm* 100, 51°42'59"N, 175°47'07"E, 86–94 m, 7 Jul 2002, 1 indet. in alcohol, USNM 1076928; *Sea Storm* 101, 51°45'27"N, 175°40'06"E, 83 m, 7 Jul 2002, 1 female, 2 males, USNM 1123024, -26, -27; *Sea Storm* 105, 52°08'59"N, 175°06'47"E, 201 m, 8 Jul 2002, 1 male, USNM 1123023; *Sea Storm* 111, 52°16'20"N, 175°29'13"E, 137 m, 9 Jul 2002, 1 male and 1 female in alcohol, USNM 1123000, 1123013; *Sea Storm* 113, 52°03'07"N, 177°23'55"E, 128 m, 10 Jul 2002, 1 male, USNM 1123007; *Sea Storm* 115, 52°04'16"N, 177°19'04"E, 165 m, 10 Jul 2002, 1 indet. in alcohol, USNM 1123020; *Sea Storm* 116, 52°04'10"N, 177°14'25"E, 87–94 m, 11 Jul 2002, 1 female, USNM 1123006; *Sea Storm* 118, 52°00'14"N, 177°49'40"E, 104–111 m, 11 Jul 2002, 1 female, USNM 1123283; *Sea Storm* 122, 52°02'49"N, 179°25'18"E, 143 m, 13 Jul 2002, 6 male colonies, some in alcohol, and SEM stub 1499, USNM 1123043–44, 1123046, 1123048–49; *Sea Storm* 124, 52°16'24"N, 179°55'47"E, 88 m, 13 Jul 2002, 1 male in alcohol; *Sea Storm* 129, 51°53'24"N, 179°44'07"E, 84–93 m, 14 Jul 2002, 1 male, USNM 1123032; *Sea Storm* 130, 52°12'10"N, 176°12'40"E, 86 m, 15 Jul 2002, 1 female, USNM 1123002; *Sea Storm* 146, 52°15'55"N, 174°48'02"W, 113 m, 20 Jul 2002, 1 male in alcohol, USNM 1123019; *Shishaldin*, 51°53.84'N, 177°44.88'W, 374 m, Feb 2000, 1 male, USNM 1122495; *Shishaldin*, 52°27'14"N, 179°30'54"E, 218 m, 2 Feb 2000, 1 female, USNM 1122521; *Vesteraalen* 3, 52°38'12"N, 169°46'35"W, 75 m, 21 May 2000, 1 female, USNM 1123373; *Vesteraalen* 5, 54°40'43"N, 169°06'11"W, 102 m, 21 May 2000, 1 female, 1 male, USNM 1123372; *Vesteraalen* 8, 53°09'28"N, 167°04'14"W, 154 m, 22 May 2000, 1 male in alcohol, USNM 1123371; *Vesteraalen* 941–36, 52°56'N, 169°31'W, 0–227 m, 10 Jun 1994, 1 male, USNM 96261; *Vesteraalen* 941–151, 52°10'N, 179°44'E (topotypic), 0–90 m, 10 Jun 1994, 1 female, USNM 96267; *Vesteraalen* 941–153, 52°10'N, 179°43'E (topotypic), 0–94 m, 10 Jun 1994, 2 female, USNM 96268; *Vesteraalen* 941–167, 51°54'N, 178°20'E, 0–150 m, 19 Jul 1994, 3 male, SEM stub 1498, USNM 96264;

Vesteraalen 941–185, 52°03'N, 176°31'E, 0–91 m, 24 Jul 1994, 1 female, USNM 96263; *Vesteraalen* 941–210, 53°07'N, 170°56'E, 0–92 m, 30 Jul 1994, 2 males, USNM 96266; University of Washington, 51°32'N, 179°15'W, 278–289 m, 1 Sep 1968, 1 female, USNM 96250; McClusky, coll., 51°53'14"N, 179°47'50"E, 351–393 m, 30 Mar 2000, 1 female, USNM 1122448; off Sharma, AK, depth unknown, 2 female, USNM 76539;

Description. Colonies variable in shape, ranging from planar (Fig. 11G) to multiplanar to bushy (Fig. 11A, E), the growth form possibly moderated by commensals (see Remarks). Branch tips usually blunt and circular to slightly flattened in cross section, 2.5–3.0 mm in diameter, more slender if affected by commensals; branch anastomosis not uncommon. Largest colony (USNM 96523) 28 cm in height; colonies attached by massive, dense basal branches up to 4.2 cm in diameter (USNM 96262). Coenosteum reticulate-granular in texture, coenosteal strips 50–60 μm wide, separated by thin slits about 10 μm wide; granules rounded, 7–8 μm in diameter. All colonies infested with spionid polychaetes (*Polydora*) and their characteristic binary axial tubes (see Remarks). Coenosteum pale orange.

Cyclosystems circular, 0.9–1.1 mm in diameter, and slightly (about 1 mm) raised above coenosteum; cyclosystems uniformly arranged on three or all four sides of a branch, rarely in a linear fashion. Gastropores 0.35–0.40 mm in diameter. Gastropore tubes cylindrical, curved (Fig. 13J), and long (often 2–3 times length of gastrostyle), the length and curvature of tube making it difficult to see gastrostyle tip when viewed from above. Ring palisade absent or diffuse, the elements about 50 μm in height and diameter. Gastrostyles lanceolate and up to 0.7 mm in height, bearing long spines up to 70 μm in length.

Dactylotomes 80–110 μm in width; dactylostyles poorly developed, the cylindrical elements about 20 μm in height and 8 μm in diameter. Range of dactylopores per cyclosystem 6–13 ($n = 50$, average = 9.22 ($\sigma = 1.43$), and mode = 10). Supernumerary dactylopores quite common on coenosteum (Fig. 13A–C) and even on pseudosepta, presenting as small (0.09–0.11 mm in diameter), circular, slightly raised (rimmed) pores. Pseudosepta variable in width (Fig. 13D), in the same cyclosystem varying between 0.11 and 0.30 mm; adcauline diastemas also frequently present, each about twice pseudoseptal width.

Female ampullae (Fig. 13J) superficial hemispheres 0.8–1.1 mm in diameter; efferent pores rarely expressed, but if present, occurring in lateral position and somewhat recessed into coenosteum, each about 0.25 mm in diameter. Male ampullae (Fig. 13K) primarily internal, visible on coenosteal surface only as low mounds 0.3–0.5 mm in diameter, often with a small (30 μm diameter) apical efferent pore. Male ampullae often occur in high density clusters, directly adjacent to one another.

Remarks. Virtually every colony of *Stylaster brochi* examined contained commensal spionid polychaetes of the genus *Polydora* (K. Fauchald, per. comm.), such that this character was informally used to distinguish it from other *Stylaster* species, such as *S. campylecus*. Almost every branch is bored axially by this robust worm, which forms binary longitudinal tubes 0.7–0.8 mm in diameter that are figure

8-shaped in cross section (Fig. 15K). Several worms may occur in the same colony, each having one or more efferent openings somewhere on the colony surface. It is easy to conjecture an advantage to the worm in this relationship, i.e. a secure place to live and protection from predators, but it is difficult to imagine an advantage to the stylasterid, as the worm must weaken the strength of the branches as well as compete for the same filtered plankton in the water. Thus, it would seem that these polychaetes are parasites on the coral. Large colonies of *S. brochi* also form substantial three dimensional habitats for a variety of invertebrates, including sponges, bryozoans, hydroids, barnacles, bivalves, and ophiuroids. Heavy encrustation by sponges seems to promote a more delicate colony growth form, in which the terminal branches are only 1.5 mm in diameter (Fig. 11E). In a monograph otherwise devoted to stylasterids from Sagami Bay, Japan, Eguchi (1968) inexplicably described *Allopora abei* from “the Aleutian Islands.” Although the type was not examined, and Cairns (1983b) suggested a synonymy with *Stylaster polyorchis* Fisher, 1938, the illustrations provided by Eguchi suggest a synonymy with *S. brochi*. Of 86 colonies examined, 35 are female, 40 male, and 11 indeterminate, resulting in a fairly equal sex ratio. *S. brochi* is one of the most common and one of the largest growing stylasterids in the Aleutian Islands. It can be distinguished from other Alaskan species in *Stylaster* (Group A) by its usually planar growth mode, higher number of dactylopores per cyclosystem, and variable width of its pseudosepta (see Table 2 for additional comparisons).

Distribution. Widespread throughout Aleutian Islands from west of Attu Island to Unalaska (including Petrel Bank), with one disjunct record near Sharma (near Anchorage); 75–351 m, but most records between 100–200 m.

***Stylaster stejneri* (Fisher, 1938)**

http://species-id.net/wiki/Stylaster_stejneri

Figs 11C, 14A–J

Allopora stejneri Fisher, 1938: 518–519, pl. 42, figs 2–2b, pl. 56.—Boschma 1957: 28. *Stylaster stejneri*: Cairns 1983b: 429.—Wing and Barnard 2004: 11, 28.—Heifetz et al. 2005: 134, 137 (listed).—Stone and Shotwell 2007: 108 (listed).—Jameison et al. 2007: 224 (listed).

Type material. Holotype, 1 female colony, and SEM stub 1500, dry, USNM 43271 (Fig. 11C). **Type locality.** *Albatross* 4777: 52°11'N, 179°49'E (Petrel Bank, Aleutian Islands), 79–95 m.

Material examined. Holotype.

Description. The holotype (Fig. 11C) is a small, arborescent colony 6 cm in height and 7 cm in width, with a basal branch diameter of 9.3 mm. Branches cylindrical and blunt tipped, the tips measuring 3–5 mm in diameter. Coenosteum reticulate-granular in texture, coenosteal strips 50–55 µm wide, separated by slits 13–15 µm wide. Coe-

Table 2. Tabular Key of the Northeast Pacific Species of *Sylaster* (AI= Aleutian Islands, BB=Bowers Bank, PB=Petrel Bank, GOA=Gulf of Alaska, CS=Cyclosystem, PS=Pseudosepta, GS=gastrostyle, DP=Dactylopoire)

	<i>S. brochi</i>	<i>S. stejnegeri</i>	<i>S. verrilli</i>	<i>S. repandus</i>	<i>S. venustus</i>	<i>S. californicus</i>	<i>S. campylecus</i>	<i>S. leptostylus</i>	<i>S. trachystomus</i>	<i>S. parageus parageus</i>	<i>S. parageus columbiansis</i>	<i>S. alaskanus</i>	<i>S. elassotomus</i>
Colony Shape; Branching	Planar to bushy; Branch tips blunt	Arborescent; Branch tips blunt	Arborescent; Branch tips blunt	Lamellar (solid)	Planar to slightly bushy; Slender, blunt	Arborescent, large; Blunt tipped	Planar; Delicate	Planar, dichotomous; Delicate	Planar; Delicate branches	Bushy (multi-flabellate); Very delicate	Bushy; Delicate	Planar; Branches forming sieve-like reticulum	Bushy; Delicate
Coenosteal Color	Orange	Orange, Pink	Light orange	Light orange, pink	Violet, Pink	Rose, red, purple	White, orange, pink	White	Pale orange, pink	White	White	Orange, pink, white	White
Spionid Worm Tubes	Present	Present	Present	Present	Present	Present	Absent	Absent	Common	Common	Absent	Absent	Absent
Arrangement of Cyclosystems	All sides of branch	All sides of branch	All sides of branches	Transverse rows	All sides of branches	All sides of branches	Branch edges and anterior face	Branch edges and anterior face	Branch edges and anterior face	Branch edges and anterior face	Branch edges and anterior face	Only on branch edges	Only on branch edges
Cyclosystems: Shape and Size	Circular; 0.9-1.1 mm	Circular; 1.0-1.2 mm	Circular; 1.0-1.2 mm	Circular; 1.0-1.15 mm	Polygonal, raised; 0.7-0.9 mm	Circular or irregular; 0.7-1.0 mm	Variable, slightly flared; 1.0-1.3 mm	Circular; 1.0-1.1 mm	Asymmetrical, slightly curved; Not visible	Circular; 0.9-1.0 mm	Circular to elliptical; 1.1-1.5 mm	Circular to irregular; 0.9-1.3 mm	Circular to elliptical; 1.0-1.2 mm
Gastropore Tube; Shape; Visibility of GS Tip	Cylindrical; Difficult to see	Cylindrical; curved; Not visible	Straight upper; cylindrical (lower); Visible	Funnel-shaped; straight; Visible	Straight rounded upper; cylindrical (lower); Visible	Straight, funnel-shaped; Visible	Cylindrical; Difficult	Funnel-shaped above; cylindrical below; Visible	Funnel-shaped above, cylindrical below; Visible	Cylindrical; curved; visible	Funnel-shaped upper; cylindrical lower; Visible	Cylindrical; straight; Easily visible	Cylindrical, cal, curved; Rarely seen
Ring Palsade	Absent or rudimentary	Absent	Robust	Rudimentary	Robust	Present	Absent to rudimentary	Absent	Rudimentary	Well-developed	Well-developed	Well-developed	Rudimentary

	<i>S. brochi</i>	<i>S. stejnegeri</i>	<i>S. verrillii</i>	<i>S. repandus</i>	<i>S. venustus</i>	<i>S. californicus</i>	<i>S. campylactus</i>	<i>S. leptostylus</i>	<i>S. trachystomus</i>	<i>S. parageus parageus</i>	<i>S. parageus columbiensis</i>	<i>S. adaskanus</i>	<i>S. elastosomus</i>
Dactylopores/Cylosystem; Supernumerary Dactylopores	6-13 (x=9.22); Common	5-11 (x=6.46); Common	5-10 (x=7.10); Absent	1-11 (x=3.94); Absent	4-8 (x=5.70); Absent	3-8 (x=5.52); Common	7-17 (x=11.94); Absent	7-12 (x=9.76); Absent	8-18 (x=11.82); Rare	5-11 (x=8.54); Rare	6-13 (x=9.38); Rare	7-14 (x=11.30); Absent	11-17 (x=14.40); Common
Dactylostyles	Rudimentary	Rudimentary	Robust	Robust	Robust	Robust	Rudimentary	Moderate	Rudimentary	Well-developed	Well-developed	Rudimentary	Rudimentary
Ampullae:	Superficial;	Superficial,	Both	Superficial;	Both	Both	Both	Both	Superficial,	Superficial;	Both super-	Superficial,	Unknown;
Female:	Partially	ridged;	internal	Primarily	internal	internal	superficial	superficial	Superficial;	Partially	facial	ridged;	Superficial
Male:	internal	Unknown	internal	internal	internal	internal	superficial	superficial	Superficial	internal	superficial	Superficial	Superficial
Other Characters	Pseudosepta of variable width	Coenosteal papillae common	Coenosteal papillae common	Coenosteal papillae common	Coenosteal papillae common; Css occasionally linked	Coenosteal papillae common; Css often linked	Coenosteal strips linear near css	Ps porous and wide; Coenosteal ridges	Ps porous and wide; Coenosteal ridges	Gastropores only 0.25-0.30 mm in diameter	Gastropores 0.45-0.50 mm in diameter	Coenosteal papillae occasionally	Dactylostyles very shallow




Figure 14. Holotype of *Styaster stejnegeri*, USNM 43271: **A** stereo view of branch with cyclostyles, female ampullae, and supernumerary dactylopores **B** stereo view of a cyclostyle **C–E** cyclostyles with adjacent supernumerary dactylopores **F** coenosteal texture **G** longitudinal section of a gastropore tube **H** dactylostyles **I–J** ridged female ampullae.

nosteum also covered with small papillae (nematopores). Binary spionid worm tubes present along branch axes. Coenosteum light orange to pink.

Cyclostyles on all sides of branches, circular, 1.0–1.2 mm in diameter. Gastropores circular, 0.35–0.45 mm in diameter. Gastropore tube cylindrical and usually slightly curved such that gastrostyle tip is not visible from apical view; ring palisade absent. Gastrostyle lanceolate.

Cyclostyles flush to only slightly raised above coenosteum; surface of pseudosepta porous. Dactylotome width 0.08–0.11 m. Range of dactylopores per cyclostyles 5–11 ($n = 50$, average = 6.46 ($\sigma = 1.03$), and mode = 6). Small (0.11–0.14 mm diam-

eter), circular supernumerary dactylopores fairly common (Fig. 14A–C). Dactylostyles rudimentary (Fig. 14H).

Female ampullae (Fig. 14A, I–J) superficial hemispheres up to 1.1 mm in diameter, often bearing low ridges radiating from their apex. Male ampullae unknown.

Remarks. Although only one specimen is known of this species, it does appear to be distinctive. Perhaps most similar to *S. brochi*, it differs from that species in having flush cyclo systems, fewer dactylopores per cyclo system, and ridged female ampullae (Table 2).

Distribution. Known only from one specimen from the type locality.

***Stylaster verrillii* (Dall, 1884)**

http://species-id.net/wiki/Stylaster_verrillii

Figs 11B, F, H, 15A–M

Allopora verrillii Dall, 1884: 111–113.—Fisher, 1931: 391–392.

Allopora moseleyi Dall, 1884: 113.—Fisher, 1931: 391–392.

Allopora verrilli: Fisher 1938: 521–522, pl. 54, fig. 3, pl. 57, pl. 76, figs 5–6.—Nau-mov 1960: 567, text-fig. 411.

Stylaster (Allopora) norvegicus forma *pacifica* Broch, 1936: 52, text fig. 15c-d (in part: record from Snake Island, Strait of Georgia, British Columbia).

Stylaster (Allopora) verrillii: Broch 1942: 6.

Allopora norvegicus pacifica.—Lowenstam 1964: 382 (mineralogy).

Stylaster verrillii: Cairns 1983b: 429.—Heifetz 2002: 22 (listed).—Wing and Barnard 2004: 11, 28, fig. 31.—Heifetz et al. 2005: 134, 137.—Stone and Shotwell 2007: 108 (listed).—Brooke and Stone 2007: 522, fig. 2B.—Jameison et al. 2007: 224 (listed).—Lindner et al. 2008: 3, supplemental Table S1 (phylogeny and DNA sequences).

Stylaster ?norvegicus.—Jameison et al. 2007: 224 (listed).

Type material. *Allopora verrillii*: 5 badly worn, dry, female syntypes, the largest 44 mm in diameter (USNM 4193, Fig. 11H). All branches are eroded from the colony, thus resembling an encrusting colony. Collected together with four dry topotypic specimens (USNM 8850).

Type locality. Chika Island, Akutan Pass, near Unalaska, Aleutian Islands, on beach.

Allopora moseleyi: Holotype, 1 small (18 mm) dry colony, USNM 6851 (Fig. 11F).

Type locality. Kiska Harbor, Kiska Island, Aleutian Islands, on beach.

Material examined. Types of the two species; *Alb*-4777, 52°11'N, 179°49'E, 95 m, 5 June 1906, 1 female, 1 male, 1 indet., and SEM stub 1501, USNM 53394 and 76524 (mentioned by Fisher, 1938); unnumbered *Albatross* station, Sucia Islands, near San Juan Island, Washington, 16 Sep 1890, 2 female, 11 male colonies, and SEM stub 1513, USNM 76526–28 (mentioned by Fisher, 1938); *Delta* 5622, 51°57'36.54"N, 176°49'58.32"W, 140 m, 24 Jul 2002, 1 male, and SEM




Figure 15. *Styaster verrillii*, **A, F, L–M** male from USNM 76524, **B, G–J** male from USNM 1123517, **C–E** USNM 1123299, **K** female from USNM 76528: **A** stereo view of blunt branch tip showing cyclostyles and spionid tubes **B–C** cyclostyles **D–E** coenosteal texture **F, M** dactylostyle **G–J** lateral view of gastrostyles (**H** a stereo view) **K** stereo view of axial spionid tubes and cross section of female ampullae **L** male ampulla.

stub 1503, USNM 1123299; *Delta* 5625, 51°57'42"N, 176°50'01"W, 155 m, 25 Jul 2002, 1 male, USNM 1027819; *Patricia Lee*, 51°53.44'N, 179°47.7'E, 298 m, 1 indet., AB00–41b; *Sea Storm* 122, 52°02'49"N, 179°25'18"E, 143 m, 13 Jul 2002, 2 female colonies in alcohol, USNM 1123058 (attached to *Distichopora borealis*, USNM 1123061); *Sea Storm* 123, 52°10'53"N, 179°37'02"E, 124 m, 13 Jul 2002, 3 female, 2 male, 1 indet., USNM 1123053, 1123056; *Sea Storm* 129, 51°52'24"N, 179°44'07"E, 84–93 m, 14 Jul 2002, 1 male in alcohol, USNM 1123051; *Sea Storm* 155, 52°38'43"N, 172°27'27"W, 393 m, 22 Jul 2002, 1 female, USNM 1122763; *Vesteraalen* 941–40, 52°53'N, 169°59'W, 0–62 m, 11 Jun 1994, 1 female, USNM 96260; *Vesteraalen* 941–151, 52°10'36"N, 179°43.7'E, 87–92 m, 10 Jul 1994, 1 male, and SEM stub 1502, USNM 1123537; south of Semisopochnoi Islands, 366 m, 1 indet., USNM 88368; Ralston Island, near Juneau, depth unknown, 1 indet., AB04–21; Stubbs Island, BC, depth unknown, 2 indet., AB02–02; Wooden Island, AK, depth unknown, 2002, 1 dry, AB02–150b; Middle Cross Sound, AK, 15–21 m, 27 Jul 1978, 1 indet., AB78–120a; North Pass, Lincoln Island, AK, depth unknown, 2 indet. in alcohol, AB05–37.

Description. Colonies arborescent, dichotomously branching to form a three dimensional bush; branch tips blunt to slightly clavate, 3.0–4.5 mm in diameter. Largest colony (USNM 1027819, Fig. 11B) 3.5 cm tall, 7.0 cm wide, and attached by a basal branch 11 mm in diameter. Colonies usually attached to stones or bivalve shells. Coenosteum reticulate-granular in texture, coenosteal strips 52–55 μ m wide, separated by thin slits 10–13 μ m wide. Strips covered with small (8–10 μ m in diameter) spines, occurring 4–5 across width of a strip, conferring a fine, granular or "sugary" texture. Coenosteal papillae common. Most colonies infested with spionid worms and their characteristic binary tubes (Fig. 15A, K). Coenosteum light orange.

Cyclosystems occur on all sides of branches, circular, and 1.0–1.2 mm in diameter. Gastropore circular and small (0.25–0.30 mm in diameter). Gastropore tubes funnel-shaped near surface but cylindrical adjacent to gastrostyle, straight, and often quite long in thick branches. Ring palisade robust near gastrostyle tip, composed of squat, clavate elements about 40 μ m in diameter. Gastrostyles lanceolate to elongate (up to 1.5 mm, Fig. 15H), with a pointed tip; gastrostyles covered with anastomosing, longitudinal, spiny ridges, the spines quite long (up to 65 μ m in length) and sharp.

Cyclosystems flush to only slightly raised above coenosteum. Dactylotomes 0.07–0.10 mm in width, sometimes becoming obsolete within a system by infilling of the axial slit, resulting in just an apical pore. Dactylostyles robust (Fig. 15F, M), composed of tall slender cylindrical elements up to 60 μ m in height and 10–15 μ m in diameter. Range of dactylopores per cyclosystem 5–10 ($n = 50$, average = 7.10 ($\sigma = 1.16$), and mode = 7). Pseudosepta triangular; diastemas rare.

Female ampullae primarily internal (Fig. 15K), with only a slight superficial swelling; internal diameter of ampullae 0.7–0.8 mm. Efferent pores of female ampullae




Figure 16. *Stylander venustus*, **A, E–I** USNM 76545, **B–D** holotype, MCZ 5525: **A** blunt branch tip **B–D** cyclosystems (**C** a stereo view) **E** coenosteal texture **F** stereo view of a gastrostyle and ring palisade **G** two dactylostyles **H** cross section of two internal male ampullae **I** stereo view of branch cross section showing many internal male ampullae.

never observed. Male ampullae (Fig. 15H, L) also primarily internal, visible on surface only as small dimples about 0.4 mm in diameter; internal cavity diameter 0.3–0.5 mm.

Remarks. *Stylander verrillii* is similar to *Stylander venustus* (Figs 16A–I, 17F–G), but differs in having orange, bushy coralla (not pink to purple planar colonies, Fig. 17G); larger cyclosystems (1.0–1.2 mm vs 0.7–0.9 mm in diameter); larger diameter and blunt distal branches; triangular (not rectangular, Fig. 16B) pseudosepta; and straight-sided gastropore tubes (not hemispherical, Fig. 16B)(also see Table 2). Both species are confined to relatively shallow water and have overlapping distributions off British

Columbia and Washington. According to the ICZN (1999) article 31.1.3, the original spelling *verillii* should be preserved in preference to *verilli*.

Distribution. Known from Kiska Harbor, Aleutian Islands to Sucia Islands, Washington; 21–393 m, although most records between 60–155 m.

***Stylaster repandus* Lindner & Cairns, sp. n.**

urn:lsid:zoobank.org:act:255EF5E7-5BFA-428A-A5B9-CC47C1A28385

http://species-id.net/wiki/Stylaster_repandus

Figs 27A–C, 28A–M

Type material. Holotype: 1 large dry male colony, and SEM stub 1504, USNM 1122740 (Fig. 27A–B). Paratypes: 52°21.20'N, 171°02.77'W, 475 m, 1 Sep 2000, 1 dry female colony, and SEM stub 1505, USNM 1122739; Amukta Pass, depth unknown, 1996, 2 dry male colonies, USNM 1122741. **Type locality.** 52°17.86'N, 170°58.28'W (southeast of Amukta Island), 375 m, 12 Sep 2000.

Etymology. The specific name *repandus* (from the Latin, meaning “bent backwards”, “undulate”) refers to the complexly folded coenosteal lamellae of this species.

Material examined. Types.

Description. Colonies firmly attached to a hard substrate by a robust basal stem and encrusting base, the base being 9 cm in diameter and the basal stem 5 cm in diameter in the case of the holotype. Immediately above basal stem the colony divides into two or three lamellae (Fig. 27C), each lamella (or sheet) of coenosteum increasing its surface area by folding its surfaces into a complex three-dimensional structure similar to that of *Cyclobelia*. Short (up to 15 mm) cylindrical branches project from plane of colony, but always in response to housing a parasitic spionid worm tube. Holotype (Fig. 27A–B) 23 cm tall and 19 cm in width. Coenosteal lamellae thin at edge, only 2.0–2.5 mm thick, and thus easily damaged during collection; basal lamellae up to 10 mm in thickness. All colonies examined heavily infested with spionid worm tubes. Coenosteum reticulate-granular in texture, the strips being about 30 µm wide, bordered by slits about 9–19 µm thick; granules spinose. Coenosteum covered uniformly with small papillae (Fig. 28B). Coenosteum light orange, but central core a light shade of pink.

Cyclosystems equally common on both faces of lamellae, and even occur on lamellar edges, often arranged in transverse rows of up to 15 cyclosystems that parallel lamellar edge; diameter of cyclosystems 1.0–1.15 mm. Gastropore tube funnel-shaped near branch surface, and straight but constricted, the upper diameter being about 0.34 mm in diameter, the constriction corresponding to a diffuse ring palisade, the blunt elements being up to 60 µm tall and 40 µm in diameter. Gastrostyles lanceolate, pointed, and bear longitudinal, spinose ridges. Illustrated style (Fig. 28D) 0.53 mm in height and 0.26 mm in basal diameter.

Cyclosystems almost flush with coenosteum, the dactyloporite slits raised only slightly above coenosteum; dactyloporites 0.15–0.17 mm in width. Based on 74 cyclosystems the range of dactyloporites per cyclosystem is 1–11 (average = 3.94 ($\sigma = 2.05$),

and mode = 4). In many cyclosystems there is an adcauline diastema (Fig. 28E), often produced by the infilling of 1–3 adcauline dactylopores, these dactylopores becoming obsolete. Dactylostyles robust, composed of an almost unilinear arrangement of cylindrical, blunt to clavate elements up to 60 μm tall and 13–15 μm in diameter (Fig. 28D, H, J–K).

Female ampullae (Fig. 28L–M) low swellings on coenosteum 0.9–1.1 mm in diameter (internal diameter 0.7–0.8 mm), each with a peripheral efferent pore 0.3–0.4 mm in diameter that faces upward. Male ampullae (Fig. 28I) densely arranged, superficially visible only as inconspicuous mounds 0.5–0.6 mm in diameter (internal diameter 0.4–0.5 mm), often with an irregularly shaped apical efferent pore about 60 μm in diameter.

Remarks. Among the 80 Recent species in the genus *Stylaster* (Appeltans et al. 2011: www.marinespecies.org), *S. repandus* is the only one to have a lamellate growth form. It is very similar in growth form to *Cyclohelia lamellata* and *Errinopora undulata*, and is one of five stylasterid species to have lamellate colonies (see Discussion of *Errinopora undulata*).

Distribution. Known only from three localities in the vicinity of Amukta Island, Aleutian Islands; 375–475 m.

Stylaster (Group B)

Diagnosis. Species of *Stylaster* in which coralla form branching colonies, and in which cyclosystems are arranged primarily on branch edges, but also on anterior and occasionally posterior sides of branches; branches usually delicate.

Discussion. Called the “annectant” group by Cairns (1983b) because it links the morphology of Groups A and C regarding cyclosystem arrangement.

Stylaster campylecus (Fisher, 1938)

http://species-id.net/wiki/Stylaster_campylecus

Figs 17A–D, 18A–M

Allopora campyleca Fisher, 1938: 505–506, pls. 34, 36.—Broch 1942: 101.—Boschma 1957: 19–20.—Not Lowenstam 1964: 382 (probably *S. parageus parageus*).

Allopora polyorchis Fisher, 1938: 503–505, pl. 35, fig. 1, pl. 37, 38.—Naumov 1960: 581.—Boschma 1957: 27.

Allopora campyleca tyloata Fisher, 1938: 509–510, pl. 41, figs 2–2e.—Naumov 1960: 580.

Allopora moseleyana Fisher, 1938: 512–514, pl. 49, fig. 2, pl. 51 (not pl. 50), pl. 53, fig. 1 (not forma *leptostyla*).—Boschma 1957: 23.—?Naumov 1960: 577.

?*Allopora campyleca campyleca*.—Naumov 1960: 578.

- Stylaster campylecus*.—Cairns 1983b: 430.—Not Cairns and Macintyre 1992: 100 (probably *S. parageus parageus*).—Brooke and Stone 2007: 523524, fig. 2C.—Not Lindner et al. 2008: 3, Table S1 (= *S. brochi*).
- Stylaster moseleyanus*.—Cairns 1983b: 429.—Wing and Barnard 2004: 11, 28, fig. 30.—Heifetz et al. 2005: 124, 137 (listed).—Stone and Shotwell 2007: 108 (listed).—Jamieson et al. 2007: 224 (listed).
- Stylaster polyorchis*.—Cairns 1983b: 430.—Cairns and Macintyre 1992: 100–101 (mineralogy).—Heifetz 2002: 22 (listed).—Wing and Barnard 2004: 11, 28.—Heifetz et al. 2005: 134, 137 (listed).—Stone and Shotwell 2007: 108 (listed).—Jamieson et al. 2007: 224 (listed).—Not Lindner et al. 2008: 3, Table S1 (= *S. brochi*).
- Stylaster campylecus tylotus*.—Cairns 1983b: 430.—Wing and Barnard 2004: 27 (listed).—Heifetz et al. 2005: 134 (listed).—Stone and Shotwell 2007: 108 (listed)—Jamieson et al. 2007: 224 (listed).
- Stylaster campylecus campylecus*.—Wing and Barnard 2004: 11, 27, fig. 32.—Heifetz et al. 2005: 134 (listed).—Stone and Shotwell 2007: 108 (listed)—Jamieson et al. 2007: 224 (listed).

Type material. *Allopora campyleca*: Holotype: *Alb*-3480, 1 dry male colony, now in two pieces, length 16 cm, and SEM stub 1524–25, 1545, USNM 42870 (Fig. 17A). Paratypes: *Alb*-3480, 7 dry colonies and 7 in alcohol, and SEM stub 1526, USNM 43767, 52262, 52265, 58099, and CAS 28297; *Alb*-2852 (missing in 2011); *Alb*-2858, 3 dry colonies (not considered conspecific with *S. campyleca*); *Alb*-3599, 2 dry colonies (1 male is *S. campylecus*, 1 female is *S. brochi*), USNM 76814; *Alb*-4230, 1 dry male colony (not considered to be conspecific with *campyleca*); *Alb*-4302, 2 female dry colonies, USNM 76816. **Type locality.** *Alb*-3480: 52°06'N, 171°45'W (Amukta Pass, Aleutian Islands), 518 m.

Allopora polyorchis: Holotype: *Alb*-3480, 1 large dry colony now in two pieces, and SEM stubs 1527–28, USNM 43266 (Fig. 17B). Paratypes: *Alb*-3480: about 15 dry branch fragments, USNM 76540–76542, CAS 28293. **Type locality.** *Alb*-3480: 52°06'N, 171°45'W (Amukta Pass, Aleutian Islands), 518 m.

Allopora campyleca tyloca: Syntypes: *Alb*-4781, 4 dry female colonies, the largest 15 cm in height, and SEM stub 1539, USNM 43263, and 12 dry male branches, and SEM stub 1540, USNM 86000 (Fig. 17C). **Type locality.** *Alb*-4781: 52°14'30"N, 174°13'E (off Agattu I., Near Islands, Aleutians), 882 m.

Allopora moseleyana: Holotype: *Alb*-4781: 1 dry male colony now in two pieces, largest 13 cm in length, USNM 42869 (Fig. 17D). Paratypes: *Alb*-4781, 1 dry male colony, USNM 86003; *Alb*-3480, 3 female and 1 male dry colonies, and SEM stub 1519, USNM 76556, 76557; *Alb*-3480, 2 dry female and 1 male colonies (considered as *S. leptostylus*), USNM 76555. **Type locality.** *Alb*-4781: 52°14'30"N, 174°13'E (off Agattu I., Near Islands, Aleutians), 882 m.

Material examined. Types of *A. campyleca*, *A. polyorchis*, *A. campyleca tyloca*, and *A. moseleyana*; *Alb*-3480, numerous branch fragments (topotypic), USNM 76565;




Figure 17. Color figures of skeleton of various Alaskan stylasterids: **A–D** *Stylaster campylecus* **E** *S. leptostylus* **F–G** *S. venustus* **H** *S. trachystomus* **A** holotype, USNM 42870 **B** holotype of *Allopora polyorchis*, USNM 43266 **C** syntype of *A. c. tyloa*, USNM 86000 **D** holotype of *Allopora moseleyanus*, USNM 42869 **E** holotype of *A. moseleyanus leptostylus*, USNM 43270 **F** holotype, MCZ 5525 **G** three branches, USNM 76545 **H** holotype, USNM 43265.

Alaskan Leader-35, 53°01'42"N, 170°05'59"W, 200–400 m, 4 Jun 2000, 1 female, USNM 1122510; *Alaskan Leader* 64, 53°07'N, 166°53'42"W, 324–766 m, 21 Jun 2002, 1 female, USNM 1122484; *Alb*-4781, type locality, 1 female, USNM 76700; *Ballyhoo*, 54°47'20"N, 178°46'53"E, 254 m, 5 Jun 2000, 1 female, USNM 1122535; *Ballyhoo*, 54°42.44'N, 178°46.83'E, 250 m, 5 Jun 2000, 1 female, AB00–0013; *Delta*, 51°50'55"N, 179°49'26"W, 125 m, 17 Jul 2002, 1 indet., USNM 1123463; *Delta Alfa*, western Aleutian Islands, depth unknown, 1961, 1 female, USNM 52447; *Dominator* 971–181, 51°27'43"N, 176°38'20"E, 384 m, 27 Jul 1997, 2 female, 2 male, USNM 1123363; *Jason* 2095–7-7, 51°48.693'N, 173°49.965'W, 843 m, 25 Jul 2004, 1 male, AB09–0013; *Jason* 2102–7-8, 51°30.56'N, 177°55.36'W, 824 m, 2 Aug 2004, 6 male, AB09–0012; *Jason* 2101–9-1, 51°31.43'N, 177°57.5'W, 494 m, 2 Aug 2004, 1 male, AB09–0015; *Jason* 2103–15–1, 51°50.968'N, 179°51.007'E, 678 m, 1 male, AB08–0037; *Jason* 2104–1-5, 51°43.83'N, 179°36.0'W, 1011 m, 5 Aug 2004, 1 female, 1 male, AB08–0038 and AB09–0016; *MF* 833–38, 51°42'54"N, 178°51'30"W, 582 m, 3 Aug 1983, 1 male in alcohol, USNM 77055; *MF* 833–47, 51°55'36"N, 176°52'48"W, depth unknown, 5 Aug 1983, 1 female, USNM 96529; *North Pacific*, 52°04'22"N, 176°58'01"E, 384 m, 20 Oct 2000, 1 male, USNM 1122539; *Ocean Olympic*, 52°09'06"N, 176°09'48"E, 384 m, 2000, 1 male, USNM 1122536; *Pacific Knight* 941–49, 52°46'N, 171°45'W, 0–340 m, 14 Jun 1994, 1 female, USNM 96528; *Pacific Knight* 941–75, 52°31'N, 173°30'W, 0–213 m, 20 Jun 1994, 1 male, USNM 96527; *Pacific Knight* 941–204, 53°06'N, 171°42'E, 0–455 m, 31 Jul 1994, 1 female, 1 male, USNM 96249 and 1123534; *Patricia Lee*, 52°21'09"N, 179°32'52"E, 375 m, 2000, 1 female, USNM 1122500; *Sea Storm* 90, 51°36'30"N, 177°10'48"W, 202–217 m, 4 Jul 2002, 1 male in alcohol, USNM 1123012; *Sea Storm* 91, 51°40'51"N, 177°10'49"W, 82 m, 4 Jul 2002, 1 male, USNM 1123284; *Sea Storm* 92, 51°33'34"N, 177°36'59"W, 367 m, 4 Jul 2002, 1 female in alcohol, 1 male, USNM 1123028–29; *Sea Storm* 107, 52°10'28"N, 175°14'14"E, 214 m, 8 Jul 2002, 1 female, 2 male in alcohol, USNM 1123001, -05, -31; *Sea Storm* 108, 52°11'32"N, 175°17'E, 208 m, 8 Jul 2002, 1 female, 1 male, both in alcohol, USNM 1122998 and 1123041; *Sea Storm* 109, 52°17'16"N, 175°20'56"E, 238 m, 8 Jul 2002, 2 indet. in alcohol, USNM 1122997 and 1123536; *Sea Storm* 111, 52°16'20"N, 175°59'13"E, 137 m, 9 Jul 2002, 1 male, 1123034; *Sea Storm* 112, 52°15'12"N, 176°00'42"E, 137 m, 9 Jul 2002, 1 indet., USNM 1122999; *Sea Storm* 134, 52°14'38"N, 176°01'25"E, 138 m, 15 Jul 2002, 1 indet. in alcohol, USNM 1123033; *Sea Storm* 138, 52°13'50"N, 175°14'31"E, 146 m, 16 Jul 2002, 1 female, 1 male, USNM 1122769 and 1122965; *Sea Storm* 148, 52°28'16"N, 173°19'10"W, 194 m, 21 Jul 2002, 2 male, USNM 1122748 and 1122752; *Sea Storm* 150, 52°30'47"N, 173°29'W, 213 m, 21 Jul 2002, 1 female, USNM 1122758; *Sea Storm* 151, 52°33'40"N, 173°19'10"W, 203 m, 21 Jul 2002, 1 female, 1 male, USNM 1122755–6; *Sea Storm* 155, 52°38'43"N, 172°16'23"W, 393–401 m, 22 Jul 2002, 2 indet., USNM 1122761, -64; *Shishaldin*, 54°07'N, 179°45'E, 366 m, 20 Feb 2000, 1 female, USNM 1122529; *Shishaldin*, 54°24'03"N, 179°36'11"E, 313 m, 25 Feb 2000, 1 female, USNM 1122532; *Shishaldin*, 53°56'54"N, 179°49'31"E, 318 m, 14 mar 2000, 1 indet., USNM 1122568;

Chew, coll., 51°32'N, 179°15'E, 218–289 m, 1 Sep 1968, 3 female, USNM 76573 and 96532; Haaga, coll. 53, 51°24'12"N, 177°37'20"W, 12 Jun 2000, 1 female, USNM 1122515; Haaga, coll., 54, 51°45'48"N, 178°09'47"W, 441–791 m, 10 Jun 2000, 1 female, 1 male, USNM 1122496 and AB02–30; McCloskey, coll., 52°03'26"N, 179°12'20"E, 475 m, 27 Apr 2000, 2 indet., USNM 1122449 and 1122461; McCloskey, coll., 52°17'45"N, 179°28'57"E, 28 May 2000, 1 female, USNM 1122964; McCloskey, coll., 51°16'46"N, 178°56'01"E, 373 m 21 May 2000, 2 indet., USNM 1122463; Miller, coll., 51°31'09"N, 176°53'14"W, 375 m, 2000, 1 female, USNM 1122498; Myer, coll., 52°15'N, 173°30'W, 208–274 m, 20 Nov 1973, 4 female, USNM 76567; Slear, coll., 52°07'54"N, 177°14'46"E, 238 m, 8 Dec 2000, 1 female, USNM 1122453; Slear, coll., 51°54'04"N, 179°17'28"E, 567 m, 9 Dec 2000, 2 female, USNM 1122450, -51; Stone, coll., 51°51.133'N, 179°50.9'E, 630 m, 8 Aug 2005, 1 female, AB05–0055; Tierny, coll., Cross Sound, Alexander Archipelago, 334 m, 30 Jul 1953, 1 indet., USNM 76569;

Description. Corallum essentially uniplanar or multiplanar, with occasional short side branches oriented perpendicular to flabellum. Branches anastomose only in larger colonies, the largest colony being the holotype of *Stylaster polyorchis*, which is 28 cm tall and 35 cm broad, having a massive, dense basal branch diameter of 3.9 cm (Fig. 17B). Distal branches circular to slightly flattened in cross section; larger-diameter branches often rectangular in cross section, the longer axis of the rectangle oriented perpendicular to plane of flabellum. Commensal spionid worm tubes not present. Coenosteum reticular-granular in texture, the coenosteal strips 50–55 μm in width, the slits being a discontinuous series of elongate pores about 15 μm in width; strips covered by small granules. Although generally reticulate in texture, coenosteal strips parallel and straight on exsert abaxial side of each cyclosystem (Fig. 18A). In some specimens the coenosteum is porcellaneous (*polyorchis* form, Fig. 18I), whereas in others (typical form, Fig. 18G–H) it appears to be more porous. Coenosteum white, pale orange, and pale pink.

Cyclosystems linearly arranged on both edges of branches, as well as on anterior face, and occasionally on posterior face. Those on branch edges often closely spaced, sometimes directly adjacent to one another (Fig. 18C) or even coalescent. Cyclosystems circular, elliptical, or irregular in shape, 1.0–1.3 mm in diameter, slightly flared, and standing slightly exsert from branch; gastropores circular, 0.40–0.45 mm in diameter. Gastropore tubes cylindrical and usually slightly curved (Fig. 18M) on distal branches, such that gastrostyle tip is difficult to see; ring palisade often absent or only poorly developed. Gastrostyles lanceolate, about 0.5 mm in height, and occupying only lower quarter of gastropore tube; H:D about 2.3–2.5. Gastrostyle bears spines up to 0.4 mm in length.

Dactylotomes 0.10–0.11 mm in width; dactylostyles inconspicuous. Range of dactylopores per cyclosystem is 7–17 ($n = 50$, average = 11.94 ($\sigma = 2.4$), mode = 10). Supernumerary dactylopores absent. Pseudosepta slender and fairly uniform in width, 0.11–0.13 mm in width, an adcauline diastema of about twice pseudoseptal width sometimes present.




Figure 18. *Stylaster campylecus*, **A** male syntype of *tylota* form, USNM 86000, **B–H, M** male holotype of *campylecus*, USNM 42870, **I** male holotype of *polyorchis* form, USNM 43266, **J** female syntype of *tylota* form, USNM 43263, **K** male holotype of *moseleyanus* form, USNM 42869, **L** female paratype of *campylecus*, USNM 52262: **A–B** distal branches showing male ampullae and striate coenosteum **C** stereo view of two aligned cyclosystems **D** rudimentary dactylostyle **E–F** cyclosystems **G–H** porous coenosteal texture **I** porcellaneous coenosteal texture **J** longitudinal section of a cyclosystem **K** gastrostyle **L** female ampulla **M** stereo view of a damaged cyclosystems showing curved gastropore, small dactylostyles, and lack of a ring palisade.

Female ampullae (Fig. 18L) large, smooth, superficial hemispheres 1.0–1.1 mm in diameter, often having a lateral efferent pore about 0.25 mm in diameter. Male ampullae (Fig. 18A–B) superficial swellings 0.4–0.5 mm in diameter, often clustered on anterior face.

Remarks. Fisher (1938: 505) discussed the similarities of *S. campylecus* and *S. polyorchis*, but concluded they were different species based on six minor differences: *S. polyorchis* had smaller cyclo systems, straighter gastropore tubes, no ridges on the inside of the gastropore tube, often linked cyclo systems, a ring palisade, and wrinkled female ampullae. Based on more detailed SEM observations on many more specimens, these differences are considered to be intraspecific variation, the holotype of *S. campylecus* itself having some linked cyclo systems and wrinkled female ampullae. The ring palisade is absent in most coralla, but in some specimens is weakly developed. *S. campylecus tylo-tus* was also compared to but differentiated from typical *S. campylecus* by Fisher (1938) in having a more delicate corallum, a larger ring palisade, a stouter gastrostyle, larger male ampullae, and more dactylo pores per cyclo system, but closer examinations also shows all these character to be within the range of variation of typical *S. campylecus* (see Table 2). Likewise, no significant differences could be found between *S. moseleyanus* and *S. campylecus*, and thus these four taxa are considered to be the same. Interestingly, types of three of the four taxa were collected at Albatross station 3480, prompting Fisher (1938:514) to remark about “the extraordinary number of species and subspecies dredged at station 3480,” considering the possibility that hybridism might be taking place. *S. campylecus* and *S. brochi* are the two most commonly collected stylasterids in the Aleutian Islands. *S. campylecus* can be distinguished from other Alaskan species in *Stylaster* (Group B) by its slightly flared cyclo systems and the linearity of the coenosteal strips near the cyclo systems (Table 2). The corallum was found to be 100% aragonitic according to Cairns and Macintyre (1992). Of 94 colonies examined, 44 are female, 39 male, and 11 indeterminate, resulting in a fairly equal sex ratio.

Distribution. Known from throughout the Aleutian Islands from Agattu Island to Unalaska, including Petrel and Bowers Banks, two disjunct records in Alexander Archipelago; 82–1011 m, but most records from 150–500 m.

***Stylaster leptostylus* (Fisher, 1938), new rank**

http://species-id.net/wiki/Stylaster_leptostylus

Figs 17E, 19A–I

Allopora moseleyana forma *leptostyla* Fisher, 1938: 514, pl. 52, figs. 1–3.

Allopora moseleyana Fisher, 1938: in part (pl. 50).

Stylaster moseleyanus forma *leptostylus*.—Cairns 1983b: 429.

Type material. Holotype: *Alb*-3480, 1 dry male colony 8.5 cm in height, and SEM stub 1544, USNM 43270 (Fig. 17E). Paratypes: *Alb*-3480, 4 female, 2 male colonies,




Figure 19. *Stylaster leptostylus*, **A–B, D–I** paratypes from USNM 76817, **C** holotype, USNM 43270: **A** stereo view of a male branch **B** a cyclostyle **C** gastrostyle **D** stereo view of a gastrostyle and female ampullae **E–F** coenosteal texture **G–H** gastrostyle **I** two dactylostyles.

dry, and SEM stub 1536–38, USNM 76817. **Type locality.** *Alb-3480*: 52°06'00"N, 171°45'00"W (off Seguam Island, Amukta Pass), 518 m.

Material examined. Types. *Alb-3480*, 2 female, 1 male colonies, dry, USNM 76555 (paratype of *A. moseleyana*).

Description. Corallum uniplanar, branching equal and dichotomous; no anastomosis of branches. Largest colony (the holotype, Fig. 17E) 8.5 cm in height and 7.0 cm in width, with a basal branch diameter of 8.4 mm. All branches circular in cross section. Commensal spionid polychaetes absent. Coenosteum reticulate-granular in texture, the coenosteal strips 45–55 μm in width, the slits about 15 μm in width; strips; strips not linear near cyclostyles. Coenosteum white.

Cyclosystems occur on branch edges and anterior face, but rarely on posterior face, those on edges not aligned in rows. Cyclosystems circular, 1.0–1.1 mm in diameter, and raised only slightly above branch coenosteum. Gastropores about 0.35 mm in diameter, leading to a slightly curved, funnel-shaped upper tube that narrows to a cylindrical lower part that houses the gastrostyle (Fig. 19G–H); ring palisade absent. Gastrostyles elongate-lanceolate, up to 0.6 mm in length, occupying lower half of gastropore tube, its pointed tip easily seen in an intact cyclosystem; H:D about 3.5.

Dactylotomes 0.10–0.13 mm wide; dactylostyles moderate in size, elements up to 50 μ m in length. Range of dactylopores per cyclosystem 7–12 ($n = 50$, average = 9.76 ($\sigma = 1.92$), mode = 9). Supernumerary dactylopores absent. Pseudosepta slender and fairly uniform in width (0.10–0.11 mm), an adcauline diastema of twice that width sometimes present.

Female ampullae (Fig. 19D) large, smooth, superficial hemispheres 0.9–1.1 mm in diameter, occasionally having a lateral efferent pore about 0.20 mm in diameter. Male ampullae (Fig. 19A) superficial swellings 0.45–0.55 mm in diameter, often clustered on anterior and posterior branch faces; apical efferent pores about 25 μ m in diameter.

Remarks. Fisher (1938) originally differentiated forma *leptostyla* from typical *Allopora moseleyana* (= *Stylaster campylecus*) based on its more slender gastrostyle and smaller male ampullae. We cannot find a difference in ampulla size between forma *leptostyla* and typical *S. campylecus*, but the gastrostyle of *leptostyla* is longer and more slender than that of *S. campylecus*, occupying more of the gastropore tube (and thus having a higher H:D ratio) and being more visible from external view. In addition to this difference, forma *leptostyla* differs from *S. campylecus* in having equal, non-anastomosing branching; branches circular in cross section; less exsert cyclosystems; non-linearly arranged cyclosystems; fewer dactylopores per cyclosystem; and consistently reticulate coenosteal strips (see Table 2). For these reasons this form is elevated to species rank but is considered to be closely related to *S. campylecus*.

Distribution. Known only from type locality.

Stylaster trachystomus (Fisher, 1938), new rank

http://species-id.net/wiki/Stylaster_trachystomus

Figs 17H, 20A–M

Allopora campyleca trachystoma Fisher, 1938: 510–511, pl. 45, fig. 2, pl. 46, pl. 54, figs 1–1b.—?Naumov 1960: 579.—Lowenstam 1964: 382 (mineralogy).

Stylaster campylecus trachystomus.—Cairns 1983b: 430.—Cairns and Macintyre 1992: 100–101 (mineralogy).—Wing and Barnard 2004: 27.—Heifetz et al. 2005: 134 (listed).—Stone and Shotwell 2007: 108 (listed).—Jameison et al. 2007: 224 (listed).




Figure 20. *Stylaster trachystomus*, **A–L** holotype, USNM 43265, **M** male paratype, USNM 76811: **A** stereo view of a cyclosystem showing internal decurrent ridges within gastropore **B** a cyclosystem **C** a damaged cyclosystem showing decurrent ridges **D** stereo view of longitudinal section of a cyclosystem and some female ampullae **E** a gastrostyle and ring palisade **F** porous pseudosepta **G** stereo view of ridged coenosteum and a small female ampulla **H** branch segment **I** coenosteal texture **J** rudimentary dactylostyle **K–L** clusters of female ampullae **M** male ampullae.

Type material. Holotype: *Alb*-4784, 1 dry female colony 9 cm in height, and SEM stubs 1516, 1541, USNM 43265 (Fig. 17H). Paratypes: *Alb*-4784, 2 female, 2 male, 1 indet. colonies, and SEM stub 1517–18, dry, USNM 76811. **Type locality.** *Alb*-4784: 52°55'40"N, 173°26'E (off East Cape, Attu Island, Aleutians), 247 m.

Material examined. Types; *Delta* 5999–8E-3, 51°21.042'N, 179°30.483'W, 115 m, 4 Jul 2003, 1 male, AB10–0001; *Delta* 6230–20–18, 52°28.142'N, 173°35.882'W, 190 m, 8 Jul 2004, 5 male branch fragments in alcohol, AB10–0004; *Shishaldin*, 54°54.09'N, 178°37.27'E, 366 m, 11 Feb 2000, 1 female, 2 male, AB00–45.

Description. Corallum uniplanar, occasionally with short side branches oriented perpendicular to flabellum; branch anastomosis occurs only in large-diameter basal branches. Largest colony (paratype) 12 cm tall and 8 cm wide, with a basal branch diameter of 3×1.5 cm. Distal branches circular to slightly flattened in cross section. Commensal spionid worm tubes common. Coenosteum reticulate-granular in texture, the coenosteal strips 70–80 μ m wide, the slits discontinuous and about 10–15 μ m wide, the small granules producing a rough texture. On exsert cyclo systems, coenosteal strips are parallel and straight. Most coralla bear some short coenosteal ridges, called “spinous outgrowths” by Fisher (1938: 511), these thin ridges aligned with the branch (up to 1 mm long, 1 mm tall, and only 0.1 mm in width) or aligned with some of the pseudosepta on exterior wall of a cyclo system (Fig. 20G–H). Coenosteum pale pink-orange.

Cyclo systems arranged uniformly on branches, not linearly, but fewer in number on posterior face. Cyclo systems rarely circular in shape, rather elliptical or asymmetrical, the longer axis of the ellipse up to 1.8 mm. Gastro pore tubes cylindrical, slightly curved, and quite deep (up to 3 mm); a well-developed ring palisade present near gastro style tip, composed of squat elements about 50 μ m in height and width. Gastro styles lanceolate, up to 0.67 mm, having a H:D of about 3.1, and occupying lower quarter of gastro pore tube. Because of curvature of the long gastro pore tube, the short gastro style, and the wide pseudosepta, the gastro style tip is rarely seen when viewed from above.

Dactylo tomes 0.085–0.10 mm in width, supernumerary dactylo pores being quite rare, usually associated with pseudosepta. Dactylo style inconspicuous (Fig. 20J). Range of dactylo pores per cyclo system 8–18 ($n = 50$, average = 11.82 ($\sigma = 2.15$), and mode = 11). Pseudosepta slender (0.07–0.21 mm wide), adcauline diastemas twice that width sometimes present; surface of pseudosepta quite porous (Fig. 20F).

Female ampullae (Fig. 20D, G, K–L) superficial hemispheres 1.0–1.3 mm in diameter, having a knobby or papillose surface; efferent pores not evident in material at hand. Male ampullae (Fig. 20M) also superficial swellings 0.50–0.55 mm in diameter. Both types of ampullae clustered on anterior and posterior branch faces.

Remarks. Fisher (1938) described this taxon as a subspecies of *S. campylecus*, distinguishing it from the nominate subspecies by having spongy pseudosepta (Fig. 20F), decurrent pseudoseptal ridges on the inside of the gastro pore tube (Fig. 20C), and spiny coenosteal outgrowths (Fig. 20G–H). *S. trachystomus* is otherwise similar to typical *S. campylecus* in gastro pore tube and gastro style shape and number of dactylo pores

per cyclosystem, but is also similar to *S. brochi* in arrangement of cyclosystems and in hosting commensal spionid polychaetes; it also has unique characteristics as mentioned above (see also Table 2). Although few colonies are known of this species, because it does present a unique set of characteristics, it is considered valid and herein elevated to species rank. The corallum was found to be 100% aragonitic according to Cairns and Macintyre (1992).

Distribution. Aleutian Islands: from off Attu Island and north of Amalia Island (Andreanof Islands), including Bowers Bank; 115–366 m.

***Stylaster parageus parageus* (Fisher, 1938), new rank**

http://species-id.net/wiki/Stylaster_parageus_parageus

Figs 21A, 22A–J

Allopora campyleca paragea Fisher, 1938: 507–509, pl. 41, figs 1–1d, pl. 43.—Thompson and Chow 1955: 30 (mineralogy).

Stylaster (Allopora) boreopacificus forma *typica*.—Broch, 1936: 56–60, in part: specimens from Alaska, pl. 8, fig. 22, pl. 9, figs 23, pl. 10, figs 24–25, text figs 17c–d.

?*Stylaster campylecus*.—Lowenstam 1964: 382 (mineralogy).

Stylaster campylecus parageus.—Cairns 1983b: 430.—Cairns and Macintyre 1992: 100–101 (mineralogy).—Wing and Barnard 2004: 27 (listed), not fig. 32.—Heifetz et al. 2005: 134 (listed).—Stone and Shotwell 2007: 108 (listed).—Jameison et al. 2007: 224 (listed).

Type material. Holotype: 1 dry male colony 13 cm in height, plus many tiny branch fragments, and SEM stub 1514, USNM 42871 (Fig. 21A). Paratypes: *Alb*-4245, 1 female colony, and SEM stub 1515, USNM 76812; near Sitka, Alaska, coll. E. R. Ricketts, 1 female, deposition unknown (not seen); Alaska, deposition unknown (not seen); Yakutat Bay, Alaska, 152 m, specimens reported by Broch (1936), Zoological Museum Copenhagen (not seen). **Type locality.** Tenakee Springs, near Juneau, Alaska, depth unknown.

Material examined. Types; *Alaskan Leader* 21–91A, 59°31'18"N, 144°42'42"W, 401–600 m, 31 Jul 2002, 1 male, USNM 1122485; Lambert, coll., 55°18'N, 129°57'18"W, 23 m, 28 Mar 1976, 1 male in alcohol, USNM 76976; Freege and Worth, coll., 57°48'N, 134°02'W, 30 m, 17 Apr 2000, 4 branches in alcohol, USNM 1122523; “an Indian”, coll., Sitka Bay, depth unknown, about 1884, 2 male branches, USNM 4192.

Description. Corallum consisting of overlapping flabella, also with short branchlets oriented perpendicular to flabella, producing what Fisher (1938:507) calls “subflabellate” or a “flattened bush.” Branches do not anastomose, are circular to slightly flattened in cross section, and distally are quite delicate. Largest colony (the holotype, Fig. 21A) 13 cm tall and wide, with a basal branch diameter of 3.2 cm. Commensal spionid worm tubes common. Coenosteum reticulate-granular in texture (Fig. 22E–F), the coe-




Figure 21. Color figures of skeleton of various Alaskan stylasterids: **A** *Stylaster parageus parageus*, **B** *S. parageus columbiensis* **C** *S. classotomus* **D** *S. crassisepalum* **E–G** *S. alaskanus*. **A** holotype, USNM 42871 **B** holotype, USNM 1122462 **C** holotype, USNM 43268 **D** holotype, USNM 1122531 **E** holotype of *S. gemmascens alaskanus*, USNM 43269 **F** holotype of *S. cancellatus*, USNM 43267 **G** large colony, USNM 1122454.




Figure 22. *Stylaster parageus parageus* **A–E, G–H, J** holotype (male), USNM 42871 **F I** female paratype, USNM 76812: **A** stereo view of three cyclosystems and male ampullae **B** a cyclosystem **C** gastrostyle tip surrounded by ring palisade **D** stereo view of a gastrostyle **E–F** coenosteal texture **G–H** rudimentary dactylostyles **I** female ampullae **J** cross section of internal male ampullae.

nosteal strips 60–65 μm in width, the slits only 6–8 μm wide, the strips covered with low rounded granules, altogether presenting a smooth dense aspect. Coenosteum white.

Cyclosystems occur on branch edges and anterior face, but rarely on posterior face. Cyclosystems circular in shape, small (0.9–1.0 mm in diameter, not 0.6 mm as stated by Fisher, 1938), and only slightly elevated above coenosteum; gastropores circular and quite narrow (0.25–0.30 mm in diameter, Fig. 22B). Gastrostyle tubes cylindrical, slightly curved, such that gastrostyle tip rarely seen in view from above; well-developed ring palisade present (Fig. 22C), consisting of elongate elements oriented longitudinally,

each about 45 μm in length and 30 μm in width. Gastrostyles lanceolate, up to 0.5 mm in height, and occupying the lower half of third of the gastropore tube; H:D about 2.7.

Dactylotomes 0.09–0.10 mm wide, the inner slit very shallow, resulting in a ring of essentially apical dactylopores surrounding a thick-walled gastropore tube (Fig. 22B). Dactylostyles well developed, the elements up to 35 μm in height and 10–11 μm in diameter (Fig. 22H). Range of dactylopores per cyclosystem 5–11 ($n = 50$, average = 8.54 ($\sigma = 1.20$), mode = 8). Supernumerary dactylopores present but not common. Pseudosepta 0.12–0.13 mm in width; diastemas rare.

Female ampullae (Fig. 22I) superficial hemispheres 0.8–1.0 mm in diameter, the lateral efferent pores being 0.15–0.20 mm in diameter. Male ampullae (Fig. 22A, J) partially submerged in coenosteum (internal) on distal branches, entirely internal on larger-diameter branches, the outer diameter being about 0.5 mm, the internal diameter of internal ampullae about 0.42 mm; male ampullae often clustered on branch faces.

Remarks. Although Fisher (1938:508) included this taxon as a subspecies of *S. campylecus*, considering it to be “the southern shallow-water race of *campyleca*,” there are sufficient differences to warrant raising this subspecies to species rank (Table 2). Distinctive features include its bushy delicate colony, extremely small gastropores surrounded by a thick wall, and relatively low number of dactylopores per cyclosystem. Geographically it occurs only off southeastern Alaska in relatively shallow water, not in the Aleutian Islands. Comparisons to the other subspecies are made in the following account.

Distribution. Bays and inland passages of southeastern Alaska from off Kayak Island to just north of Dixon Entrance (i.e., Prince of Wales Islands and Portland Canal); 23–401 m.

***Stylaster parageus columbiensis* Lindner & Cairns, ssp. n.**

urn:lsid:zoobank.org:act:8B19F022-F84F-4AA7-B16B-E29DB8A8756D

http://species-id.net/wiki/Stylaster_parageus_columbiensis

Figs 21B, 23A–J

Allopora campyleca Fisher, 1938: 506, in part (specimen from *Alb*-4230).

Type material. Holotype: FOMC 1162 (209), male colony, dry, and SEM stubs 1389–91, USNM 1122462 (Fig. 22B). Paratypes: *Alaskan Leader* 21–108, 54°27'N, 133°55'48"W, 215–563 m, 10 Jul 2002, 1 male, USNM 1122483; *Alb*-4230, 55°N, 131°W, 198–439 m, 7 Jul 1903, 1 male, USNM 76815 (paratype of *A. campylecus*); FOMC 1158 (206), 48°07'546"N, 125°05'46"W, 246 m, 10 Jul 2008, 1 male in alcohol, USNM 1122465; FOMC 1159 (161), 48°07'56"N, 125°05'45"W, depth unknown, 10 Jul 2008, 3 males, USNM 1122472; FOMC (162), 48°07'56"N, 125°05'46"W, depth unknown, 10 Jul 2008, 1 male in alcohol, USNM 1122471; FOMC 1159 (166), 48°07'46"N, 125°05'41"W, 273 m, 10 Jul 2008, 1 male in alcohol, USNM 1122473; FOMC 1159 (169), 48°07'48"N,




Figure 23. *Stylaster parageus columbiensis*, **A–H, J** holotype (male), USNM 1122462, **I** female paratype, USNM 1117112: **A** stereo view of branch segment including male ampullae **B** a cyclosystem **C** longitudinal fracture of a cyclosystem **D** stereo view of a gastrostyle and rudimentary ring palisade **E–F** rough and smooth coenosteal texture **G** gastrostyle tip and ring palisade **H** two dactylostyles **I** female ampulla **J** male ampullae.

125°05–41"W, 264 m, 10 Jul 2008, 1 male in alcohol, USNM 1122467; FOMC 1162 (203 and 205), 48°08'26"N, 125°11'28"W, 265 m, 12 Jul 2008, 2 males in alcohol, USNM 1122464, and -68; Lindner, coll., AL470 (41), Race Rocks, near Sooke Community, Straits of Juan de Fuca, British Columbia, depth unknown, Jul 2002, 1 male, USNM 1096625; Lindner, coll., AL466 (25), off Triangle Island, British Columbia, depth unknown, 1 male in alcohol, USNM 1096628; *Ropos* 956 (103), 48.15605°N, 124.9973°W, 288 m, 30 May 2006, 1 male, USNM 1117117; *Ropos* 957 (116), 48°08'42"N, 125°11'26"W, 285 m, 31 May 2006, 1 female in al-

cohol, SEM stub 1402, USNM 1117112. **Type locality.** 48°08'31"N, 125°11'01"W (off Cape Flattery, Washington), 273 m.

Etymology. Named for the region from which it is primarily known: British Columbia.

Material examined. Types.

Description. Corallum shape and branching similar to that of typical subspecies: bushy or flattened bushy with delicate terminal branches. Holotype 7 cm tall and 7 cm wide, with a basal branch diameter of 10.1 × 8.1 mm; largest corallum (USNM 1096625, Fig. 21B) 8.5 × 8.0 cm, with a massive basal branch diameter of 3 cm. Commensal spionid worm tubes absent. Coenosteum reticulate-granular in texture, the coenosteal strips 50–60 μm wide, bordered by slits about 10 μm wide, the strips originally covered with irregularly-shaped nodules that are subsequently covered with smoother granular coenosteum (Fig. 23E shows the transition). Coenosteum white.

Cyclosystems occur exclusively on edges of distal branches, but also on anterior face of larger-diameter branches. Cyclosystems circular to elliptical in shape, the larger axis ranging from 1.1 to 1.5 mm, the abaxial edge slightly raised above coenosteum; gastropores circular, 0.45–0.50 mm in diameter. Gastropore tubes somewhat inflated in upper half, changing to a narrow cylinder proximally; a well-developed ring palisade present as in the typical subspecies, the narrow pointed gastrostyle tip projecting through the ring palisade constriction into upper chamber (Fig. 23G) and thus easily visible from above. Gastrostyles elongate-lanceolate (Fig. 23D), occupying lower half of gastropore tube, and up to 0.67 mm in length, having a H:D ratio of 3.1–3.7.

Dactylotomes 0.09–0.10 mm in width, the inner slit short but longer than that of typical subspecies. Dactylostyles well developed, the cylindrical elements up to 38 μm in height and about 14 μm in diameter. Range of dactylopores per cyclosystem 6–13 ($n = 50$, average = 9.38 ($\sigma = 1.43$), and mode = 9). Supernumerary dactylopores rare. Pseudosepta 0.17–0.28 mm in width; diastemas uncommon.

Female ampullae (Fig. 23I) smooth superficial hemispheres 0.9–1.0 mm in diameter, with lateral efferent pores about 0.23 mm in diameter. Male ampullae (Fig. 23J) superficial swellings up to 0.5 mm in diameter, with small apical efferent pores. Both types of ampullae clustered on anterior and posterior faces.

Remarks. Subspecies *columbiensis* resembles the typical subspecies in many ways, including colony shape and branching, and ring palisade shape, but also differs in a number of small but consistent characteristics (Table 2). Subspecies *columbiensis* has larger cyclosystems and gastropores, a slightly higher average number of dactylopores per cyclosystem, a more commodious upper gastropore chamber and deeper dactylotome slits, and a more elongate gastrostyle that is easily seen from above. Dixon Entrance appears to be the border between the two subspecies, the typical subspecies occurring to the north and *columbiensis* occurring to the south of this body of water. Because of the overall similarity of the two taxa, and

their consistent minor differences, including their geographic separation, *columbiensis* is considered as a subspecies or *S. parageus*. Of 16 colonies examined, 15 are male and one is female.

Distribution. Entire coast of British Columbia (Canada) from Dixon Entrance to off Cape Flattery, Washington (USA); 246–285 m.

Stylaster (Group C)

Diagnosis. Species of *Stylaster* in which coralla form branching colonies, and in which cyclosystems are arranged exclusively on branch edges; branching delicate.

***Stylaster alaskanus* Fisher, 1938**

http://species-id.net/wiki/Stylaster_alaskanus

Figs 21E–G, 24A–M

Stylaster gemmascens alaskanus Fisher, 1938: 500–502, pl. 47–48, pl. 54, fig. 2.—Boschma 1957: 11.

Stylaster cancellatus Fisher, 1938: 502–503, pl. 35, figs 2–2c, pl. 39–40.—Boschma 1957: 4.—Cairns 1983b: 430 (listed).—Heifetz 2002: 22 (listed).—Wing and Barnard 2004: 11, 27, fig. 33.—Heifetz et al. 2005: 22, 134, 137 (listed).—Stone and Shotwell 2007: 108 (listed).—Brooke and Stone 2007: 525–530, fig. 2D, 3B.—Jameison et al. 2007: 224 (listed).—Lindner et al. 2008: 3, supplemental Table S1: 2 (phylogeny and DNA sequence).

Stylaster alaskanus: Cairns 1983b: 430 (elevation to species rank).—Wing and Barnard 2004: 11, 27.—Stone and Shotwell 2007: 108 (listed).

Stylaster alaskana: Heifetz et al. 2005: 133 (listed).—Jameison et al. 2007: 224 (listed but misspelled).

Type material. *Stylaster gemmascens alaskanus*: Holotype, *Alb*-3480, 1 dry male branch fragment 4.5 cm in length, USNM 43269 (Fig. 21E); 5 (perhaps one broken from the four mentioned by Fisher) dry paratype branches (2 female, 2 male, 1 indet.), and SEM stubs 1511–12, *Alb*-3480, USNM 86006. **Type locality.** *Alb*-3480: 52°06'N, 171°45'W (Amukta Pass, Aleutian Islands), 518 m.

Stylaster cancellatus: Holotype, *Alb*-3480, 1 dry female branch fragment 9 cm in height, and SEM stub 1509, USNM 43267 (Fig. 21F); 7 dry paratype branches (2 female, 5 male), and SEM stub 1510, *Alb*-3480, USNM 86005. **Type locality.** *Alb*-3480: 52°06'N, 171°45'W (Amukta Pass, Aleutian Islands), 518 m.

Material examined. Types of *S. alaskanus* and *S. cancellatus*; *Alaskan Leader* 35, 51°50.9'N, 175°08'W, 178–365 m, 14 Jun 2002, 1 male, AB02–31; *MF* 833–38, 51°42'54"N, 178°51'30"W, 582 m, 3 Aug 1983, 2 female, USNM 96526; *MF*




Figure 24. *Styaster alaskanus*, **A–C, M** male paratype of *S. cancellatus*, USNM 86005 **D–E, H, J, L** female paratype of *S. gemmascens alaskanus*, USNM 86006, **F–G, I, K** holotype (female) of *S. cancellatus*, USNM 43267: **A** stereo view of branch tip with male ampullae **B–C, J** cyclosystems and male ampullae **D** stereo view of cyclosystem and female ampulla with a distinct efferent pore **E** cyclosystem and female ampulla **F** coenosteal texture **G** stereo view of gastrostyle and ring palisade **H** a cyclosystem **I** rudimentary dactylostyle **K–L** female ampullae **M** cluster of male ampullae.

833–56, off Great Sitkin I., 249 m, 6 Aug 1983, 2 male, USNM 77045; *Alaska Sea*, 52°09'24"N, 173°44'19"E, 318 m, 5 Feb 2000, 1 indet., USNM 1122566; *Alb-3480*, see above, 8 female and 1 male (non-type), USNM 76722; *Dominator* 971–135, 51°37'22"N, 178°26'28"W, 163 m, 14 Jul 1997, 2 female, USNM 1123365; *Dominator* 971–201, 51°54'28"N, 175°49'00"E, 378 m, 1 Aug 1997, 1 indet., USNM 1123361; *Ocean Olympic*, 52°04.78'N, 177°13.39'E, 256 m, 1 female, AB00–57; *Pacific Knight* 941–204, 53°06'N, 171°42'E, 455 m, 31 Jul 1994, 4 female some in alcohol, USNM 96246 and 1123533; *Sea Storm* 36, 53°05'45"N, 171°41'56"E, 453–458 m, 19 Jun 2002, 1 female, 1 male, USNM 1123277–78; *Sea Storm* 108, 52°11'32"N, 175°16'54"W, 208–215 m, 8 Jul 2002, 1 indet. in alcohol, USNM 1027822; *Sea Storm* 138, 52°13'50"N, 175°24.2'E, 146 m, 16 Jul 2002, 2 female, 2 male, USNM 1122766–68, 1122898; *Sea Storm* 148, 52°28'16"N, 173°29'35"W, 194 m, 21 Jul 2002, 2 female, USNM 1122757; *Shishaldin*, 54°24'04"N, 177°37'43"E, 154 m, 9 Feb 2000, 1 female, USNM 1122533; *Shishaldin*, 54°24'03"N, 179°36'11"E, 313 m, 25 Feb 2000, 4 indet., USNM 1122532; McClusky, coll., 52°01'47"N, 179°56'14"W, 362 m, 5 May 2000, 1 female, USNM 1122454; McClusky, coll., 51°17'58"N, 179°00'48"E, 265 m, 25 May 2000, 3 male, USNM 1122455 and AB01–92; University of Washington, 51°32'N, 179°15'W, 278–289 m, 1 Sep 1968, 4 male, USNM 76632 and 96530; A. Vatter, coll., 52°04'25"N, 177°11'00"E, depth unknown, 1 male, USNM 1122466.

Description. Colonies uniplanar or multiplanar, additional flabella usually oriented perpendicular to original flabellum. Branches highly anastomotic, smooth large-diameter branches forming a framework for infilling by smaller-diameter branches, resulting in a sieve-like reticulum (Fig. 21G). Largest colony (USNM 1122768) 28 cm in height, having a massive, dense basal branch diameter of 5 cm. Distal branches circular to rectangular in cross section. Spionid worm tubes do not occur in this species. Coenosteum reticulate-granular in texture, the coenosteal strips 50–60 µm wide, bordered by short slits 6–8 µm wide; strips covered with small granules 6–7 µm in width; strips arranged linearly adjacent to cyclostyles. In some colonies, such as the type of *S. alaskanus*, linearly arranged, apically perforate processes up to 0.2 mm tall (Figs 24D–E, J) occur on coenosteum, these structures of unknown function. Coenosteum light orange, pink, or white.

Cyclostyles circular, elliptical or irregular in shape, 0.9–1.3 mm in greater diameter, occurring exclusively on branch edges; gastropores circular, 0.40–0.45 mm in diameter. Gastropore tubes straight and funnel-shaped, having a well-developed ring palisade (Fig. 24G) composed of rounded elements about 40 µm in diameter. Gastrostyles slender and pointed, up to 0.5 mm in height, the gastrostyle tip easily seen when viewed from above. Very rarely two gastrostyles may be present in one gastropore tube.

Dactylotomes 80–85 µm in width; dactylostyles inconspicuous. Range of dactylotomes per cyclostyle 7–14 ($n = 50$, average = 11.30 ($\sigma = 1.57$), and mode = 13). Supernumerary dactylotomes absent. Pseudosepta of cyclostyles usually only slightly exert, but if two cyclostyles are in close proximity, pseudosepta may become quite exert. Pseudosepta of uniform thickness; diastemas, if present at all, are narrow.

Female ampullae (Fig. 24D–E, K–L) prominent hemispheres up to 0.9 mm in diameter, often bearing several low ridges or short spines; lateral efferent pore about 0.25 mm in diameter. Male ampullae (Fig. 24A–B, J, M) superficial, densely clustered, and also irregular in shape, up to 0.60 mm in diameter, each with a small efferent pore 40–60 μm in diameter. Efferent pores sometimes at the apex of a small curved spine.

Remarks. Originally described as a subspecies of *S. gemmascens* by Fisher (1938), Cairns (1983b) later raised *alaskanus* to species status. Herein it is synonymized with *S. cancellatus*, but since it has page priority in Fisher (1938), *S. alaskanus* is considered to be the senior synonym (see synonymy above). The types of *S. alaskanus* differ from most other specimens assigned to this species in having numerous coenosteal papillae, called “thorny outgrowths” by Fisher (1938: 501), which are often aligned in short ridges. Also, the coenosteum is white and the branches of the type series do not anastomose. However, all these differences are considered to be intraspecific variation, as some other specimens assigned to this species have a white corallum, and small colonies, like the types of *S. alaskanus*, usually do not have anastomosing branches. Furthermore, rarely some coralla of “*cancellatus*” have coenosteal papillae, which may be a reaction to an unfavorable environmental habitat. Thus, the type series of *S. alaskanus*, which was collected at the same station as the type of *S. cancellatus*, is considered to be a somewhat aberrant specimen of what was more widely known as *S. cancellatus*. A similarly shaped, sieve-like, reticulate colony is known for *Errinopsis reticulum* Broch, 1951, however, in that species the reticulum is composed of equally-sized branches, whereas in *S. alaskanus* the reticulum is composed of thick and thin branches. *S. alaskanus* is distinguished from all other species in its group by having a corallum shaped as a sieve-like reticulum. It is further distinguished from those Alaskan species of group C by having ridged female ampullae (Table 2). Among the 59 specimens examined, 25 are female, 22 male, and 12 indeterminate, a fairly even sex ratio.

Distribution. Known from throughout Aleutian Islands from west of Attu Island to Amukta Pass, including Bowers Bank; 146–582 m.

Stylaster elassotomus Fisher, 1938

http://species-id.net/wiki/Stylaster_lassotomus

Figs 21C, 25A–G

Stylaster elassotomus Fisher, 1938: 499–500, pl. 41, fig. 3, pl. 42, figs 1–1c, pl. 49, fig. 1.—Boschma 1957: 7.—Wing and Barnard 2004: 11 (key), 28.—Heifetz et al. 2005: 134137 (listed).—Stone and Shotwell 2007: 108 (listed).—Jameison et al. 2007: 224 (listed).

Type material. Holotype: 1 dry male colony, and SEM stub 1507, USNM 43268 (Fig. 21C). The pale pink paratype 5 cm in length (USNM 86007) mentioned by Fisher is an unidentified *Stylaster* having little in common with the holotype. **Type locality.** *Albatross* 4781, 52°14'30"N, 174°13'E (off Agattu Island), 882 m.




Figure 25. Holotype (male) of *Stylaster elassotomus*, USNM 43268: **A** stereo view of curved gastropore tube **B** cyclosystems and supernumerary dactylopores **C** poorly preserved gastrostyle and ring palisade **D** truncated gastrostyle showing ring palisade **E** coenosteal texture **F** stereo view of cyclosystem showing shallow dactylotomes **G** male ampulla.

Material examined. Types.

Description. Holotype (Fig. 21C) colony bushy, 5.5 cm tall and 4.5 cm wide, attached by a basal branch 6 mm in diameter. Branches do not anastomose and are circular in cross section. Coenosteum reticular-granular in texture, the strips 55–60 μm in width, separated by slits 10–14 μm wide; strips covered with small angular granules. Spionid worm tubes absent. Coenosteum white.

Cyclosystems circular to slightly elliptical, 1.0–1.2 mm in diameter, arranged exclusively on branch edges in an alternating fashion, most projecting perpendicular to branch. Gastropores circular and quite large (0.5–0.6 mm in diameter), occupying up

to 55% of cyclosystem diameter (Fig. 25F). Gastropore tube cylindrical, curved, and long, such that gastrostyle tip is rarely seen when viewed from above (Fig. 25A). A delicate ring palisade (Fig. 25C–D) occurs near gastrostyle tip, composed of slender cylindrical elements up to 60 μm long and 25 μm in diameter. Gastrostyle slender (H:D = 4.4), the illustrated style 0.53 mm in height, occupying lower quarter of gastropore tube (Fig. 25A).

Dactylotomes 60–80 μm wide but very short as well as having short internal slits, thus contributing to the large size of the gastropore tube (Fig. 25F); dactylostyles inconspicuous. Range of dactylopores per cyclosystem 11–17 ($n = 21$, average = 14.40 ($\sigma = 1.36$), and mode = 15). Supernumerary dactylopores common (Fig. 25B), 70–100 μm in diameter. Pseudosepta same width as dactylotomes; diastemas rare.

Female ampullae unknown. Male ampullae (Fig. 25G) hemispherical, 0.5–0.6 mm in diameter.

Remarks. Despite access to a diverse stylasterid collection from the Aleutian Islands, no unequivocal additional specimens could be identified as *S. elassotomus*, even the paratype of the species not considered to be conspecific. Its distinctive characteristics of having a highly curved gastropore tube, slender ring palisade elements, and a very large gastropore surrounded by very short dactylotomes and pseudosepta distinguish it from all other Alaskan species (see Table 2).

Distribution. Known only from type locality.

***Stylaster crassiseptum* Lindner & Cairns, sp. n.**

urn:lsid:zoobank.org:act:D638F3B7-D697-40BE-B502-5D4F5F9E519B

http://species-id.net/wiki/Stylaster_crassiseptum

Figs 21D, 26A–J

Type material. Holotype: dry female branch fragment 12 cm in length, plus many smaller broken pieces, and SEM stubs 1520–1521, 1546, USNM 1122531 (Fig. 21D). Paratypes: *North Pacific*, 52°04'17"N, 176°59'21"E, 366 m, date unknown, 1 female, USNM 1122511; *Shishaldin*, 54°00'41"N, 179°46'52"E, 291 m, 14 Mar 2000, 1 female, USNM 1122497; *Shishaldin*, 53°56'24"N, 179°49'31"E, 318 m, 14 Mar 2000, 1 male, and SEM stub 1522–23, USNM 1122519; *Shishaldin*, 51°48.49'N, 174°29.92'W, 531m, 2000, 1 female, USNM 112527. **Type locality.** *Shishaldin* station, 53°59'50"N, 179°46'52"E (off Bowers Bank, Aleutian Islands), 291 m, 5 Mar 2000.

Etymology. The specific name *crassiseptum* (from the Latin *crassus*, meaning “thick” + *septum* meaning “wall”), in reference to the wide pseudosepta of the cyclosystems.

Material examined. Types.

Description. Colonies primarily uniplanar, having no branch anastomosis. Holotype (Fig. 21D) a branch fragment 12 cm in length and 1 cm in basal branch diameter; largest specimen (USNM 1122527) an intact colony 24 cm in height and 19 cm wide, with a basal branch diameter of 4.5 cm. Distal branches circular in cross section, basal branches somewhat rectangular in cross section, the longer axis perpendicular to colony plane;




Figure 26. *Stylaster crassisseptum* **A, D, H–I** male paratype, USNM 112519, **B–C, E–G, J** holotype (female), USNM 1122531: **A** stereo view of cyclosystems and a cluster of male ampullae **B** branch segment showing sympodial arrangement of cyclosystems **C–D** cyclosystems **E** thick pseudosepta and thin dactyloctomes **F** coenosteal texture **G** robust dactylostyle **H** stereo view of gastropore tube and male ampulla **I** gastrostyle surrounded by well-developed ring palisade **J** female ampulla.

symbiotic polychaetes absent. Coenosteum reticulate-granular in texture, coenosteal strips 60–70 μm in width, separated by very narrow slits only 3–5 μm wide; coenosteal strips not linearly arranged near cyclosystems. Coenosteal granules very low and smooth, conferring a shiny or porcellaneous texture to branches. Coenosteum dense and uniformly pale orange.

Cyclosystems circular, slightly exsert, and relatively small (0.7–1.0 mm in diameter), occurring exclusively on branch edges (Fig. 26A) in alternating fashion (Fig. 26B), most projecting perpendicular to branch. Cyclosystems well spaced, separated by 1–3 cyclosystem diameters from one another on one side of a branch. Gastropores circular, about 0.5

mm in diameter; gastropore tubes cylindrical and slightly curved, having a well-developed ring palisade (Fig. 26I) composed of numerous large squat elements up to 35 μm in height and 50 μm in diameter. Gastrostyles lanceolate and slender, the figured style being 0.48 mm in height and only 0.15 mm in diameter (Fig. 26I), the tip usually seen when viewed from above. Gastrostyle covered with blunt spines up to 33 μm in length.

Dactylotomes slender and uniform in width (60–70 μm); dactylostyles quite robust (Fig. 26G), the large cylindrical elements (up to 45 μm tall and 13 μm in diameter) almost completely filling the lower dactylopore, but because of the narrow dactylopores and exsert pseudosepta, the dactylostyles are barely visible in an intact cyclo-system. Range of dactylopores per cyclo-system 6–12 ($n=60$, average = 8.89 ($\sigma = 1.4$), mode = 9). Supernumerary dactylopores not present. Pseudosepta solid, exsert, and variable in width (0.15–0.31 mm in width), or up to five times width of adjacent dactylotomes (Fig. 26C–E). Diastemas rare.

Female ampullae (Fig. 26J) smooth superficial hemispheres 0.9–1.1 mm in diameter, with a lateral efferent pore about 0.3 mm in diameter. Male ampullae (Fig. 26A, H) often clustered, slightly irregular in shape, 0.45–0.50 mm in diameter, often with one or more tiny (28–30 μm in diameter) apical efferent pores.

Remarks. Although similar to *S. alaskanus*, *S. crassiseptum* is distinguished by having non-anastomosing branches, a curved gastropore tube, relatively wide pseudosepta, and more robust dactylostyles (Table 2).

Distribution. Aleutian Islands from off Kiska to off Atka Islands, Bowers Bank; 291–531 m.

Genus *Stylantheca* Fisher, 1931

<http://species-id.net/wiki/Stylantheca>

Diagnosis. Colonies encrusting, forming thin laminae on rocks and shells, rarely forming short stubby branches. Coenosteum reticulate-granular and of many colors; coenosteal papillae quite common. Gastro- and dactylopores arranged in cyclo-systems, often with more than one gastrostyle per cyclo-system; pores of the peripheral type; supernumerary dactylopores rare. Gastropore tube single-chambered, but may be constricted by a ring palisade. Dactylostyles usually robust. Ampullae usually internal.

Discussion. As summarized by Cairns (1983b), species attributed to this genus have been placed in the genera *Allopora*, *Stylaster*, and as a subgenus of *Stylaster*. *Stylantheca papillosa* (Dall, 1884) is indeed very similar to *Stylaster* (Group A, i.e., *Allopora*). At present the only characters differentiating *Stylantheca* from *Stylaster* (Group A) is its tendency to form encrusting colonies, and its tendency to have more than one gastrostyle per cyclo-system, although there are exceptions to both of those criteria. Deeper water colonies of *S. papillosa* often have short stubby branches, and the northern range of *S. papillosa* sometimes has only one gastrostyle per cyclo-system (see Remarks of *S. papillosa*). It is likely that *Stylantheca* belongs to the *Stylaster* genus, perhaps as a fifth group, i.e., “Group E”, of species sensu Cairns (1983b) that would include encrusting species with a tendency to

have multiple gastrostyles per cyclostyles. But until more specimens have been collected to study morphological variation, we choose to retain the genus *Stylanthea*.

Type species. *Stylanthea porphyra* Fisher, 1931, by monotypy.

Distribution. From the Alaska Peninsula to Lower California, 0–27 m.

***Stylanthea papillosa* (Dall, 1884)**

http://species-id.net/wiki/Stylanthea_papillosa

Fig 27D–G, 29A–L

Allopora papillosa Dall, 1884: 113–114.—Fisher 1938: 527–528, pl. 54, fig. 4, pl. 59, fig. 3.

Stylanthea porphyra Fisher, 1931: 395–397, pl. 15, figs 1, 1a, pl. 16, figs 5, 5a–b, pl. 17, figs 6, 6a–c.—Boschma 1956: F100, text fig. 81–1a–b; 1960: 426–427, text figs 1e–g.—Cairns 1983b: 430, 481–483, figs 18A–I, 24H, 27G, J.—Wing and Barnard 2004: 10, 27.—Heifetz et al. 2005: 133 (listed).—Stone and Shotwell 2007: 108 (listed).—Whitmire and Clarke 2007: 154 (listed).—Jamieson et al. 2007: 224 (listed).

Allopora petrograpta Fisher, 1938: 530–531, pl. 54, figs 5, 5a, pl. 59, fig. 4.—Frichman 1974: 245–258 (larval development).

Allopora porphyra: Fisher 1938: 528–530, pl. 59, figs 1–2, pl. 60, pl. 61, figs 1, 1a, pl. 70, figs 2, 2a.—Thompson and Chow 1955: 30 (mineralogy).

Stylaster (Allopora) porphyrus: Broch 1942: 102.

Stylaster (Stylanthea) porphyra: Boschma 1951: 39, text fig. 5b.

Stylanthea petrograpta: Cairns 1983b: 430.—Wing and Barnard 2004: 10, 27.—Heifetz et al. 2005: 133 (listed).—Stone and Shotwell 2007: 108 (listed).—Whitmire and Clarke 2007: 154 (listed).—Jamieson 2007: 224 (listed).—Lindner et al. 2008: 3, and supplemental Table 1: 2 (phylogeny and DNA sequences).

Stylanthea papillosa: Cairns 1983b: 430.—Wing and Barnard 2004: 10, 27.—Heifetz et al. 2005: 133 (listed).—Stone and Shotwell 2007: 108 (listed).—Jamieson et al. 2007: 224 (listed).

Stylaster porphyra: Jamieson et al. 2007: 224 (listed).

Type material. *Allopora papillosa*: holotype, a small (11 mm long) dry encrusting fragment, USNM 6852 (Fig. 27D).

Type locality. Coal Harbor, Unga Island, Shumagin Islands, Alaska Peninsula, 11 m.

Stylanthea porphyrea: holotype (USNM 43018, Fig. 27G) and 13 paratype colonies (USNM 43019, 43276, 43277 and SEM stub 136), all dry.

Type locality. Pescadero Point, Carmel Bay, California (36°33'42"N, 121°57'13"W), intertidal.

Allopora petrograpta: two small (15 and 17 mm in length) dry, male encrusting fragments, syntypes, and SEM stub 1506, USNM 43272 (Fig. 27F). **Type locality.** Kyack Island, mouth of Sitka Harbor, Alaska, intertidal.




Figure 27. Color figures of skeleton of various Alaskan stylasterids: **A–C** *Stylaster repandus* **D–G** *Stylaster papillosus* **H–I** *Crypthelia trophostega*: **A–B** lateral and apical views of holotype, USNM 1122740 **C** small paratype colony, USNM 1122741 **D** holotype of *Allopورا papillosa*, USNM 6852 **E** encrusting colony, AB02–0005 **F** syntype of *Allopورا petrograpta*, USNM 43272 **G** holotype of *S. porphyrea*, USNM 43018 **H** holotype, USNM 42876 **I** arborescent colony, USNM 1122435.

Material examined. Types of the three named species; Middle of Cross Sound, AK, 15–21 m, 27 Jul 1978, 5 female colonies, AB78–120; South of Yasha Island, Chatham Strait, AK, depth unknown, 31 Jul 1976, 8 female colonies in alcohol, and SEM stubs 1532–34, AB76–55; Wooden Island, AK, depth unknown, 2002, 1 indet., AB02–150; Wooden Island, AK, depth unknown, 15 May 2009, 1 male in alcohol, AB09–10; Race Rocks, Victoria, British Columbia, depth unknown, 2002, 8 colonies dry and in alcohol, USNM 1073478; Seontary Island, Victoria, BC, 10 m, Jul 2002, 1 large male colony, USNM 1096626; 48°18'N, 123°31.8'W, 12.1 m, 6 Sep 1973, 1 female colony, USNM 76559; cactus island Channel, BC, 15 m, 1 male, AB02–0006; Lion's Gate Bridge, Burrard Inlet, British Columbia, 18 m, Jan 2002, 2 indet., AB02–0005; Turn Island, Washington, 0–27 m, 6 Jul 1995, 10 colonies, USNM 1084659; Puget Sound, Washington, 27 m, 2, CAS 117458–59; Squaw Island, Oregon, 20 Jul 1962, 1 female colony, USNM 45685; 43°18'15"N, 124°23'56"W (Charleston, Oregon), intertidal, 2 indet in alcohol, USNM 1086321; Pillar Point, Half Moon Bay, California, intertidal, 18 Feb 1996, 16 colonies, USNM 1084663; Pigeon Point, California, intertidal, 12 Jul 1995, 12 colonies (forma *porphyra*), USNM 1084662.

Description. The typical form is variable in colony shape, those specimens living in a high energy shallow-water environment usually being encrusting, the layer of coenosteum sometimes less than 1 mm thick (Fig. 27E), the gastrostyle base almost resting on the hard substrate. In deeper water, colonies produce short (up to 17 mm), knobby, clavate, cylindrical branches, which originate from the basal encrustation. Only rarely will these branches bifurcate. Although the types of *S. papillosa* and *S. petrograpta* are small, beach-worn fragments (Figs 27D, F), the species produces mats up to 30 cm in diameter. Parasitic spionid polychaetes (*Polydora* ?*allopors* Light, 1970) usually present, forming binary tubes/paired burrows throughout the coenosteum. Coenosteum reticulate-granular in texture, the strips about 50–55 µm in width, separate by slits 12–15 µm wide. Short, porous, conical papillae (nematopores?) common on coenosteum, each about 0.13 mm in diameter and equally tall (Fig. 29B, D, F–G). Coenosteum purple, pink, red, and occasionally white, the tips of the clavate branches usually white.

Cyclosystems uniformly arranged on encrustations and on all sides of branchlets; diameter of cyclosystems 0.9–1.2 mm. Gastropore tube highly constricted, the upper larger portion being infundibuliform to spherical, the lower chamber spherical and almost completely occupied by the gastrostyle(s), the two portions of the chamber constricted near gastrostyle tip. Gastropore tube diameter about 0.3 mm, the tube constriction about half that diameter. Just above the constriction numerous small (up to 45 µm in length) papillae form a wide ring palisade (Fig. 29K) further reducing access to lower chamber. Gastrostyles quite variable and sometimes irregular in shape, including globose (Fig. 29K), lanceolate, or triangular. Regardless, gastrostyles bear quite long (up to 50 µm long and 15 µm in diameter) spines that completely obscure the underlying gastrostyle shaft. Usually there is only one gastrostyle per cyclosystem, but occasionally the upper chambers of two cyclosystems are linked together resulting in two gastrostyles in one cavity, and rarely there appears to be 2 gastrostyles in one normal gastropore chamber.




Figure 28. *Stylander repandus*, **A–K, M** holotype (male), USNM 1122740, **L** female paratype, USNM 1122739: **A–C** cyclosystems and coenosteal papillae **D** gastrostyle and robust dactylostyle **E** stereo view of a cyclosystem **F–G** coenosteal texture **H** stereo view of a gastrostyle and adjacent dactylostyle **I** stereo view of plate cross section showing some gastrostyles and male ampulla **J** a robust dactylostyle **K** elements of a dactylostyle **L** cross section of a female ampulla **M** cross section of a male ampulla.

In well-preserved colonies, dactylopores are slightly raised above surface of coenosteum; dactylopores 0.10–0.12 mm wide. Number of dactylopores per cyclo-system variable, depending on size of cyclo-system, but ranges from 2–12, with common modes of 4, 5 and 8 (see Remarks). Dactylostyles robust (Fig. 29A, C, H), the cylindrical elements up to 55 μm long and 13–17 μm in diameter, not unlike the gastrostyle elements.

Female ampullae (Fig. 29L) primarily internal, with little superficial relief, even by an efferent pore; internal diameter 0.7–0.9 mm. Male ampullae also internal, the internal diameter being 0.4–0.5 mm.

Remarks. Although the holotype of *Allopora papillosa* is a small, worn colony measuring only 9.4 \times 6.6 mm in width and containing only 28 cyclo-systems (Fig. 27D), it is not difficult to see the similarity to the two syntypes of *Allopora petrograpta* Fisher, 1938 (Fig. 27F), also worn specimens (13 and 16 mm in width, constituting over 100 cyclo-systems), even though they were collected from shallow water on opposite sides of the Gulf of Alaska. All of the cyclo-systems of the type of *Stylantheca papillosa* have only one gastrostyle, whereas all but four of the cyclo-systems of the syntypes of *S. petrograpta* have one gastrostyle, the other four having two. The more difficult synonymy is with the more southerly species *S. porphyra* Fisher, 1931, described from shallow water in Carmel Bay, California. *S. porphyra* is very similar to the encrusting forms of *S. papillosa*, differing only in usually having more than one gastrostyle per cyclo-system, sometimes as many as 8. Cairns (1983b) reported an average of 3.3 gastrostyles per cyclo-system (mode = 3) for the holotype. Another typical population of *S. porphyra* is reported herein from Monterey Bay (USNM 1084662), also having multiple gastrostyles per cyclo-system, but just 50 km to the north (USNM 1084663) are colonies otherwise identical to *S. porphyra* that have only one gastrostyle per cyclo-system. Fisher (1938:530) was uncertain as to the systematic importance of the number of gastrostyles per cyclo-systems, suggesting that this character might have subgeneric or even generic discriminating power. We suggest that it is a matter of intraspecific variation and thus synonymize *S. porphyra* with *S. papillosa*, resulting in only one valid species in the genus *Stylantheca*. The number of dactylopores per cyclo-system of *S. porphyra* ranges from 6 to 16 ($n = 34$, average = 9.11, $\sigma = 2.08$, mode = 10) (Cairns, 1983b), whereas it is lower for the type of *S. petrograpta* ($n = 30$, range = 2–7, average = 5.07, $\sigma = 0.94$, mode = 5) and slightly lower still for the types of *S. papillosa* ($n = 14$, range = 3–5, average and mode = 4.0, $\sigma = 0.68$). Another typical specimen (AB76–55) having only 1 gastrostyle per cyclo-system has a range of 5–12 dactylopores per cyclo-system ($n = 30$, average = 7.43, $\sigma = 1.61$, mode = 8). The higher number of dactylopores per cyclo-system of *S. porphyra* is explained by its larger cyclo-systems that house multiple gastrostyles, but in general the number of dactylopores per cyclo-system cannot be used as a diagnostic character for this species. Southern populations that have a slightly larger cyclo-systems and more than one gastrostyle per cyclo-system can be referred to as the “*porphyra*” form, a conclusion first suggested by Cairns (1983b: 483). The type of *S. porphyra* is illustrated by Cairns (1983b) and as Fig. 27G herein.




Figure 29. *Stylianteca papillosa*, **A–G, K** topotypic specimen, USNM 43276, **H–J** syntype of *Allopore petrograpta*, USNM 43272, **L** female colony, AB76–55: **A** stereo view of a cyclostyle **B, D, F** coenosteal papillae (nematopores) **C** robust dactylostyle **E** supernumerary dactylopore **G** three cyclostyles and many papillae **H** stereo view of longitudinal section through cyclostyle **I** gastrostyle **J** gastrostyle spines **K** stereo view of squat gastrostyle and ring palisade **L** stereo view of longitudinal section of two cyclostyles and a female ampulla.

Two other eastern Pacific species are similar to *Stylanthea papillosa* in color and cyclo-system and gastrostyle shape: *Stylaster venustus* (Verrill, 1870) and *Stylaster californicus* (Verrill, 1866). *S. venustus* (Figs 16A–I, 17F–G) is known from Monterey Bay, California (USA) to Vancouver Island, BC (Canada) at depths of 10–108 m and, although it may originate with an encrusting base (Fig. 17F, MCZ 5525, holotype), it quickly forms small colonies with delicate branches (Fig. 17G). Sixteen locality records are present at the NMNH, most from off Washington and Oregon (USA), but this species is not treated in this review except to illustrate some of its characters (Table 2). On the other hand, the more southern *S. californicus*, known from off southern California (USA) from approximately 21–45 m (see Boschma 1957), forms large colonies with robust branches; this species is not further discussed in this review except for the table of comparisons (Table 2).

Distribution. Widespread from the Alaska Peninsula (Shumagin Islands) to Monterey Bay, California, although not yet reported from the northern Gulf of Alaska. Common in the inner passages of Alaska and British Columbia; intertidal to 27 m. Forma *porphyra* known only from Monterey Bay area.

Genus *Crypthelia* Milne Edwards & Haime, 1849

<http://species-id.net/wiki/Crypthelia>

Diagnosis. Colonies usually uniplanar and quite delicate. Coenosteum linear-imbri-cate and white; nematopores common. Gastro- and dactylo-pores arranged in cyclo-systems that are arranged unifacially on colony, only rarely on both faces; cyclo-systems covered all or in part by a fixed lid. Gastropore tube double-chambered, lacking gastro-and dactylostyles. Ampullae usually superficial with a variety of locations of efferent pores.

Discussion. *Crypthelia* is the second-most species-rich genus in the family, currently boasting 31 Recent and one fossil species (Appeltans et al. 2011: www.marine-species.org). They are easily recognized by the prominent fixed lid that covers each cyclo-system. They are more characteristic of deep water environment and hold the record for the deepest stylasterid species.

Type species. *Crypthelia pudica* Milne Edwards and Haime, 1849, by monotypy.

Distribution. Eocene to Recent: cosmopolitan, but not off continental Antarctica, 183–2789 m.

***Crypthelia trophostega* Fisher, 1938**

http://species-id.net/wiki/Crypthelia_trophostega

Figs 27H–I, 30A–L

Crypthelia trophostega Fisher, 1938: 533–535, pl. 62, figs 1–8, pl. 63.—Cairns 1986: 24 (ampulla code); 1991: 386 (off Pribilof Island).—Cairns and Macintyre 1992:

102–103 (mineralogy).—Heifetz 2002: 22 (listed).—Wing and Barnard 2004: 10, 27, fig. 29.—Heifetz et al. 2005: 133, 136 (listed).—Stone and Shotwell 2007: 107 (listed).—Brooke and Stone 2007: 530, figs 2L, 3F.—Jameison et al. 2007: 224 (listed).—Lindner et al. 2008: 3, and supplemental Table S1: 1 (phylogeny and DNA sequences).

Cryptohelia trophostega: Thompson and Chow 1955: 30 (mineralogy).

Type material. Holotype: *Alb*-3480, 1 large dry male colony, 15 cm in width, SEM stubs 1482–1484, USNM 42876 (Fig. 27H). Paratypes: *Alb*-3480, 2 dry colonies, male and female, USNM 52264; *Alb*-3480, 1 female colony in alcohol, USNM 43769; *Alb*-3480, 1 dry colony, CAS 69609.

Type locality. *Alb*-3480: 52°06'N, 171°45'W (Amukta Pass), 517 m.

Material examined. *Alaskan Leader* 40, 52°09'18"N, 175°40'42"W, 174 m, 8 Jun 2002, 1 indet., USNM 1137354; *Alaskan Leader* 54–14, 51°44.4'N, 178°16.7'W, 567–680 m, 11 Jun 2002, 1 indet., AB02–29; *Alb*-4771, 54°30'N, 179°17'E, 779 m, 4 Jun 1906, 2 males, USNM 76780; *Alb*-4780, 52°01'N, 174°39'E, 1913 m, 7 Jun 1906, 2 female, USNM 62371; *Alb*-4781, 52°14'30"N, 174°13'E, 882 m, 7 Jun 1906, 3 male, USNM 43770; *Delta* 6230–20–19, 52°28.142'N, 173°35.882'W, 190 m, 8 Jul 2004, 1 female, AB; *Dominator* 971–142, 51°43'10"N, 178°35'E, 215 m, 17 Jul 1997, 1 male, USNM 1123356; *Dominator* 971–181, 51°27'43"N, 178°35'E, 384 m, 27 Jul 1997, 1 indet., USNM 1123354; *Dominator* 971–201, 51°54'28"N, 378 m, 1 Aug 1997, 1 male, USNM 1123355; *Dominator* 971–126, 51°34'16"N, 177°47'36"W, 237 m, 12 Jul 1997, 1 indet. in alcohol, USNM 1123353; *Jason* II-2095–2-9–5, 51°48.682'N, 173°50.070'W, 843 m, 26 Jul 2004, 1 male, AB08–0035; *Jason* II-2099–17–1, 51°30.101'N, 177°02.354'W, 1453 m, 30 Jul 2004, 1 indet. in alcohol, AB09–0029; *Ocean Olympic*, 52°22.5'N, 176°03.5'E, 237 m, 1 female, AB 00–0024; *Patricia Lee*, 51°59'N, 179°30.08'E, 457 m, 1 male, AB00–0042; *Sea Storm* 93, 51°50'59"N, 178°26'02"E, 390 m, 5 Jul 2002, 1 male in alcohol, USNM 1122897; *Sea Storm* 108, 52°11'32"N, 175°28'18"E, 208 m, 8 Jul 2002, 1 male, USNM 1122893; *Sea Storm* 109, 52°17'16"N, 175°20'56"E, 238 m, 8 Jul 2002, 1 female in alcohol, USNM 1122894; *Sea Storm* 133, 52°13'40"N, 176°02'19"E, 148 m, 15 Jul 2002, 1 female and 1 male in alcohol, USNM 1122889–90; *Sea Storm* 138, 52°13'50"N, 175.242'E, 146 m, 16 Jul 2002, 1 female and 1 male, USNM 1122891–92; *Sea Storm* 148, 52°28'16"N, 173°19'10"W, 194 m, 2 female, SEM stub 1485, USNM 1122887–88; *Sea Storm* 150, 52°30'47"N, 173.2935°W, 220 m, 21 Jul 2002, 1 female, USNM 1075952; *Sea Storm* 151, 52°33'40"N, 173.3195°W, 203 m, 21 Jul 2002, 1 male, USNM 1122895; *Sea Storm* 155, 52°38'43"N, 172°16.38'W, 393–401 m, 22 Jul 2002, 1 indet., USNM 1122896; *Shishaldin*, 54°12'37"N, 179°30'05"E, 179 m, 1 female and 1 male, USNM 1122435 and 1122446; *Shishaldin*, 54°07'N, 179°45'E, 366 m, 20 Feb 2000, 1 female, SEM stubs 1486–87, USNM 1122487; *Vesteraalen* 941–138, 51°28'N, 178°38'W, 0–311 m, 7 Jul 1994, 1 male, USNM 96248; *Vesteraalen* 941–167, 51°54'N, 178°20'E, 0–150 m, 19 Jul 1994, 1 female, USNM 96240; 51°50'32"N, 176°00'08"E, 272 m, 5 Dec 2000, 1 male, USNM 1122441;




Figure 30. *Cryphelia trophostega*, **A–B, F, H–J** female colony, USNM 1122887 **C–E, G, K–L** holotype (male), USNM 42876: **A–B** cyclosystem and lid, covered with numerous nematopores **C** under side of cyclosystem lid showing numerous male efferent pores **D–E** coenosteal texture **F** slender pseudosepta bearing nematopores **G** three nematopores **H** stereo view of longitudinal section of cyclosystem showing multiple female efferent pores under lid **I** enlargement of efferent pores of figure H **J** fractured female lid **K** stereo view of fractured cyclosystem showing multiple male efferent pores (enlarged in fig. C) **L** stereo view of cyclosystem missing its lid revealing several male ampullae.

“Gulf of Alaska”, 219–274 m, 1 female, USNM 77415; “Bering Sea”, 1 male in alcohol, USNM 76570; University of Washington, 51°32'N, 179°15'W, 278–289 m, 1 Sep 1968, 1 female, USNM 62372.

Description. Colonies variable in shape, usually uniplanar (Fig. 27H) but sometimes forming three dimensional bushes (Fig. 27I) or multiplanar colonies. Largest colony known (USNM 1122446) is multiplanar, 20 cm tall and 12 cm wide, with a basal diameter of 11.5 mm, this colony only slightly larger than the holotype. Branching irregularly dichotomous and often anastomotic, which reinforces strength of a uniplanar colony. Coenosteum linear-imbricate in surface texture, the parallel strips 50–70 μm wide. Strips covered with irregularly shaped platelets, which are rarely continuous across a strip but dissected into 2–5 smaller sections, each of which has an irregular leading edge and only a slight imbricating overlap with more distal platelet, altogether producing a rough microtexture. Shallow, circular (0.06–0.13 mm in diameter) nematopores very common, occurring on branch coenosteum, cyclosystem lids, and even on pseudosepta (Fig. 30A–C, F–G, J), most concentrated around cyclosystem edge. Nematopores flush with coenenchyme or encircled with a very low rim. Coenosteum white.

Cyclosystems unilinearly arranged either bifacially or on one face, both conditions sometimes occurring on same colony. Unifacially arranged cyclosystems seem to be more common on uniplanar colonies, whereas the bifacial arrangement favors bushy coralla. Cyclosystems circular to slightly elliptical in outline (2.2–2.6 mm in diameter), the greater axis of ellipse being perpendicular to branch axis. Cyclosystems slightly flared, each covered with a horizontal lid that covers most of cyclosystem, sometimes even fusing to opposite side of cyclosystem. Lids usually greatly inflated, containing the male or female ampullae. Cyclosystems have a range of 13–23 dactylopores ($n = 50$, average = 18.66 ($\sigma = 1.96$), and mode = 19). Upper gastropore chamber spherical, about 0.9–1.0 mm in diameter, which is separated from lower, flattened chamber by a circular gastropore ring constriction 0.4–0.6 mm in diameter, the lower chamber being about 0.8–0.9 mm in diameter (Fig. 30H). Dactylopores uniform in width, 0.14–0.16 mm; pseudosepta quite slender, having a thin (20–60 μm) blade-like inner edge (Fig. 30F). Upper edges of pseudosepta not exsert, but gradually attenuate in size near cyclosystem edge.

Female ampullae (Fig. 30H–J) smooth, massive, hemispherical swellings in cyclosystem lid, often two ampullae contained in same lid (Fig. 30I). Efferent pores quite large (up to 0.5 mm in diameter), opening beneath lid (type A ampulla formula of Cairns, 1986, Fig. 30H–I). Male ampullae (Fig. 30K–L) irregularly shaped swellings in cyclosystem lid, up to 11 occurring in one lid. Efferent pores smaller (10–12 μm in diameter), also opening beneath the lid (type A2 ampulla formula of Cairns 1986)(Fig. 30C, K).

Remarks. Among the 31 Recent species of *Cryptothelia* (Cairns et al. 1999; www.marinespecies.org), *C. trophostega* is unique in having a tendency to have a bifacial cyclosystem arrangement and for having more than one female ampullae per cyclosystem lid. Furthermore, only one other species is known to have the A–A2 ampulla formula (see Cairns, 1986), that being *C. pudica* Milne Edwards and Haime, 1849.

C. trophostega differs from *C. pudica* in having bifacial cyclo systems, a rougher coenosteal texture, more nematopores, larger cyclo systems, and lower lids. The corallum was found to be 100% aragonitic according to Cairns and Macintyre (1992). Of the 42 specimens examined: 18 are female, 18 male, and 6 indeterminate in gender.

Distribution. Aleutian Islands from Near Islands to Amuka Pass, Petrel and Bowers Banks, off Pribilof Bank; 146–1913 m, although most records between 200 and 400 m.

Acknowledgements

We are grateful to the following people who generously loaned or gave us specimens to study: Anita Brinckmann-Voss, Arthur Schultz, Garry Fletcher, Steve Blair, ‘Skip’ Zenger, Jay Orr, Dale Calder, Bob van Syoc, Bob Stone, Bruce Wing, Kirsten Lindstrom, Mary Sue Brancato, Adam Baldinger, and John Pearse. We are indebted to Tim Coffey for constructing most of the figures and to Walter Larrimore for taking the color photographs. This work was partially funded by a National Science Foundation grant (NSF PEET DEB-9978086) and by a scholarship from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES BEX0917-0) to A. Lindner. We also thank the reviewers for comments that improved the manuscript.

References

- Appeltans W, Bouchet P, Boxshall GA, Fauchald K, Gordon DP, Hoeksma BW, Poore GCB, van Soest RWM, Stohr S, Walter TC, Costello MJ (2001) World Register of Marine Species. www.marinespecies.org [Accessed on 2011.09.19]
- Avisé JC (2000) Phylogeography: the history and formation of species, second edition. Harvard University Press, Cambridge, 447 pp.
- Baum DA, Shaw KL (1995) Genealogical perspectives on the species problem. In: Hoch PC, Stephenson AG (Eds) Experimental and Molecular Approaches to Plant Biosystematics, Missouri Botanical Garden, St Louis, 289–303.
- Bernecker M, Weidlich O (1990) The Danian (Paleocene) coral limestone of Fakse, Denmark: a model for ancient aphotic, azooxanthellate coral mounds. *Facies* 22: 103–138. doi: 10.1007/BF02536947
- Bernecker M, Weidlich O (2005) Azooxanthellate corals in the Late Maastrichtian – Early Paleocene of the Danish basin: bryozoan and coral mounds in a boreal shelf setting. In: Freiwald A, Roberts JM (Eds) Cold-water Corals and Ecosystems. Springer-Verlag, Berlin, 3–25. doi: 10.1007/3-540-27673-4_1
- Boschma H (1956) Milleporina and Stylasterina. In: Moore RC (Ed) Treatise on Invertebrate Paleontology, Part F, University of Kansas Press, Lawrence, Kansas, F90-F106.
- Boschma H (1957) List of the described species of the order Stylasterina. *Zoologische Verhandlungen* 33: 1–72.

- Boschma H (1959) Revision of the Indo-Pacific species of the genus *Distichopora*. *Bijdragen tot de Dierkunde* 29: 121–171.
- Boschma H (1960) *Gyropora africana*, a new stylasterine coral. *Proceedings Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam. Series C* 63(4): 423–434.
- Broch H (1935) Einige Stylasteriden (Hydrokorallen) der Ochotskischen und Japanischen See. *Explorations des mers d' URSS* 22: 58–60.
- Broch H (1936) Untersuchungen an Stylasteriden (Hydrokorallen). Teil 1. *Skrifter utgitt av Norske Videnskaps-Akademi i Oslo. 1. Matematisk-Naturvidenskapelig Klasse* 8: 1–103.
- Broch H (1942) Investigations of Stylasteridae (Hydrocorals). *Skrifter utgitt av Norske Videnskaps-Akademi I Oslo. 1. Matematisk-Naturvidenskapelig Klasse* 3: 1–113.
- Broch H (1951) Stylasteridae (Hydrocorals) from the Southern Ocean. *Discovery Reports* 26: 33–46.
- Brooke S, Stone R (2007) Reproduction of deep-water hydrocorals (family Stylasteridae) from the Aleutian Islands, Alaska. *Bulletin of Marine Science* 81(3): 519–532.
- Cairns SD (1983a) Antarctic and Subantarctic Stylasterina (Coelenterata: Hydrozoa). *Antarctic Research Series* 38: 61–163. doi: 10.1029/AR038p0061
- Cairns SD (1983b) A generic revision of the Stylasterina (Coelenterata: Hydrozoa). Part 1. Description of the genera. *Bulletin of Marine Science* 33(2): 427–508.
- Cairns SD (1986) Stylasteridae (Hydrozoa: Hydroida) of the Galápagos Islands. *Smithsonian Contributions to Zoology* 426: 1–42. doi: 10.5479/si.00810282.418
- Cairns SD (1991) *Cyclohelia lamellata*, new genus and species of Stylasteridae (Cnidaria: Hydrozoa) from the Bering Sea. *Pacific Science* 45(4): 383–388.
- Cairns SD (1992) A generic revision of the Stylasteridae (Coelenterata: Hydrozoa). Part 3. Keys to the genera. *Bulletin of Marine Science* 49(1–2): 538–545.
- Cairns SD (2005) Revision of the Hawaiian Stylasteridae (Cnidaria: Hydrozoa: Athecata). *Pacific Science* 59(3): 439–451. doi: 10.1353/psc.2005.0034
- Cairns SD (2011) Global diversity of the Stylasteridae (Cnidaria: Hydrozoa: Athecata). *PLoS ONE* 6(7): 1–13. doi: 10.1371/journal.pone.0021670
- Cairns SD, Hoeksema BW, Land J van der (1999) Appendix: list of extant stony corals. *Atoll Research Bulletin* 459: 13–46.
- Cairns SD, Macintyre IG (1992) Phylogenetic implications of the calcium carbonate mineralogy in the Stylasteridae (Cnidaria: Hydrozoa). *Palaios* 7: 96–107. doi: 10.2307/3514799
- Cunningham CW, Buss LW (1993) Molecular evidence for multiple episodes of pedomorphosis in the family Hydractiniidae. *Biochemical Systematics and Ecology* 21(1): 57–69. doi: 10.1016/0305-1978(93)90009-G
- Dall WH (1884) On some Hydrocorallinae from Alaska and California. *Proceedings of the Biological Society of Washington* 2: 111–115.
- Eguchi M (1968) The Hydrocorals of Sagami Bay. In: Eguchi M (Ed) *The Hydrocorals and Scleractinian Corals of Sagami Bay*, Maruzen Co., Tokyo, 1–53.
- Fisher WK (1931) Californian hydrocorals. *The Annals and Magazine of Natural History* 8: 391–399. doi: 10.1080/00222933108673409
- Fisher WK (1938) Hydrocorals of the North Pacific Ocean. *Proceedings of the United States National Museum* 84(3024): 493–554. doi: 10.5479/si.00963801.84-3024.493

- Frichman HK (1974) The planula of the stylasterine hydrocoral *Allopora californica* Fisher: its structure, metamorphosis and development of the primary cyclosystem. *Proceedings of the Second International Coral Reef Symposium* 2: 245–258.
- Gouinotte JM, Orr J, Cairns SD, Freiwald A, Morgan L, George RY (2006) Will human-induced changes in seawater chemistry alter the distribution of deep-sea scleractinian corals? *Frontiers in Ecology and the Environment* 3(4): 141–146. doi: 10.1890/1540-9295(2006)004[0141:WHCISC]2.0.CO;2
- Gray JE (1831) Description of a new genus (*Stylaster*) of star-bearing corals. *Zoological Miscellany*, 36–37.
- Gray JE (1847) An outline of the arrangement of stony corals. *The Annals and Magazine of Natural History* (1)19: 120–128. doi: 10.1080/037454809496460
- Heifetz J (2002) Coral in Alaska: distribution, abundance, and species associations. *Hydrobiologia* 471: 19–28. doi: 10.1023/A:1016528631593
- Heifetz J, Wing BL, Stone RP, Malecha PW, Courtney D (2005) Corals of the Aleutian Islands. *Fisheries Oceanography* 14: 131–138. doi: 10.1111/j.1365-2419.2005.00371.x
- International Commission on Zoological Nomenclature (1999) *International Code of Zoological Nomenclature*, fourth edition. International Trust for Zoological Nomenclature, London, 306 pp.
- Jamieson GS, Pellegrin N, Jesson S (2007) Taxonomy and zoogeography of cold-water corals in coastal British Columbia. In: George RY, Cairns SD (Eds) *Conservation and Adaptive Management of Seamount and Deep-Sea Coral Ecosystems*, University of Miami, Miami, 215–229.
- Lamarck JBPA de (1816) *Histoire Naturelle des Animaux sans Vertèbres*, volume 2. Verdier, Paris, 568 pp.
- Light WJ (1970) *Polydora alloporis*, sp. n., a commensal spionid (Annelida, Polychaeta) from a hydrocoral off Central California. *Proceedings of the California Academy of Sciences* 37(14): 459–472.
- Lindner A (2005) *Evolution and Taxonomy of Stylasterid Corals (Cnidaria: Hydrozoa: Stylasteridae)*. PhD thesis, Chapel Hill, North Carolina, Duke University, 165 pp.
- Lindner A, Cairns SD, Cunningham CW (2008) From offshore to onshore: multiple origins of shallow-water corals from deep-sea ancestors. *PLoS One* 3(6): e2429. doi:10.1371/journal.pone.0002429
- Lowe TP (1967) *A Comparative Study of Stony Hydroids, Order Stylasterina, of Southern South America and Adjacent Antarctic Regions*. MS thesis, Washington DC, USA, George Washington University, Washington D.C., 116 pp.
- Lowenstam HA (1964) Coexisting calcites and aragonites from skeletal carbonates of marine organisms and their strontium and magnesium contents. In: Miyake Y, Koyama T (Eds) *Recent Researches in the Fields of Hydrosphere, Atmosphere, and Nuclear Geochemistry*, Maruzen C., Ltd, Tokyo, 373–403.
- Milne Edwards H, Haime J (1849) Mémoire sur les polypiers appartenant a la famille des Oculinides, au groupe intermédiaires des Pseudoastréides et a la famille des Fongides. *Compte Rendu Hebdomadaire des Séances de l'Académie des Sciences, Paris*, 29: 67–73.
- Milne Edwards H, Haime J (1850) Introduction In: *A Monograph of the British Fossil Corals. Part 1*. Paleontographical Society, London, i-lxxxv.

- Mori K (1980) Professor Motoki Eguchi, April 21, 1905 – March 4, 1978: obituary, bibliography and check-list of corals. *Scientific Reports of the Tohoku University (Geology)* 50 (1/2): 1–18.
- Namikawa H, Mawatari SF (1992) Role of the tentaculozooids of the polymorphic hydroid *Stylactaria conchicola* (Yamada) in interactions with some epifaunal space competitors. *Journal of Experimental Marine Biology* 62: 65–75. doi: 10.1016/0022-0981(92)90125-T
- Naumov DV (1960) Hydroids and Hydromedusae of the USSR. Keys to the fauna of the USSR published by the Academy of Sciences of the USSR 70: 1–660.
- Stone RP, Shotwell SK (2007) State of deep coral ecosystems in the Alaskan region: Gulf of Alaska, Bering Sea and the Aleutian Islands. In: Hourigan TF et al. (Eds) *The State of Deep Coral Ecosystems of the United States*. NOAA Technical Memorandum CRCP-3, NOAA, Silver Spring, 65–108.
- Swofford DL (1999) PAUP*. Phylogenetic analysis using parsimony, version 4.0. Sinauer Associates, Sunderland, Massachusetts.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* 24: 4876–4882. doi: 10.1093/nar/25.24.4876
- Thompson TG, Chow TJ (1955) The strontium-calcium atom ratio in carbonate-secreting marine organisms. *Deep-Sea Research (Supplement)* 3: 20–39.
- Verrill AE (1866) Synopsis of the polyps and corals of the North Pacific Exploring Expedition, with descriptions of some additional species from the west coast of North America. Part III. Madreporaria. *Proceedings of the Essex Institute* 5: 17–50.
- Verrill AE (1868–1870) Review of the corals and polyps of the west coast of America. *Transactions of the Connecticut Academy of Arts and Sciences* (2): 377–558.
- Whitmire CE, Clarke ME (2007) State of deep coral ecosystems in the U.S. Pacific coast: California to Washington. In Hourigan TF et al. (Eds) *The State of Deep Coral Ecosystems of the United States*. NOAA Technical Memorandum CRCP-3, NOAA, Silver Spring, 109–154.
- Wing GL, Barnard DR (2004) A field guide to Alaskan Corals. NOAA Technical Memorandum NMFS-AFSC-146, NOAA, US Dept of Commerce, 67 pp.

New records of water mites of the family Sperchontidae Thor, 1900 from China (Acari, Hydrachnidia), with descriptions of two new species

Xu Zhang^{1,2†}, Dao-Chao Jin^{2,‡}

1 The Provincial Key Laboratory of Agricultural Pest Management in Mountainous Region, Institute of Entomology, Guizhou University; Guiyang 550025, China **2** Anhui Key Laboratory of Plant Resources and Biology, Huaibei Normal University; Huaibei 235000, China

† [urn:lsid:zoobank.org:author:CF3C3580-FF77-44BC-98FB-477999A7AD4E](https://doi.org/urn:lsid:zoobank.org:author:CF3C3580-FF77-44BC-98FB-477999A7AD4E)

‡ [urn:lsid:zoobank.org:author:FC06EF2C-1B36-4BFB-9E33-D9AF34725F40](https://doi.org/urn:lsid:zoobank.org:author:FC06EF2C-1B36-4BFB-9E33-D9AF34725F40)

Corresponding author: Dao-Chao Jin (dcjin@gzu.edu.cn)

Academic editor: V. Pešić | Received 26 August 2011 | Accepted 17 November 2011 | Published 22 December 2011

[urn:lsid:zoobank.org:pub:9C3BD933-1D7E-4158-9D60-B1D32FC3157B](https://doi.org/urn:lsid:zoobank.org:pub:9C3BD933-1D7E-4158-9D60-B1D32FC3157B)

Citation: Zhang X, Jin D-C (2011) New records of water mites of the family Sperchontidae Thor, 1900 from China (Acari, Hydrachnidia), with descriptions of two new species. ZooKeys 158: 89–109. doi: 10.3897/zookeys.158.1970

Abstract

Five species of water mites of the family Sperchontidae Thor, 1900 are reported from China. Two of them are new to science, *Sperchon* (*Sperchon*) *orbipatella* **sp. n.** and *S.* (*Sperchon*) *urumqiensis* **sp. n.**, and the other three are new to China, i.e., *Sperchon* (*Palpisperchon*) *nikkoensis* Imamura, 1976, *S.* (*Sperchon*) *sounkyo* Imamura, 1954 and *Sperchonopsis* (*Sperchonopsella*) *whiteshellensis* Conroy, 1991. The first descriptions of the female of *S. nikkoensis* and the male of *S. sounkyo* are also given. The subgenus *Sperchonopsella* Conroy is new to the fauna of China.

Keywords

Hydrachnidia, Sperchontidae, new species, new records, China

Introduction

Water mites of the family Spermontidae Thor, 1900 are presently known from all biogeographic regions, except Antarctica, and over 200 species were reported (Cook 1974, Viets 1987, Di Sabatino et al. 2008). Up to now, 17 species were described from China: *Sperchon beijingensis* Zhang & Jin, 2010; *S. brevialpis* Jin, 1997; *S. curvipalpis* Zhang & Jin, 2010; *S. fluvialtilis* Uchida, 1934; *S. garhwalensis* Kumar, Kumar & Pesic, 2007; *S. gracilipalpis* Lundblad, 1941; *S. heteropoda* Zhang & Jin, 2010; *S. huangshanensis* Zhang & Jin, 2010; *S. lanigerus* Guo & Jin, 2011; *S. mirabilis* Lundblad, 1941; *S. oligospinis* Jin, 1997; *S. placoderma* Lundblad, 1967; *S. perspicuus* Zhang & Jin, 2011; *S. plumifer* Thor, 1902; *S. rostratus* Lundblad, 1968; *S. turfanensis* Zhang & Jin, 2010 and *Sperchonopsis echphyoma* Prasad & Cook, 1972 (Zhang and Jin 2010, Jin 1997; Zhang et al. 2010, Jin et al. 2010, Zhang et al. 2007, Zhang et al. 2011).

During checking our collection of water mites from China, five spermontid species were found. Two of them, *Sperchon* (*Sperchon*) *orbipatella* sp. n. and *S. (Sperchon) urumqiensis* sp. n., are new to science. The other three taxa, i.e., *Sperchon* (*Sperchon*) *sounkyo* Imamura, 1954, *S. (Palpisperchon) nikkoensis* Imamura, 1976 and *Sperchonopsis* (*Sperchonopsella*) *whiteshellensis* Conroy, 1991, are new records for China. The subgenus *Sperchonopsella* is new to the fauna of China.

Material and methods

Specimens were collected by Dao-Chao Jin, Jian-Jun Guo, Zhen-Zao Tian, Ai-Ping Cui and Xu Zhang during 1997–2010 from China, and preserved in Koenike's fluid and dissected as described elsewhere (e.g. Cook 1974). Terms follow Jin (1997). The following abbreviations are used:

A1, A2 = antennal glandularia 1 and 2; ACG = anterior coxal group (CxI + CxII); CxI–CxIV = coxae I–IV; D1–D4 = dorsoglandularia 1–4; E1–E4 = epimeroglandularia 1–4; L1–L4 = lateroglandularia 1–4; O1, O2 = ocularia 1 and 2; PCG = posterior coxal group (CxIII + CxIV); P-I–P-V = palpal segments 1–5; V1–V4 = venteroglandularia 1–4; I-L-1–I-L-6 = the first leg segments 1–6; II-L-1–II-L-6 = the second leg segments 1–6; III-L-1–III-L-6 = the third leg segments 1–6; IV-L-1–IV-L-6 = the fourth leg segments 1–6.

All the type specimens are deposited in the Institute of Entomology, Guizhou University, China (GUGC).

All measurements are given in μm with the mean first, followed by the range in bracket.

Systematics

Family Sperchontidae Thor, 1900

Genus *Sperchon* Kramer, 1877

Subgenus *Palpisperchon* Lundblad, 1941

Sperchon (Palpisperchon) nikkoensis Imamura, 1976


http://species-id.net/wiki/Sperchon_nikkoensis

Figures 1–10

Material examined. 2 females, Hainan Province, Bawangling National Nature Reserve, an unnamed stream (19°07'16"N, 109°04'58"E), 15 August 2005, coll. Xu Zhang; 1 male and 2 females, Guizhou Province, Leigongshan National Nature Reserve, an unnamed stream (26°21'06"N, 108°12'39"E), 3 October 2005, coll. Xu Zhang; 2 males and 2 females, Anhui Province, Jinzhai city, Shuanghe country, an unnamed stream (31°36'37"N, 115°41'30"E), 19 July 2010, coll. Xu Zhang; 1 female, Anhui Province, Anqing city, Mingtangshan scenic area, Hulu River (30°51'19"N, 116°06'06"E), 20 August 2010, coll. Xu Zhang.

Description. Male (n = 3): Idiosoma flat, 730 (730-768) in length, 583 (583-616) in width, color yellow-brown. Cuticle soft and covered with small flat papillae and fine striations in various form and size (Fig. 3). Dorsum without chitinous plate, only two pair of muscular sigillae faintly visible. Each glandularia on dorsum and venter encircled by a plate and raised conically as nipples. The anterior dorsal area before A1 protruded dorsally forward and leaf-shaped with a vein-like line in dorsal view. Coxae in four groups, surface of coxae reticulated. ACG 210 (210-218) in length, close to each other but not fused, posterior apodeme indistinct. E2 laterally between ACG and PCG; PCG 213 (213-230) in length, widely separated. Glandularia absent from CxI-II. Distance between anterior end of ACG and posterior end of PCG 438 (438-451). Genital field between CxIV of PCG with a small and rounded platelet in front. Genital valves not covering the genital acetabula, 143 (143-152) in length, 135 (135-140) in width. Three pairs of acetabula, the anterior two elliptic and the posterior more or less rounded. Cuticle with fine striations between the genital organ and coxal groups. V1 without accompanying glandularia but on small sclerites. Excretory pore surrounded by a sclerotized ring and close to the line of V2.


Infracapitulum with a short rostrum, length 292 (292-236). Chelicera total length 352 (352-387), basal segment length 283(283-308), claw length 69 (69-79), basal segment/claw length ratio 4.1 (3.9-4.1). Palp short and thick. Dorsal lengths of the palpal segments: P-I, 35 (35-37); P-II, 114 (114-121); P-III, 48 (48-53); P-IV, 89 (89-98); P-V, 52 (52-59). P-I stout and without seta. P-II thick with a long ventro-distal projection, bearing three setae, one of which nearly at the base of the projection and slightly longer than projection, the other two relatively short located approximately on the middle of projection. Nine seta on the dorsal and lateral side of the P-II. P-III shorter than P-II, with a long and thin ventrodiscal seta and three short dorsal setae. P-IV with two greatly enlarged ventral peg-like setae, close to each other.



Figures 1–3. *Spermion (Palpspermion) nikkoensis*, Imamura, 1976, Male **1** idiosoma, dorsal view **2** idiosoma, ventral view **3** decorations of cuticle.

Dorsal lengths of leg I: I-L-1, 39 (39-48); I-L-2, 72 (72-81); I-L-3, 83 (83-97); I-L-4, 137 (137-151); I-L-5, 156 (156-171); I-L-6, 127 (127-141). Dorsal lengths of leg IV: IV-L-1, 88 (88-97); IV-L-2, 95 (95-110); IV-L-3, 129 (129-145); IV-L-4, 237 (237-255); IV-L-5, 220 (220-236); IV-L-6, 192 (192-208). Ambulacrum with two claws. Claws with well protruded claw-blade and two clawets, a long dorsal and a shorter ventral one (Fig. 8).

Female (n = 3): Similar to male except for the morphology of genital field and the size of idiosoma. Idiosoma 816 (792-857) in length, 688 (643-712) in width. ACG 310 (302-317) in length, PCG 319 (308-336) in length. Distance between anterior end of ACG and posterior end of PCG 606 (587-624). Genital field 234 (226-242) in length, 215 (208-229) in width. Pregenital sclerite crescent-shaped, and more developed than the postgenital sclerite. Infracapitulum length 301 (290-316). Chelicera total length 381 (364-409), basal segment length 308 (293-328), claw length 73 (71-80), basal segment/claw length ratio 4.2 (4.1-4.2). Dorsal lengths of the palpal segments: P-I, 41 (40-47); P-II, 123 (119-128); P-III, 57 (55-62); P-IV, 94 (89-99); P-V, 57 (55-64). Dorsal lengths of the first leg: I-L-1, 54 (52-62); I-L-2, 80 (78-92); I-L-3, 103 (99-116); I-L-4, 157 (149-176); I-L-5, 179 (168-194); I-L-6, 111 (106-120). Dorsal lengths of the fourth leg: IV-L-1, 72 (70-88); IV-L-2, 123 (117-138); IV-L-3, 149 (138-162); IV-L-4, 268 (257-285); IV-L-5, 266 (259-281); IV-L-6, 208 (200-217).




Figures 4–8. *Sperchon (Palpisperchon) nikkoensis*, Imamura, 1976, Male **4** infracapitulum **5** palp **6** chelicera **7** IV-L-1–6; **8** claw.

Remarks. At present, only five species of the subgenus *Palpisperchon* Lundblad, 1941 are known: *Sperchon crassipalpis* Marshall, 1933, *S. distans* Scheffler, 1972, *S. mirabilis* Lundblad, 1941, *S. nikkoensis* Imamura, 1976 and *S. skopetsi* Tuzovskij, 1982.

Due to the shape of cuticle, E4 absent from CxIII, P-II with a very long ventrodistal projection, and P-IV with two greatly enlarged ventral peg-like setae, the specimens from China show a general conformity with *S. (P.) nikkoensis*, a species previously known only from the female sex from Japan (Imamura 1976). However, the absence of pregenital sclerite and the presence of a round platelet in front of the genital field (as reported in the original illustration, see Imamura 1976), are typical characters of the male sex (Di Sabatino et al. 2010). Therefore, the specimen described by Imamura should be a male instead of female. This is the first description of the opposite sex to that in original description (Imamura 1976), and first record of this species from China.

Distribution. China (present study); Japan (Imamura 1976).



Figures 9–10. *Sperchon (Palpisperchon) nikkoensis*, Imamura, 1976, Female **9** idiosoma, dorsal view **10** idiosoma, ventral view.

Subgenus *Sperchon* Kramer, 1877

Sperchon (Sperchon) orbipatella sp. n.

urn:lsid:zoobank.org:act:61043D65-F634-4D8C-9904-01EC2163CC42


http://species-id.net/wiki/Sperchon_orbipatella

Figures 11–21

Type series. Holotype: Male, Xinjiang Uygur Autonomous Region, Urumqi city, Baiyanggou scenic area, an unnamed stream (43°40'53"N, 88°04'59"E), 18 August 1996, coll. Dao-Chao Jin. Paratypes: 46 males and 59 females, the same data as the holotype.

Diagnosis. Cuticle covered with small papillae various in shape; excretory pore surrounded by a sclerotized ring; the second acetabulum near to the third but far away from the first one; P-II with a long ventro-distal projection and one thick seta at the base of the projection; third to fifth segments of leg I-IV with rather short plumose setae in longitudinal rows.


Description. Male (n=3): Idiosoma oval in outline, 636 (602–783) in length, 533 (510–563) in width. Cuticle yellow-brown, covered with small, various shaped papillae (Fig. 13). A1 smooth, short and thick, other glandularia thin and longer. Cuticle without an extended dorsal and ventral plate. Glandularia and O2 encircled by small rounded platelets (Fig. 11, Fig. 12). Coxae in four groups, surface of coxae reticulated. ACG 162 (153–176) in length, close to each other but not fused, posterior apodeme weakly developed. E2 laterally between ACG and PCG. PCG 239 (228–257)



Figures 11–13. *Sperchon (Sperchon) orbipatella* sp. n., Male **11** idiosoma, dorsal view **12** idiosoma, ventral view **13** decorations of cuticle.

in length. E4 close to anterior margin of CxIII. Distance between anterior end of ACG and posterior end of PCG 387 (364–412). Genital field between PCG with a small and rounded platelet in front. Genital valves not covering the genital acetabula, 170 (157–184) in length, 140 (132–165) in width. Pre- and postgenital sclerites not developed. Three pairs of small and rounded genital acetabula, the second acetabulum near to the third and far away from the first. V1 on very small sclerites and without accompanying glandularia. Excretory pore surrounded by a sclerotized ring and placed anterior to the line of V2 glands.


Infracapitulum with a relatively short and high rostrum, length 172 (165–184). Chelicera total length 223 (207–246), basal segment length 165 (155–183), claw length 58 (52–63), basal segment/claw length ratio 2.8 (2.8–3.0). Dorsal lengths of the palpal segments: P-I, 32 (25–34); P-II, 99 (80–107); P-III, 121 (113–134); P-IV, 163 (145–173); P-V, 37 (34–45). P-I short and without seta. P-II with a long ventro-distal projection bearing one thin seta and a thick peg-like seta at the base of the projection, the thick seta almost as long as the projection. Six setae on the dorsal side of the P-II, none of them plumose. The venter margin of P-III without setae, two setae on the dorsal side, one of which almost on middle and another one near to the distal end of the segment. P-IV venter with two subequal peg-like setae, well distanced from each other. Six thin setae on the dorsal side of P-IV, four of them on the terminal end of segment. Dorsal lengths of the first leg: I-L-1, 56 (47–69); I-L-2, 71 (58–86); I-L-3, 78 (62–88);



Figures 14–19. *Sperchon (Sperchon) orbipatella* sp. n., Male **14** infracapitulum **15** claw **16** chelicera **17** palp **18** IV-L-1–6 **19** dorsal seta of IV-L-5.

I-L-4, 144 (126-163); I-L-5, 145 (131-167); I-L-6, 143 (123-159). Dorsal lengths of the fourth leg: IV-L-1, 85 (76-93); IV-L-2, 107 (96-114); IV-L-3, 115 (108-127); IV-L-4, 238 (221-257); IV-L-5, 215 (197-226); IV-L-6, 205 (196-214). Third to fifth segments of leg I-IV with rather short plumose setae in longitudinal rows (Fig. 19). Claws with well protruded claw-blade and two clawlets, a long dorsal and a shorter ventral one (Fig. 15).

Female (n = 3): Similar to male except for the morphology of genital field (Fig. 21). Idiosoma 771 (725-947) in length, 645 (617-768) in width. ACG 199 (182-218) in length, PCG 272 (253-304) in length. Distance between anterior end of ACG and posterior end of PCG 448 (436-469). Genital field 209 (182-234) in length, 174



Figures 20–21. *Sperchon (Sperchon) orbipatella* sp. n., Female **20** idiosoma, dorsal view **21** idiosoma, ventral view.

(163–181) in width. Infracapitulum length 210 (197–242). Chelicera total length 283 (262–307), basal segment length 212 (195–230), claw length 71 (67–77), basal segment/claw length ratio 3.0 (2.9–3.0). Dorsal lengths of the palpal segments: P-I, 37 (32–41); P-II, 122 (113–137); P-III, 155 (140–172); P-IV, 197 (178–217); P-V, 48 (40–54). Dorsal lengths of leg I: I-L-1, 67 (54–73); I-L-2, 100 (87–113); I-L-3, 128 (112–147); I-L-4, 217 (203–246); I-L-5, 197 (184–213); I-L-6, 183 (169–212). Dorsal lengths of leg IV: IV-L-1, 95 (87–106); IV-L-2, 148 (132–167); IV-L-3, 151 (137–173); IV-L-4, 299 (267–315); IV-L-5, 280 (262–301); IV-L-6, 227 (209–246).

Etymology. The species is named after the shape of the three acetabula, “orbi-” Latin word, means rounded.

Remarks. All the species of the genus *Sperchon* have three pairs of acetabula, in most cases arranged almost equidistantly, usually the anterior two pairs of acetabula elongated while the posterior pair is somewhat rounded. However, in some species such as *Sperchon prosperoides* Tuzovskij, 1990 and *S. minutiporus* Sokolow, 1934, the acetabula are small and rounded, and the second pair shifted near to the third one but far away from the first one.

The new species resembles to *Sperchon prosperoides* Tuzovskij, 1990, from Russia, from which it can be distinguished by the morphology of the palp and genital field (Tuzovskij 2008). In *S. prosperoides*, P-III bearing two thick setae on the ventral side and only half of the genital field is located between PCG, whereas in the new species, P-III is without setae on the ventral side and the genital field is located entirely between PCG.

Distribution. China (Xinjiang Uygur Autonomous Region).

***Sperchon (Sperchon) souunkyo* Imamura, 1954**


http://species-id.net/wiki/Sperchon_souunkyo

Figures 22–31


Material examined. 25 males and 36 females, Xinjiang Uygur Autonomous Region, Altay City, Kanas Lake (48°48'59"N, 87°02'01"E), 14 August 1997, coll. Dao-Chao Jin; 2 males, Shanxi Province, Taibaishan Mountain National Forest Park, Honghua-ping area scene, an unnamed stream (34°07'16"N, 107°53'46"E), 02 September 1997, coll. Dao-Chao Jin; 3 females, Hebei Province, Wulingshan National Nature Reserve, an unnamed stream (40°37'23"N, 117°29'51"E), 03 April 2002, coll. Jian-Jun Guo; 2 males and 1 female, Guizhou Province, Xishui National Nature Reserve, an unnamed stream (28°28'19"N, 106°05'41"E), 02 Jun 2002, coll. Jian-Jun Guo; 10 males and 8 females, Guizhou Province, Kuankuoshui National Nature Reserve, an unnamed stream (28°14'25"N, 107°11'59"E), 22 May 2004, coll. Zhen-Zao Tian and Ai-Ping Cui; 11 males and 9 females, Guizhou Province, Leigongshan National Nature Reserve, an unnamed stream (26°21'24"N, 104°34'40"E), 21 May 2005, coll. Xu Zhang.

Description. Male (n = 4): Idiosoma oval in outline, 561 (523–635) in length, 396 (374–418) in width. Cuticle yellow-brown, covered with scale-shaped papillae (Fig. 24). A1 short and plumose, setae of other glandularia smooth and relatively long. Cuticle without dorsalia and ventralia except the platelets surrounding glandularia and O2 (Fig. 22, Fig. 23). Coxae in four groups, surface of coxae reticulated. ACG 157 (148–166) in length, apodeme developed. E2 laterally between ACG and PCG. PCG 167 (160–174) in length. E4 on the median and near to inner margin of CxIII. Distance between anterior end of ACG and posterior end of PCG 308 (283–317). Genital field between PCG with a small and rounded platelet in front. Genital valves not covering the genital acetabula, 113 (108–121) in length, 109 (101–114) in width. Pre- and postgenital sclerites visible though not developed. Three pairs of genital acetabula, the first pair of genital acetabula more or less rectangular, the second pair somewhat triangular, and the third pair rounded and larger than the anterior two. V1 on median size sclerites and without accompanying glandularia. Excretory pore without sclerotized ring, and anterior to the line of V2 glands.


Infracapitulum with a short rostrum, length 193 (185–203). Chelicera total length 204 (194–219), basal segment length 151 (146–160), claw length 53 (48–59), basal segment/claw length ratio 2.8 (2.7–3.0). Dorsal lengths of the palpal segments: P-I, 21 (19–23); P-II, 94 (85–102); P-III, 102 (94–108); P-IV, 97 (92–111); P-V, 24 (22–27). P-I short and without seta. P-II with a long ventro-distal projection, one thin and long seta near to the base of the projection. Four setae on the dorsal and lateral side of the P-II, one of them plumose. The ventral side of P-III nearly straight and without seta, six short smooth setae on the lateral and dorsal side. P-IV with two small peg-like ventral setae and three thin setae, the proximal peg-like seta is larger than the distal one. The dorsal of P-IV with three thin setae, two of them located distally. Dorsal lengths of the first leg: I-L-1, 44 (39–50); I-L-2, 63 (55–71); I-L-3, 68 (59–76); I-L-4, 119 (112–138); I-L-5, 109 (102–123); I-L-6, 105 (97–114). Dorsal lengths of the fourth leg: IV-L-1, 61 (57–



Figures 22–24. *Sperchon (Sperchon) sounkyo* Imamura, 1954, Male **22** idiosoma, dorsal view **23** idiosoma, ventral view **24** decorations of cuticle.



Figures 25–29. *Sperchon (Sperchon) sounkyo* Imamura, 1954, Male **25** infracapitulum **26** chelicera **27** claw **28** palp **29** IV-L-1-6.



Figures 30–31. *Sperchon (Sperchon) sounkyo* Imamura, 1954, Female **30** idiosoma, dorsal view **31** idiosoma, ventral view.

65); IV-L-2, 85 (76-95); IV-L-3, 103 (92-116); IV-L-4, 195 (183-210); IV-L-5, 204 (191-218); IV-L-6, 162 (149-178). Third to fifth segments of leg I-IV with rather short smooth setae in longitudinal rows. Ambulacrum with two claws and each claw with protruded claw-blade and two clawets, a long dorsal and a shorter ventral one (Fig. 28).

Female (n = 3): Similar to male except for the morphology of genital field and the size of idiosoma (Fig. 30, Fig. 31). Idiosoma 910 (768-944) in length, 570 (493-648) in width. ACG 182 (170-190) in length, PCG 215 (204-221) in length. Distance between anterior end of ACG and posterior end of PCG 397 (381-405). Genital field 174 (164-181) in length, 154 (144-162) in width. Pre- and postgenital sclerites more developed than in the male. Infracapitulum length 229 (213-238). Chelicera total length 270 (246-279), basal segment length 200 (183-205), claw length 70 (63-74), basal segment/claw length ratio 2.9 (2.7-2.9). Dorsal lengths of the palpal segments: P-I, 23 (20-25); P-II, 123 (118-130); P-III, 149 (142-156); P-IV, 113 (107-122); P-V, 29 (26-33). Dorsal lengths of leg I: I-L-1, 43 (37-49); I-L-2, 65 (60-74); I-L-3, 85 (79-97); I-L-4, 147 (136-154); I-L-5, 134 (125-147); I-L-6, 112 (106-120). Dorsal lengths of leg IV: IV-L-1, 75 (70-82); IV-L-2, 111 (103-119); IV-L-3, 118 (112-127); IV-L-4, 242 (221-263); IV-L-5, 223 (208-237); IV-L-6, 187 (178-196).

Remarks. *Sperchon sounkyo* Imamura, 1954, was firstly described from Japan based on a single female (Imamura 1954), and never recorded since the first description.

Although the original descriptions and illustrations of *S. sounkyo* were lacking of some specific features, such as shape of A1 (Fig. 22, Fig. 30) and leg claws (Fig. 28), most characters of our specimens (e.g., shape of cuticle and palp, E4 on CxIII, and excretory pore without sclerotized ring) are in accordance with this species. So we attribute our specimens to *S. sounkyo* Imamura, 1954. Some differences found in the body size and the number of the setae on the palpal between the Chinese specimens and Japanese specimens should be regarded as the variety between the different populations.

It is the first report from China and the first description of the male of *S. sounkyo*. This species seems to be a widely distributed species in China.

Distribution. China (present study); Japan (Imamura 1954).

***Sperchon (Sperchon) urumqiensis* sp. n.**

urn:lsid:zoobank.org:act:53DAC024-9964-47BA-BA72-B3563069B059

http://species-id.net/wiki/Sperchon_urumqiensis


Figures 32–42

Type series. Holotype: Male, Xinjiang Uygur Autonomous Region, Urumqi city, Baiyanggou scenic area, an unnamed stream (43°37'42"N, 87°57'18"E), 18 Aug 1996, coll. Dao-Chao Jin. Paratypes: 6 males and 8 females, the same data as the holotype.

Diagnosis. Cuticle covered with scale-shaped papillae; excretory pore surrounded by a sclerotized ring; P-II with a long ventro-distal projection and one heavy seta close to the ventrodistal margin of P-II; third to fifth segments of leg I-IV with rather long plumose setae in longitudinal rows.


Description. Male (n = 3): Idiosoma oval in shape, 812 (756-907) in length, 682 (663-712) in width. Cuticle yellow in colour, covered with scale-shaped papillae (Fig. 34). A1 smooth, short and thick, other glandularia thin and long. All glandularia and O2 encircled by a platelet. Some chitinous platelets and glandular platelets on dorsum fused as shown in Fig. 32. Coxae in four groups, surface of coxae reticulated. ACG 163 (158-171) in length, posterior apodeme weakly developed. E2 laterally between ACG and PCG. PCG 242 (236-254) in length. E4 close to anterior margin of CxIII. Distance between anterior end of ACG and posterior end of PCG 443 (432-464). Genital field between PCG with a small and rounded platelet in front. Genital valves not covering the genital acetabula, 188 (182-193) in length, 117 (113-124) in width. Pre- and postgenital sclerites not developed. Three pairs of acetabula, the two anterior pairs of acetabula long elliptic and the posterior pair more or less rounded. An oblong plate posterior to genital field, almost in the middle line of V1 and V3. V1 on very small sclerites and without accompanying glandularia. Excretory pore surrounded by a sclerotized ring and close to the line at V2 glands.

Infracapitulum with a short and heavy rostrum, length 189 (185-197). Chelicera total length 229 (225-237), basal segment length 174 (172-179), claw length 55 (53-



Figures 32–34. *Sperchon (Sperchon) urumqiensis* sp. n., Male **32** idiosoma, dorsal view **33** idiosoma, ventral view **34** decorations of cuticle.


58), basal segment/claw length ratio 3.2 (3.1–3.2). Dorsal lengths of the palpal segments: P-I, 22 (21–24); P-II, 99 (97–106); P-III, 118 (113–127); P-IV, 171 (165–182); P-V, 37 (34–42). P-I short and with one dorsal seta. P-II with a long ventro-distal projection bearing two thin setae, of which one longer than another. One heavy seta close to the ventrodistal margin of P-II. Seven setae on the dorsal and lateral side of the P-II, none of them plumose. The ventral side of P-III somewhat swelled and without seta, two long smooth setae on the lateral and dorsal side. P-IV slender, much longer than P-III, and with two ventral peg-like setae, of which the proximal one almost at ventral middle of the segment, and the distal one approximately equidistant from proximal one and distal end of the segment. Four thin setae on P-IV, one of them on the median dorsal, two on the dorsal distal and another one on the terminal end of the segment. Dorsal lengths of the first leg: I-L-1, 39 (36–43); I-L-2, 83 (80–90); I-L-3, 98 (90–106); I-L-4, 191 (185–200); I-L-5, 182 (177–193); I-L-6, 179 (174–187). Dorsal lengths of the fourth leg: IV-L-1, 87 (85–94); IV-L-2, 117 (111–128); IV-L-3, 154 (148–168); IV-L-4, 300 (290–317); IV-L-5, 271 (264–285); IV-L-6, 257 (249–266). Third to fifth segments of leg I–IV with rather long plumose setae in longitudinal rows (Fig. 39).



Figures 35–40. *Sperchon (Sperchon) urumqiensis* sp. n., Male **35** infracapitulum; **36** chelicera **37** claw **38** IV-L-1-6 **39** dorsal seta of IV-L-4 **40** palp.

Ambulacrum with two claws and each claw with well protruded claw-blade and two clawlets, a long dorsal and a shorter ventral one (Fig. 37).

Female (n = 3): Color, idiosoma shape, and the decorations of cuticle as in the male. Characters of the genital field, the fusion and the size of dorsalia and ventralia different from the male (Fig. 41, Fig. 42). Idiosoma 979 (953-1023) in length, 829



Figures 41–42. *Sperchon (Sperchon) urumqiensis* sp. n., Female **41** idiosoma, dorsal view **42** idiosoma, ventral view.

(805-876) in width. ACG 202 (197-212) in length, PCG 301 (295-314) in length. Distance between anterior end of ACG and posterior end of PCG 559 (547-578). Genital field 227 (223-234) in length, 245 (241-253) in width. Pre- and postgenital sclerites well developed than that in the male. Infracapitulum length 255 (250-261). Chelicera total length 317 (311-331), basal segment length 244 (241-253), claw length 73 (70-78), basal segment/claw length ratio 3.3 (3.2-3.4). Dorsal lengths of the palpal segments: P-I, 47 (45-50); P-II, 146 (140-155); P-III, 181 (174-193); P-IV, 274 (263-289); P-V, 45 (42-49). Dorsal lengths of leg I: I-L-1, 70 (64-78); I-L-2, 135 (127-148); I-L-3, 128 (122-137); I-L-4, 256 (243-171); I-L-5, 272 (260-284); I-L-6, 234 (222-247). Dorsal lengths of leg IV: IV-L-1, 115 (119-124); IV-L-2, 156 (148-169); IV-L-3, 230 (219-244); IV-L-4, 379 (366-397); IV-L-5, 363 (351-380); IV-L-6, 322 (311-340).

Etymology. The species is named after the city where it was collected.

Remarks. Due to the shape of cuticle, E4 on the CxIII and P-II with a long ventro-distal projection, the new species is similar to *Sperchon sounkyo* Imamura, 1954. From the later species, *S. urumqiensis* sp. n. can be easily distinguished by the following features: the dorsum and posterior venter with sclerotized muscle attachment plates (Fig. 32-33, Fig. 41-42) (unsclerotized in *S. sounkyo*, see Fig. 22-23, Fig. 30-31), the presence of a long and heavy seta close to the ventrodistal margin of P-II (Fig. 40) (vs. without heavy seta in *S. sounkyo*, see Fig. 27), the two peg-like setae of P-IV are sub-

equal in size (Fig. 40) (proximoventral seta larger than distoventral in *S. sounkyo*, see Fig. 27), and the excretory pore is sclerotized (Fig. 33, Fig. 42) (smooth in *S. sounkyo*, see Fig. 23, Fig. 31).

The new species also resemble *Sperchon ussuriensis* Tuzovskij, 2002 from Russia (Tuzovskij 2002). However, in *S. ussuriensis*, the anterior groups of coxae are fused and the excretory pore is unsclerotized. While in the new species, the anterior groups of coxae are not fused and the excretory pore is surrounded by a sclerotized ring.

Distribution. China (Xinjiang Uygur Autonomous Region).

Genus *Sperchonopsis* Piersig, 1896

Subgenus *Sperchonopsella* Conroy, 1991

Sperchonopsis (*Sperchonopsella*) *whiteshellensis* Conroy, 1991


http://species-id.net/wiki/Sperchonopsis_whiteshellensis

Figures 43–51

Material examined. 3 males and 6 females, Xinjiang Uygur Autonomous Region, Altay City, Kanas Lake (48°48'52"N, 86°57'47"E), 14 August 1997, coll. Dao-Chao Jin; 24 males and 33 females, Yunnan Province, Tengchong country, Gaoligongshan National Nature Reserve, an unnamed stream (25°33'12"N, 98°35'18"E), 15 July 2002, coll. Jian-Jun Guo.


Description. Male (n = 4): Idiosoma oval in outline, 578 (562–675) in length, 418 (410–457) in width. Cuticle yellow-brown in color, and covered with papillae. The eye capsules in some young specimens well developed and somewhat projected over the idiosoma margin (Fig. 43). Glandularia greatly enlarged and projected with well-developed papillae, and encircled by platelet. Chitinous platelets and glandularia on dorsum and ventrum as showed in Fig. 43 and Fig. 44. Coxae in four groups, surface of coxae reticulated. ACG 152 (151–157) in length, posterior apodeme indistinct. Anterior tip of CxI with a tuft of hair-like setae. E2 laterally between ACG and PCG. PCG 167 (164–175) in length. Glandularia absent from CxIII. Approximately two-thirds of genital field lying between PCG. Genital valves not covering the genital acetabula, 176 (174–188) in length, 145 (143–152) in width. One round platelet in front of genital field. Pre- and postgenital sclerites small. Three pairs of small and rounded genital acetabula, the second pair much close to the third one and far away from the first one. V1 on sclerites in medium size and without accompanying glandularia. Excretory pore lying between V1 and V2 and on a protuberance surrounded by a well-developed sclerotized ring.

Infracapitulum with a relatively long rostrum, length 196 (194–207). Chelicera total length 207 (204–222), claw length 151 (149–162), basal segment length 56 (55–60), basal segment/claw length ratio 2.7 (2.7). Dorsal lengths of palpal segments: P-I, 24 (24–27); P-II, 104 (101–112); P-III, 94 (92–100); P-IV, 96 (93–105); P-V, 33 (32–36). P-I short and without seta. P-II with a long ventro-distal projection bearing one long and one short setae. About four setae on the lateral and dorsal side of P-II and



Figures 43–49. *Sperchonopsis (Sperchonopsella) whiteshellensis* Conroy, 1991, Male **43** idiosoma, dorsal view **44** idiosoma, ventral view **45** infracapitulum **46** chelicera **47** palp **48** IV-L-1–6 **49** claw.

none of them plumose. P-III with five smooth stae, of them two on the venter and other three on the latero-dorsal side. P-IV without peg-like setae, but with a ventral projection bearing two setae. Two setae on the dorsal and the ventral distal of P-IV



Figures 50–51. *Sperchonopsis (Sperchonopsella) whiteshellensis* Conroy, 1991, Female **50** idiosoma, dorsal view **51** idiosoma, ventral view.

respectively. Dorsal lengths of the first leg: I-L-1, 44 (42-49); I-L-2, 43 (42-48); I-L-3, 57 (54-64); I-L-4, 97 (93-108); I-L-5, 87 (84-97); I-L-6, 51 (49-57). Dorsal lengths of the fourth leg: IV-L-1, 70 (67-77); IV-L-2, 68 (62-74); IV-L-3, 88 (85-97); IV-L-4, 168 (160-181); IV-L-5, 169 (158-184); IV-L-6, 152 (148-164). The distal edge of I-IV-L-2–I-IV-L-5 with several thick and plumose setae (Fig. 48). Ambulacrum with two claws and each claw with well protruded claw-blade and two clawets, a long dorsal and a shorter ventral one (Fig. 49).

Female (n = 3): Similar to male except the characteristics of genital field and the size of idiosoma. Idiosoma length 886 (853-936), width 626 (607-645). ACG 200 (194-205) in length, PCG 212 (204-221) in length. Distance between anterior end of ACG and posterior end of PCG 438 (424-452). Genital field 213 (207-217) in length, 170 (167-175) in width. Pregenital sclerite well developed. Infracapitulum length 266 (257-276). Chelicera total length 279 (272-289), basal segment length 193 (189-199), claw length 86 (83-90), basal segment/claw length ratio 2.2 (2.2-2.3). Dorsal lengths of the palpal segments: P-I, 36 (34-40); P-II, 155 (148-162); P-III, 132 (126-140); P-IV, 136 (125-143); P-V, 45 (42-48). Dorsal lengths of the first leg: I-L-1, 73 (66-81); I-L-2, 75 (67-83); I-L-3, 84 (77-93); I-L-4, 122 (114-131); I-L-5, 127 (119-138); I-L-6, 108 (101-117). Dorsal lengths of the fourth leg: IV-L-1, 108 (102-117); IV-L-2, 104 (98-112); IV-L-3, 126 (118-135); IV-L-4, 222 (213-233); IV-L-5, 195 (186-207); IV-L-6, 181 (173-190).

Remarks. The subgenus *Sperchonopsella* is a small group with only two known species, *Sperchonopsis whiteshellensis* Conroy, 1991 and *S. nipponicus* Uchida, 1934 (Conroy 1991).

Due to the papillate cuticle with the greatly enlarged glandularia, the second and the third pair of acetabula approached to each other and the ventral projections on P-I and P-IV, the specimens from China are similar to *S. whiteshellensis* from North America (Conroy 1991) and *S. nipponicus* from Japan (Uchida 1934). The arrangement of dorsalia and ventralia are in a good agreement with description of *S. whiteshellensis*, and differs from *S. nipponicus* (see illustration in Uchida 1934). Hence we assigned our specimens to *S. whiteshellensis*. This is the first record of the subgenus *Sperchonopsella* from China.

Distribution. China (present study); North America (Conroy 1991)

Acknowledgements

We are very grateful to Dr. Peter Tuzovskij (Institute for Biologie of Inland Waters, Borok, Jaroslavl, Russia) for providing literature and Dr. Yi Tian-Ci (Guizhou University, China) for his helps in many ways. Further, we are thankful to Dr. Wang Xing (China Agricultural University) for his help in finding the full text of literature. This study was supported by The National Science Foundation of China (No. 30840021) and the Natural Science Foundation (KJ2011B143) from Department of Education, Anhui Province, China.

References

- Conroy JC (1991) New species in the genus *Sperchonopsis* in North America with a description of a new subgenus, *Sperchonopsella*. II. *Acarologia* 32: 151–161.
- Cook DR (1974) Water mite genera and subgenera. *Memoirs of the American Entomological Institute* 21: 1–860.
- Di Sabatino A, Gerecke R, Gledhill T, Smit H (2010) Chelicerata: Acari II. In: Gerecke R (Ed) 2006. *Chelicerata: Araraneae, Acari I. Süßwasserfauna von Mitteleuropa*, Vol. 7, 2–2, Elsevier Spektrum Akademischer Verlag, München, 134pp.
- Di Sabatino A, Smit H, Gerecke R, Goldschmidt T, Matsumoto N, Cicolani B (2008) Global diversity of water mites (Acari, Hydrachnidia; Arachnida) in freshwater. *Hydrobiologia* 595: 303–315. doi: 10.1007/s10750-007-9025-1
- Imamura T (1954) Studies on water-mites from Hokkaido. *Journal of Hokkaido Gakugei University, Section B, Supplement* 1: 1–148.
- Imamura T (1976) Two New Species of Water-Mites from Nikko National Park, Japan. *Annotationes zoologicae Japonenses* 49 (4): 279–284.
- Jin DC (1997) *Hydrachnellae: morphology, systematics; a primary study of Chinese fauna*. Guizhou science and technology publishing house, Guiyang, 356pp. [in Chinese]
- Jin DC, Yi TC, Guo JJ (2010) A review of progress in taxonomy of water mites from China (Acari: Hydrachnidia). *Zoosymposia* 4: 106–119.

- Tuzovskij PV (2002) A new species of the water mite genus *Sperchon* Kramer from Primorsk territory of Russia (Acariformes: Sperchonidae). *Zoosystematica Rossica* 11 (1): 105–108.
- Tuzovskij PV (2008) The morphology of three species of adult water mites of the genus *Sperchon* (*S. kuluensis*, *S. prosperoides*, and *S. orientalis*, Acariformes, Sperchontidae). *Entomological Review* 88 (2): 139–149. doi: 10.1134/S0013873808020024
- Uchida T (1934) Some rheophilous water-mites from Japan. *Journal of the Faculty of Science, Hokkaido Imperial University, Series VI (Zoology)* 3(2): 67–116.
- Viets KO (1987) Die Milben des Süßwassers (Hydrachnellae und Halacaridae [part.], Acari). 2. Katalog. Sonderbände des Naturwiss. Vereins Hamburg 8: 1–1012.
- Zhang X, Jin DC (2010) Three new species and one new record of the subgenus *Hispidosperchon* Thor, 1901 within the genus *Sperchon* Kramer, 1877 from China. *Zootaxa* 2684: 14–24.
- Zhang X, Jin DC, Guo JJ (2011) Two new water mites species of the genus *Sperchon* Kramer, 1877 (Acari, Hydrachnidia, Sperchontidae). *Acta Zootaxonomica Sinica* 36 (2):221–226.
- Zhang X, Jin DC, Guo JJ, Yi TC (2010) Descriptions of two new species of *Mixosperchon* Viets, a newly recorded subgenus of water mites, from China (Acari, Hydrachnellae, Sperchontidae). *Acta Zootaxonomica Sinica* 35 (3): 435–439.
- Zhang X, Jin DC, Guo J J, Zhu Q (2007) Three species of the Family Sperchontidae (Acari, Hydrachnellae, Sperchontidae) as new records from China. *Sciencepaper Online*. (2007–12–28):1–5. [in Chinese] <http://www.paper.edu.cn/index.php/default/releasepaper/download/Paper/200712-845/2>

