

# **Coastal Staphylinidae (Coleoptera): A worldwide checklist, biogeography and natural history**

J. H. Frank  
&  
Kee-Jeong Ahn



Sofia–Moscow

2011

ZooKeys 107 (SPECIAL ISSUE)

COASTAL STAPHYLINIDAE (COLEOPTERA): A WORLDWIDE CHECKLIST, BIOGEOGRAPHY AND  
NATURAL HISTORY

*J. H. Frank & Kee-Jeong Ahn*

First published 2011

ISBN 978-954-642-595-9 (paperback)

Pensoft Publishers

Geo Milev Str. 13a, Sofia 1111, Bulgaria

Fax: +359-2-870-42-82

[info@pensoft.net](mailto:info@pensoft.net)

[www.pensoft.net](http://www.pensoft.net)

Printed in Bulgaria, June 2011

# Contents

<b>2</b>	<b>Introduction</b>
<b>17</b>	<b>A Checklist of coastal Staphylinidae and their distribution</b>
<b>36</b>	<b>Habits, habitats, and classificatory notes</b>
36	Microsilphinae
36	Omaliinae
37	Aphaenostemmini
37	Omaliini
38	Pselaphinae
38	Batrisitae
38	Batrisini
38	Goniaceritae
38	Brachyglutini
39	Aleocharinae
39	Aleocharini
40	Athetini
43	Diglottini
43	Falagriini
44	Homalotini
45	Liparocephalini
47	Myllaenini
49	Oxypodini
50	Phytosini
51	Incertae sedis
51	Oxytelinae
51	Oxytelini
51	Thinobiini

53	Scydmaeninae
53	Cephenniini
54	Paederinae
54	Paederini
54	Staphylininae
54	Staphylinini
<b>58</b>	<b>Discussion</b>
<b>68</b>	<b>Summary</b>
<b>68</b>	<b>Acknowledgments</b>
<b>69</b>	<b>References</b>



# Coastal Staphylinidae (Coleoptera): A worldwide checklist, biogeography and natural history

J. H. Frank<sup>1</sup>, Kee-Jeong Ahn<sup>2</sup>

**1** Entomology and Nematology Department, University of Florida, Gainesville, FL 32611-0630, USA **2** Department of Biology, Chungnam National University, Daejeon 305-764, Republic of Korea

Corresponding author: Kee-Jeong Ahn (kjahn@cnu.ac.kr)

---

Academic editor: Jan Klimaszewski | Received 8 September 2010 | Accepted 14 February 2011 | Published 16 June 2011

---

**Citation:** Frank JH, Ahn K-J (2011) Coastal Staphylinidae (Coleoptera): A worldwide checklist, biogeography and natural history. ZooKeys 107: 1–98. doi: 10.3897/zookeys.107.1651

---

## Abstract

We provide a list of the 392 described species of Staphylinidae confined to coastal habitats worldwide. The list is in taxonomic sequence by subfamily, tribe, and genus and includes 91 genera. We provide the page reference of the original description of every species and genus listed and of many synonyms. We note the existence of recent reviews, phylogenies and keys of each of the tribes and genera included. Coastal Staphylinidae contain eight subfamilies: Microsilphinae, Omaliinae, Pselaphinae, Aleocharinae, Oxytelinae, Scydmaeninae, Paederinae, and Staphylininae.

By ‘coastal habitats’ we mean habitats existing on the sea coast and subject to inundation or at least splashing by the very highest tides. This includes rocky, boulder, coral, sandy, and muddy seashores, and at least portions of salt-marshes, estuaries, and mangrove swamps. We exclude the sand dune habitat and higher parts of sea-cliffs.

The list notes distribution of all the species, first according to the ocean or sea on whose shores it has been recorded, and second by country (and for the larger countries by province or state). Although this distribution is undoubtedly incomplete, it provides a basis for future development of a dedicated database.

The ‘Habitats, Habits, and Classificatory Notes’ section is designed to provide ecologists with further taxonomic and ecological information. It includes references to descriptions of the immature stages, behavior of adults and immatures, their food, natural enemies, and habitat. We would have preferred to separate these entities, but current knowledge of ecology is developed in few instances beyond natural history.

The Pacific Ocean basin was the origin and contributed to the dispersal of the majority of specialist coastal Staphylinidae at the level of genus. However, at the level of species, species belonging to non-coastal-specialist genera are about as likely to occur on the shores of other oceans as on the shores of the Pacific. This difference is a reflection of the antiquity of coastal genera and species.

A complete bibliography, and habitat and habitus photographs of some representative coastal Staphylinidae species are provided.

### **Keywords**

seashore Staphylinidae, marine Staphylinidae, littoral Staphylinidae, intertidal Staphylinidae, habitat, behavior

## **Introduction**

We struggled to find an appropriate title for this work, but eventually rejected the expressions intertidal, marine, littoral, and seashore, all of which have been used by other authors. By “coastal” we mean species that dwell on sea coasts and are restricted to such habitats. However, we restrict the definition to habitats to those that are normally or occasionally inundated by tides, excluding cliff and dune habitats, as well as inland salt-laden habitats. Thus, those species included dwell in the intertidal zone and at the drift line, and in mangrove swamps, salt marshes, and estuaries where they may be inundated by the tides. We do not employ the terms halophile and halobiont because they refer to organisms that dwell in salt-laden habitats, which are not restricted to coastal areas; indeed saline lakes and ponds occur hundreds of kilometers from coasts, and we do not wish to consider these.

A book chapter on intertidal Staphylinidae (Moore and Legner 1976) was undoubtedly Ian Moore’s major contribution to this, his favorite, subject. Moore had already published numerous papers on intertidal Staphylinidae of the Pacific coast of North America. In this pioneering treatment he endeavored to summarize the world literature on intertidal staphylinids. To do so it was necessary to separate the literature on intertidal species from that on non-marine species with which it was intermixed. The task was daunting because of the number of species in the family (over 54,000 are now recognized), a huge polyglot literature, and the lack of habitat information in many of the early taxonomic publications. He included keys to identification of adults to the level of genus.

In the subsequent 35 years, some genera that Moore dealt with have been revised (see particularly studies by Ahn, Ashe, Assing, Gusarov, Haghebaert, Herman, Jeon, Klimaszewski, Maruyama, and Zerche), additional species have been described, synonymies have been reported, there has been much change in the higher classification of Staphylinidae, and there have been some studies of the behavior of intertidal Staphylinidae. These changes make an updated contribution worth undertaking. This contribution is not simply an updating of Moore and Legner (1976), but it additionally lists all the staphylinid species (not just genera) that are believed to be restricted to

coastal habitats. This list is augmented by page-references to their original descriptions in the literature, to references to generic revisions, and to publications on behavior of the species in question. We believe this will enable interested readers to access the original literature more readily. In this contribution, however, we do not include keys to identification of adults. For these the reader is urged to consult the cited literature. This contribution is intended for the reader who is willing and able to tackle the taxonomic literature, even if the ultimate objective is ecological or ethological.

This contribution lists some 392 species, in 91 genera, of Staphylinidae that are believed to be confined to coastal habitats. Some genera are confined to coastal habitats. Others include species that are confined to coastal habitats (and primarily only those with such restricted habitat are included in this treatment). One large genus, *Bledius* Leach, is exceptional in that its members live on banks of either freshwater or saline water bodies. Among the latter group it is in many instances unclear whether they are restricted to marine saline habitats.

Moore and Legner (1976) included not only genera that restricted to intertidal habitats, but also genera whose members had frequently or occasionally been found there. Our contribution is more selective in that it tries to admit only those species for which there is evidence of restriction to such habitats. This attempt to concentrate on true coastal species is emphasized by Smetana (2009) in his critical review of a paper published by Majka et al. (2008). Smetana's (2009) viewpoint is to exclude those species that are occasional or even frequent visitors to the coastal habitat, and to concentrate on those that are confined to the habitat. There is some difficulty in treating members of the large genus *Bledius* because some species occupy not only coastal habitats, but also inland saline habitats; we attempt to exclude species that are not restricted to coastal habitats.

The current epitome of a study of regional coastal staphylinids is that by Hammond (2000). It discusses systematics and distribution of the British species, but it includes species that are not restricted to this habitat. The converse is perhaps Hasegawa and Kanie (1992) which provided a list of Staphylinidae collected at one seashore locality in a 2-year time period, making no distinction between those species restricted to seashores and those incidentally found there, and makes no mention of their wider distribution. Lengerken (1929) published an extensive compilation of the distribution of coastal species of the North, Baltic, Irish, Mediterranean, Black, and Caspian seas. Audisio and Taglianti (2010) presented a compilation of Coleoptera found on Italian coasts, but did not indicate which ones among the included species are restricted to such habitats.

The adjective halophilous seems first to have been used in English in the late 19<sup>th</sup> century to mean plants that are salt-loving, or growing in salt marshes (OED 1971). Since then, a set of terms evolved to describe adaptations of organisms to saline environments:

1. Halobionts (obligate inhabitants of saline habitats),
2. Halophiles (facultative inhabitants of saline habitats),

3. Haloxenes (halotolerant species),
4. Incidentals (species not specifically associated with saline habitats but regularly found there).

Such terms (in German) were used by Lengerken (1929) and other authors in English and French. We do not use these terms because they do not describe exactly what we want to include (and exclude), which is species that are restricted to sea coasts. We suppose that all of the species we list are halobionts, but we exclude halobionts living on the shores of inland saline lakes.

### The checklist


This checklist is the first to attempt to enumerate all coastal staphylinids, and their distribution. Arrangement is taxonomic including subfamily, tribe, and genus; subtribes are included where defined. References to original generic and specific description are given. Generic and species synonyms are listed, each with original bibliographic reference. Listing of names of species within genera, genera within tribes, and tribes within subfamilies are mostly alphabetical, but names of subfamilies are arranged in taxonomic sequence. The arrangement followed for the higher categories is that of Lawrence and Newton (2000), Newton and Thayer (2007), and Grebennikov and Newton (2009) (Fig. 1).

To the far right of the taxonomic entries for genera (and in one instance for a tribe) reference to recent taxonomic revisions and phylogenies and keys to identification of adults [e.g., rev. Ahn 1996a; phy. Ahn and Ashe 1996b; key Moore and Legner 1976] is given.

Because this work deals with coastal species, primary geographical entries are given according to the oceans and seas on which species are found. Secondary entries are the names of the countries they inhabit, and tertiary entries (if any) are the (mainly political) subunits of larger countries, or islands belonging to the former. Compression of this information into a checklist required the use of abbreviations, which are as follows:

**A.** Codes used for oceans have 3 letters: ACO (Arctic Ocean), INO (Indian Ocean), NAO (North Atlantic Ocean), NPO (North Pacific Ocean), SAO (South Atlantic Ocean), and SPO (South Pacific Ocean). Names of seas and gulfs are spelled out: Andaman Sea, Arabian Sea, Arafura Sea, Bali Sea, Baltic Sea, Bering Sea, Bismarck Sea, Black Sea, Caribbean Sea, Celebes Sea, East China Sea, East Sea [sometimes called Sea of Japan, but that name is disputed (Wikipedia 2010)], Gulf of California (sometimes called Sea of Cortez), Gulf of Mexico, Irish Sea, Java Sea, Mediterranean Sea (here including Adriatic, Aegean, Ionian, Ligurian and Tyrrhenian Seas), North Sea, Sea of Okhotsk, Philippine Sea, Red Sea, South China Sea, Sulu Sea, Tasman Sea, and Timor Sea.

Names of seas are not used throughout. We have used the name of the ocean in the broad sense (of which the sea is part) in instances where the name of the sea is not ap-



**Figure 1.** Phylogeny of the Staphylinidae. Bold indicates eight subfamilies containing coastal species. Modified from Newton and Thayer (2007) and Grebennikov and Newton (2009).

parent from the literature. For example, some species known from New Zealand may be known from the west coast (the Tasman Sea), but if that was not apparent from the literature, we ascribed them to SPO (the South Pacific Ocean).

**B.** Country codes have 2 letters and are the International Standards Organization (ISO) abbreviations. They are given in parentheses. Those used are: AG=Antigua and

Barbuda, AL=Albania, AR=Argentina, AU=Australia, BB=Barbados, BE=Belgium, BG=Bulgaria, BM=Bermuda, BR=Brazil, BS=Bahamas, CA=Canada, CL=Chile, CN=China, CO=Columbia, CU=Cuba, CY=Cyprus, DE=Germany, DJ=Djibouti, DK=Denmark, DM=Dominica, DO=Dominican Republic, DZ=Algeria, EC=Ecuador, EE=Estonia, EG=Egypt, ER=Eritrea, ES=Spain, ET=Ethiopia, FI=Finland, FJ=Fiji, FP=French Polynesia, FR=France, GB=Great Britain, GD=Grenada, GE=Georgia, GH=Ghana, GL=Greenland, GP=Guadeloupe, GR=Greece, HR=Croatia, HT=Haiti, ID=Indonesia, IE=Ireland, IL=Israel, IN=India, IQ=Iraq, IS=Iceland, IT=Italy, JM=Jamaica, JP=Japan, KE=Kenya, KN=St. Kitts Nevis, KP=North Korea, KR=South Korea, KY=Cayman Island, LB=Lebanon, LC=St. Lucia, LK=Sri Lanka, LY=Libya, MA=Morocco, MG=Madagascar, MM=Myanmar, MR=Mauritania, MS=Montserrat, MT=Malta, MU=Mauritius, MX=Mexico, MY=Malaysia, NA=Namibia, NC=New Caledonia, NG=Nigeria, NL=Netherlands, NO=Norway, NZ=New Zealand, PE=Peru, PG=Papua New Guinea, PH=Philippines, PL=Poland, PR=Puerto Rico, PT=Portugal, RE=Reunion, RO=Romania, RU=Russian Federation, SA=Saudi Arabia, SC=Seychelles, SD=Sudan, SG=Singapore, SN=Senegal, SE=Sweden, SO=Somalia, TH=Thailand, TN=Tunisia, TR=Turkey, TT=Trinidad and Tobago, TW=Taiwan, TZ=Tanzania, UA=Ukraine, UK=United Kingdom, US=USA, UY=Uruguay, VE=Venezuela, VI=US Virgin Islands, VN=Vietnam, WS=Samoa (formerly Western Samoa, not American Samoa), YE=Yemen, YU=former Yugoslavia, ZA=South Africa.

**C.** Where places within countries are mentioned, they are given after a colon (:) following the abbreviation of the country name, and either are spelled out or are abbreviated. For the USA and Canada, the abbreviations are the 2-letter postal codes (BC=British Columbia, NB=New Brunswick, NL=Newfoundland and Labrador, NS=Nova Scotia, NT=Northwest Territories, PE=Prince Edward Island, QC=Quebec, YT=Yukon Territory); for Mexico they have 2 letters (BN=Baja California, BS=Baja California Sur, CA=Campeche, CH=Chiapas, GU=Guerrero, JA=Jalisco, MI=Michoacán, NA=Nayarit, OA=Oaxaca, QR=Quintana Roo, SI=Sinaloa, SO=Sonora, TB=Tabasco, TM=Tamaulipas, and VC=Veracruz); for Japan, designations are for major islands and island groups: (HK=Hokkaido, HN=Honshu, KY=Kyushu, RY=Ryukyu, SH=Shikoku). For Great Britain (GB) they are England, N. Ireland, Scotland, and Wales.

Compilation of the checklist is a first step in mapping of the distribution of all species. It draws upon information given in the original species descriptions as well as later sources that are believed reliable but are not cited in the bibliography. An ideal compilation would be an online database with a map for each species compiled from published collection records and linked directly to a bibliographic entry.

## **Habits and habitats**

A checklist by itself reveals nothing about how the insects live. The ultimate in autecology is the numerical assessment of the dynamics of populations. An intermediate step

is the study of habitat and behavior, by which can be learned the kinds of limitations to population size. This kind of information is sketchy for most coastal staphylinids, and is hard to present in tabular form. Therefore, this section presents textual information for genera and species whose habitat and behavior are known. Although this information is fundamental to population dynamics, it is also useful for the purposes of zoogeography.

Works on seashores and their fauna at large have traditionally been published by marine biologists. If we wanted to learn about the identification of the fauna on European shores, we might consult Barrett and Yonge (1958) or its replacement (Hayward et al. 1996), but we would be disappointed in coverage of Staphylinidae. If we wanted to read about sandy beaches and their fauna, we might consult McLachlan and Brown (2006), or Little et al. (2009) for rocky shores, or Hogarth (2007) for mangroves and seagrasses. Although we might learn much about those environments, we would again be disappointed in coverage of Staphylinidae. Not until we consult Cheng (1976) for marine insects do we gain an appreciation that many staphylinid species dwell on seashores (the included chapter by Moore and Legner), or Morris et al. (1980) for intertidal invertebrates of California (the included chapter by Evans), or Sherwood et al. (2000) for British coastal invertebrates (the included chapter by Hammond). The fault, if there is one, is the slowness in development of comprehensive treatment by entomologists of coastal staphylinids, and that the groundwork (taxonomic and behavioral studies) may not be in publications that marine biologists usually consult.

Habitats of coastal staphylinids are drifted seaweed, the intertidal zone, sandy beaches, pebble beaches, rocky shores, muddy beaches and flats, salt marshes, and mangrove swamps (Figs 2–5), but these are not necessarily mutually exclusive. Staphylinidae are very often associated with plants and algae. These may be growing in estuaries (especially marshgrasses, such as *Spartina*), on rocky shores (green, brown, and red algae), or on sandy shores (diatoms). Or they may be plants or seagrasses (*Thalassia*, *Zostera*, etc.) torn loose and deposited on any of those shores. Few staphylinid studies have identified associated plants specifically or even generically. Often, the drifted seaweeds (i.e., seaweeds deposited onshore by the tide) are called ‘wrack’. Backlund (1945) used the term to include not only the brown alga *Fucus*, but many other algae, and the sea-grass *Zostera* (a vascular plant) deposited onshore. Barrett and Yonge (1958), in contrast, referred to large brown algae including the genus *Laminaria* as ‘kelp’, confining the term ‘wrack’ to moderately sized brown algae including *Pelvetia*, *Fucus*, and *Ascophyllum*. They apply these terms to growing seaweed as well as to seaweed deposited onshore. Moore and Legner (1973) defined wrack as drifted kelp. We employ the term ‘drifted seaweed’ to all algae deposited on shorelines, and we distinguish sea-grasses and marsh-grasses.


Drifted seaweed may be ephemeral and sparse and greatly subject to drying, to deep and decomposing and more or less permanent depending upon location and tides. Backlund (1945) published an ecological study of drifted seaweed in more or less permanent beds on seashores in Sweden and Finland, including its insect inhabitants. Staphylinids may occupy the sand under sparse accretions of seaweed. In thick seaweed beds, staphylinids inhabit the seaweed together with other invertebrates. Organisms






**Figure 2.** Habitats of coastal Staphylinidae. **A** Seagrasses on a sandy beach in Florida, USA **B** Seaweeds on a sandy beach in Jejudo Island, Korea **C** Kelp on a rocky shore in Greymouth, New Zealand **D** Rocky shore in San Diego, USA **E** Seaweeds on rocks of a sandy beach in California, USA **F** Rock crevice on a rocky shore in Plymouth, England **G** Rock covered with barnacles in Baeksu, Korea **H** Close-up of barnacles.





**Figure 3.** Habitats of coastal Staphylinidae. **A** Estuary of Carmel River in California, USA **B** Salt marsh in Florida, USA **C** Mangrove forests in Cat Ba National Park, Vietnam **D** Under stones of mangrove in Cat Ba National Park, Vietnam **E** Mud flat in Gungpyongri, Korea **F** Pebbles and rocks on beach in Kaikoura, New Zealand **G** Seagrasses on a sandy beach in Haiphong, Vietnam **H** Dead fishes on a sandy beach in Florida, USA.





**Figure 4.** Coastal Staphylinidae. **A** *Aleochara zerchei* on a sandy beach in Donghae, Korea **B** *Aleochara puetzi* on a sandy beach in Donghae, Korea **C** *Bryothinusa koreana* under a stone on rocky headland in Dangjin, Korea **D** *Atheta tokiokai* on a sandy beach in Jejudo Island, Korea **E** *Paramblopusa borealis* under a stone on pebble beach in Alaska, USA **F** *Diaulota aokii* with barnacles on a rocky shore in Baeksu, Korea **G** *Diaulota aokii* with fresh seaweeds on a rocky shore in Jejudo Island, Korea **H** Larva of *Diaulota aokii* with barnacles on a rocky shore in Baeksu, Korea.





**Figure 5.** Coastal Staphylinidae. **A** *Phucobius simulator* on a sandy beach in Guryongpo, Korea **B** *Liusus hilleri* on a sandy beach in Donghae, Korea **C** Overwintering staphylinine species (*Cafius histrio*, *Liusus hilleri* and *Philonthus nudus*) under a wooden board on a sandy beach in Jindo Island, Korea **D** Larva of *Cafius* sp. under fresh seaweeds on a sandy beach in Jejudo Island, Korea **E** *Cafius bistriatus* on a sandy beach in North Carolina, USA **F** *Cafius seminites* under decaying seaweeds on a sandy beach in California, USA **G** *Cafius rufescens* on a sandy beach in Jindo Island, Korea **H** *Philonthus nudus* under decaying seaweeds on a sandy beach in Jindo Island, Korea.

that eat and decompose seaweed are more abundant, including amphipods (Crustacea) and seaweed flies (Diptera: Coelopidae), both fed upon by many predacious Staphylinidae, including *Cafius* spp. Amphipods (commonly called ‘beach-hoppers’ or ‘beach-fleas’) are innocuous with respect to humans, but coelopids, (i.e., *Coelopa* spp.) are more problematic. “The flies normally pass their whole lives on the weed [and] are abundant in the wrack all the year around [and] their numbers are not appreciated until something makes them come out into the open [and] they may fly in a band, a little above the ground [and] occasionally they stray far inland [and] can be a great nuisance when large numbers... congregate in shops, garages, and particularly dry-cleaners’ (Oldroyd 1965: 173–174).” By feeding on these flies, *Cafius* spp. and other staphylinids can be considered as beneficial species. Poleward movement of seaweed deposits and their coelopid inhabitants has been noted and attributed to global warming, and there is a suspicion that *Laminaria* is in decline and that *Fucus* is expected to decline in the future in the British Isles (Edward et al. 2007).

Rocky shores offer refuges for specialist staphylinids (those not found in other habitats) in crevices or empty barnacle shells that trap pockets of air. Such shorelines often support the growth of algae, and these algae exhibit zonation according to species (Jones 1968). Staphylinids likewise distribute themselves according to such zones where they find refuge among algal holdfasts (Jones 1968; Topp and Ring 1988b). Rocky shores not only provide a substrate for barnacles and living algae, with which some staphylinids are associated, but they also frequently include tidal pool habitats. Tidal pools are inhabited by a few staphylinids, e.g., *Rothium*. A few species such as *Micralymma marinum* (which is believed to prey on Collembola) attain higher elevations above sea level on rocky cliffs (Thayer 1985).

Coral reefs, even five km from the shoreline, are habitat for a species of *Brachyprotonomaea* (Sawada 1956). Its food there remained unknown until the abundant Collembola present at one such site suggested a probable food source (Ahn et al. 2003). Off the coast of the Aru Islands in the Indonesian Archipelago, among coral polyps, Fauvel (1878a) found unusual staphylinids and described the genera *Corallis* and *Polypea*. Their food and way of life have not yet been determined.

Shores may be graded from solid rock to boulders, cobbles, pebbles, gravels, and sand, and even finer particles typical of mud flats, salt marshes, and mangrove swamps. All of those substrates have their complement of coastal Staphylinidae.

Table 1 is our attempt to summarize information about habitats across all genera; it is incomplete because the available information is incomplete. We summarize existing information not just about habitats but also about behavior and physiology in the section on Habits, Habitats, and Classificatory Notes under the name of each genus.

## Zoogeography

Based on taxonomy and distribution, we provide some provisional ideas about the dispersion of the taxa.

**Table 1.** Genera of the Staphylinidae containing coastal species with their known numbers and habitats.

Subfamily	Tribe	Genus	No. species	Tidal zone	Habitat	Microhabitat
Microsilphinae		<i>Microsilpha</i>	1	not known	sand spit	not known
Omaliinae	Aphaenostemmini	<i>Giulianium</i>	3	HM	SB	UD, UP
		<i>Crymus</i>	2	not known	not known	US
	Omaliini	<i>Macralymma</i>	1	not known	SB	US
		<i>Microlymma</i>	2	ML	RH	RC
		<i>Omaliomimus</i>	10	ML	not known	US
		<i>Omalius</i>	4	not known	not known	US, UG, UD
		<i>Arbromelus</i>	1	HM	MA	UD
		<i>Barriscenites</i>	2	HM	MA	UD
		<i>Barriscenus</i>	1	HM	MA	UD
		<i>Brachygluta</i>	6	not known	SM	UG, UP
Pselaphinae	Barrisini	<i>Briana</i>	1	PH	MA	UP, UD
		<i>Brianaxis</i>	1	not known	not known	UD, UP, US
	Brachyglutini	<i>Mangalobothrus</i>	3	HM	MA	cavities in log
		<i>Nisaxis</i>	2	not known	SM	UD, UG
		<i>Pedinops</i>	1	not known	coral reef	not known
		<i>Physoplectus</i>	4	not known	not known	UP
		<i>Prosthecarthron</i>	1	not known	MF, SM	UP
		<i>Aleochara</i>	16	PH, HM	SB, RH	US, UD
		<i>Acticola</i>	1	not known	not known	US
		<i>Adota</i>	6	PH	SM, SB	US, UD
		<i>Atheta</i>	6	PH	SM, ES, SB	US, UD
		<i>Brundinia</i>	2	not known	SM, ES	UD
Aleocharinae	Aleocharini					
	Athetini					

Subfamily	Tribe	Genus	No. species	Tidal zone	Habitat	Microhabitat
		<i>Halobrecta</i>	7	not known	SM, ES	US, UD
		<i>Hydrosmeeta</i>	1	PH	SB	US
		<i>Iotarphia</i>	1	not known	not known	not known
		<i>Osakatheta</i>	1	HM	ES, MF	UP
		<i>Pontomalota</i>	2	PH, HM	SB	US, UD
		<i>Psammopora</i>	1	not known	not known	not known
		<i>Psammotibia</i>	5	PH, HM	SB	US
		<i>Tarphiota</i>	3	PH, HM	SB	US, UD
		<i>Thinusa</i>	2	PH, HM	SB	US, UD
	Diglotini	<i>Diglotia</i>	8	PH	SB, SM, ES, BS	UP, UB
	Falagriini	<i>Bryobiota</i>	2	PH, HM	SB	US, UD
		<i>Myrmecopora</i>	14	PH, HM	SB	US, UD
	Homalotini	<i>Cameronium</i>	5	HM	RH	RC, US
		<i>Heterota</i>	10	PH	SB	US, UD
		<i>Linoglossa</i>	1	not known	MA	not known
		<i>Paractocharis</i>	3	PH	SB	US
		<i>Pseudopasilia</i>	1	not known	BS	UP, US
		<i>Thirubiosus</i>	1	not known	SB	US
	Liparocephalini	<i>Amblopusa</i>	5	HM	BS	UP, UB, US
		<i>Baeostethus</i>	1	PH	SB	UP
		<i>Diaulota</i>	8	ML, VL	RH	EB, RC, RA
		<i>Halorhadinus</i>	3	HM	BS	UP, US
		<i>Ianmoorea</i>	1	PH	SB	UB
		<i>Liparocephalus</i>	4	ML, VL	RH	RC, RA

Subfamily	Tribe	Genus	No. species	Tidal zone	Habitat	Microhabitat
Oxytelinae	Myllaenini	<i>Paramblopusa</i>	2	HM	SB	UP
		<i>Brachtypronmaea</i>	4	ML	RH	coral reef
		<i>Bryoibinus</i>	30	HM, ML	SB, BS, MF, MA	UP, UB, RC, EB
		<i>Corallis</i>	1	VL	not known	under coral polyp
		<i>Lautaea</i>	1	not known	MA, MF	not known
		<i>Myllaena</i>	1	PH	BS	UP
		<i>Polypea</i>	1	VL	not known	under coral polyp
		<i>Rothium</i>	6	HM, ML	RH	RA
	Oxypodini	<i>Chilodera</i>	1	not known	not known	US
		<i>Dasydera</i>	1	not known	not known	US
		<i>Gyronotus</i>	1	not known	not known	not known
		<i>Oreuryalea</i>	1	PH	SB	US, UD
	Phytosini	<i>Actocharis</i>	2	PH	SB	UB
		<i>Arena</i>	2	PH	SB, MF	UB
		<i>Euphytosus</i>	1	not known	not known	not known
		<i>Phytosus</i>	8	not known	SB	UB, US, UD
Oxytelinae	Incertae sedis	<i>Salinamexus</i>	3	PH	SB	US, UD, UP
		<i>Anoryllus</i>	1	PH	SB	US, UD
	Oxytelini	<i>Blediorogus</i>	4	PH	SB	US
		<i>Pareiobledius</i>	3	PH	SB	US
		<i>Sartallus</i>	1	PH	SB	US, UD
		<i>Bledius</i>	57	PH, HM	MF, SM, ES	UB, US, UD
	Thinobiini	<i>Carpelimus</i>	1	PH	MF, SM, ES	UB, US, UD
		<i>Teropalpus</i>	9	PH	SB	US, UD

Subfamily	Tribe	Genus	No. species	Tidal zone	Habitat	Microhabitat
Scydaeninae	Cephenniniini Paederini	<i>Thinobius</i>	3	PH, HM	BS, SB, SM, MF	US
		<i>Cephennodes</i>	1	PH	BS	UP
		<i>Chetocephalus</i>	1	PH	SB	US
		<i>Medon</i>	4	PH	SB	US, UD
		<i>Ophioomma</i>	1	PH	SB	UD
		<i>Sunius</i>	2	PH	SB, MA	US
		<i>Bisnius</i>	1	PH	SB	US
		<i>Caftus</i>	44	PH	SB	US, UG, UD
		<i>Gabronthus</i>	1	not known	not known	not known
		<i>Hadropinus</i>	1	PH	SB	US
Staphylininae	Staphylinini	<i>Hadrotes</i>	2	PH	SB	US
		<i>Heterothops</i>	1	PH	SB	US, UD
		<i>Liusus</i>	2	PH	SB	US, UG, UD
		<i>Orthidus</i>	1	PH	SB	US, UP
		<i>Philonthus</i>	1	PH	SB, SM	US, UG, UD
		<i>Phucobius</i>	8	PH	SB	US, UG, UD
		<i>Quediocafus</i>	3	PH	SB	UG
		<i>Remus</i>	4	PH	SB	US, UG, UD
		<i>Thinocafus</i>	1	PH	SB	US
		<i>Thinopinus</i>	1	PH	SB	US

PH: proximal to high tide zone; HM: high to mid tide zone; ML: mid to low tide zone; VL: very low tide zone. BS: boulder shores (band of gravels/pebbles/cobbles); ES: estuary; MA: mangrove; MF: mud/sand flats; RH: rocky headland; SB: sandy beach; SM: salt marsh. EB: Inside of empty barnacles/shells; RA: rock with algae; RC: rock crevices; UB: under beach sand; UD: under debris; UG: under seagrasses; UP: under stones (gravels/pebbles/cobbles); US: under seaweeds.



## A Checklist of coastal Staphylinidae and their distribution

### MICROSILPHINAE

*Microsilpha* Broun 1886: 889

*M. litorea* Broun 1886: 890 - SPO (NZ)

### OMALIINAE

#### APHAENOSTEMMINI

*Giulianium* Moore 1976: 56 [rev Ahn and Ashe 1999]

*G. alaskanum* Ahn and Ashe 1999: 162 - NPO (US: AK; JP: HK)

*G. campbelli* Moore 1976: 57 - NPO (US: CA)

*G. newtoni* Ahn and Ashe 1999: 163 - NPO (US: CA)

#### OMALIINI

*Crymus* Fauvel 1904b: 92

= *Arpediopsis* Cameron 1917a: 124

= *Arpediomimus* Cameron 1917f: 277

*C. antarcticus* Fauvel 1904b: 93 - SAO (South Georgia; Falkland Islands)

= *falklandicus* (Cameron) 1917a: 125

*C. kronii* (Kiesenwetter) 1877: 161 SPO (NZ: Antipodes Island, Auckland Island, Campbell Island, South Island)

= *longiceps* (Broun) 1914: 89

*Macralymma* Cameron 1945c: 179

*M. punctiventre* Cameron 1945c: 179 - SPO (NZ)

*Micralymma* Westwood 1838: 129

*M. marinum* (Strøm) 1783: 65 - NAO (CA: NB, NL, NS, QC; US: MA, ME, NH; FR; GB: England, N. Ireland, Scotland; GL; IE; IS; NO), North and Baltic and Irish Seas (BE; DE; GB: England, Scotland, N. Ireland, Wales; NL; SE; RU: Karelia)

= *brevipenne* (Gyllenhal) 1810: 234

= *johnstonis* Westwood 1838: 130

= *stimpsonii* LeConte 1863: 57

*M. laticolle* Motschulsky 1860: 549 ACO (RU: Siberia) [probably does not belong to *Micralymma*]

*Omaliumimus* Jeannel 1940: 117

*O. actobius* (Broun) 1893: 1035 - SPO (NZ)

*O. albipennis* (Kiesenwetter) 1877: 162 - SPO (AU: Macquarie Island; NZ: Campbell Island, Auckland Island)

= *variipennis* (Lea) 1920: 30

= *flavipennis* Cameron 1948: 723

*O. carinigerus* (Broun) 1893: 1036 - SPO (NZ)

*O. chalmeri* (Broun) 1893: 1037 - SPO (NZ)

*O. conicus* (Fauvel) 1878b: 484 - SPO (NZ)

- O. laetipennis* (Broun) 1910: 19 - SPO (NZ)  
*O. litoreus* (Broun) 1886: 942 - SPO (NZ)  
*O. robustus* (Broun) 1911: 96 - SPO (NZ: Chatham Islands, Pitt Island)  
*O. setipes* (Broun) 1909b: 230 - SPO (NZ)  
*O. venator* (Broun) 1909a: 98 SPO (AU: Macquarie Island; NZ: mainland, Antipodes Island, Auckland Island, Campbell Island, Snares Island)  
*Omalius* Gravenhorst 1802: 111  
*O. algarum* Casey 1885: 316 - NPO (CA: BC; US: CA, OR)  
*O. laeviusculum* Gyllenhal 1827: 464 NAO (FR; GB: England, Scotland; IE; IS; NO), ACO (RU), North Sea (BE; DE; DK; GB: England, Scotland; NL), Irish Sea (GB: England, Wales; IE), Baltic Sea (FI; SE)  
*O. riparium* Thomson 1857: 224 - NAO (ES; FR; GB: England, Scotland; MA; PT; DK: Faroes), Irish Sea (GB: England, Wales; IE), North Sea (BE; DE; DK; FR; GB: England, Scotland; NL), Baltic Sea (DE; DK; EE; FI; PL; SE), Mediterranean Sea (DZ; ES; FR; IT; MA; YU)  
*O. rugulipenne* Rye 1864: 58 - NAO (GB: England, Scotland), Irish Sea (GB: England; IE), North Sea (BE; DE; FR; GB: England, Scotland; NL)

## PSELAPHINAE

### BATRISITAE

#### BATRISINI

- Arthromelus* Jeannel 1949: 149  
*A. quadratus* Tanokuchi 1989: 88 - South China Sea (SG)  
*Batriscenites* Jeannel 1952: 96  
*B. celer* Tanokuchi 1989: 91 - South China Sea (SG)  
*B. humicola* Tanokuchi 1989: 95 - South China Sea (SG)  
*Batriscenus* Raffray 1903: 48  
*B. foveiterminalis* Tanokuchi 1989: 97 - South China Sea (SG)

## GONIACERITAE

### BRACHYGLUTINI

- Brachygluta* Thomson 1859: 54  
*B. abdominalis* (Aubé) 1833: 27 - NAO (eastern US; CA: NB, NS)  
*B. cavicornis* (Brendel) 1865a: 30 - NAO (eastern US: NY, MD, DC, VA)  
*B. curvicera* (Motschulsky) 1854: 4 - NAO (eastern US: NY)  
*B. floridana* (Brendel) 1865b: 257 - NAO (eastern US: NY, MD, VA, NC, SC, FL)  
*B. luniger* (LeConte) 1849: 87 - NAO (eastern US: MA, NY, NJ, MD, VA)  
*B. ulkei* (Brendel) 1866: 193 - NAO (eastern US: MD, DC, DE, VA, SC, GA, FL)  
*Briara* Reitter 1882: 207  
     = *Gonatocerus* Schaufuss 1880: 30 [preoccupied]  
     = *Berlara* Reitter 1882: 206

- B. bella* (Tanokuchi) 1989: 101 - South China Sea (SG)
- Briaraxis* Brendel 1894: 158
- B. depressa* Brendel 1894: 159 - Caribbean Sea (US: FL; TT: Tobago, Costa Rica)
- Mangalobythus* Tanokuchi 1989: 104
- M. acutifolius* Tanokuchi 1989: 109 - South China Sea (TH)
- M. furcifer* Tanokuchi 1989: 106 - South China Sea (SG)
- M. murphyi* Tanokuchi 1989: 111 - South China Sea (SG)
- Nisaxis* Casey 1886: 183
- N. maritima* Casey 1887: 468 - Gulf of Mexico (US: LA, MS, TX)
- N. tomentosa* (Aubé) 1833: 33 - NAO (eastern US: CT, NY, NJ, DE, MD, DC, NC), Gulf of Mexico (FL, AL, MS, TX), Caribbean Sea
- = *minuta* (Brendel) 1865a: 30
- = *cinninata* Casey 1887: 466
- Pedinopsis* Newton and Chandler 1989: 43
- = *Pedinopsis* Raffray 1890: 102
- = *Halohermatus* Sawada 1991: 148
- P. regulus* (Sawada) 1991: 150 - NPO (JP: RY)
- Physoplectus* Reitter 1882: 210
- = *Halorabyxis* Jeannel 1954: 338
- = *Thalassomerus* Sawada 1992: 55
- P. irritans* Chandler 2001: 349 - SPO (AU: Queensland)
- P. miyakei* (Sawada) 1992: 58 - NPO (JP: RY)
- P. reikoeae* (Sawada) 1992: 56 - NPO (JP: HN)
- P. vinsoni* (Jeannel) 1954: 341 - INO (MU)
- Prosthecarthron* Raffray 1915: 2
- P. sauteri* Raffray 1915: 3 - NPO (KP; JP: HN, KY, SH, RY), SPO (TW; VN)
- = *palpalis* (Löbl) 1974: 97; 1977: 236

## ALEOCHARINAE

### ALEOCHARINI

- Aleochara* Gravenhorst 1802: 67 [rev Nearctic Klimaszewski 1984, southern Africa Klimaszewski and Jansen 1994, Palearctic Assing 1995]
- A. (Coprochara) salsipotens* Bernhauer 1912c: 209 - SAO (NA; ZA), INO (ZA)
- A. (Coprochara) squalithorax* Sharp 1888: 282 - East Sea (JP: HN; KR)
- A. (Coprochara) sulcicollis* Mannerheim 1843: 225 NPO (CA: BC; US: AK, CA, OR, WA; MX: BN, BS), SPO (CL)
- A. (Emplenota) albopila* (Mulsant and Rey) 1852: 171 - Mediterranean Sea (FR; GR; IT; YU), Black Sea (BG), NAO (ES: Canary Islands)
- A. (Emplenota) curticens* Klimaszewski 1984: 101 - NPO (CA: BC; US: CA)
- A. (Emplenota) fucicola* Sharp 1874: 9 - NPO, South China Sea, East Sea (CN: Hong Kong; JP: HN; KR)
- = *variolosa* Weise 1877: 89

- A. (Emplenota) litoralis* (Mäklin) 1853: 182 - NAO (CA: NB, NL, NS, QC; US: FL, MA, NJ, NY, RI), NPO (CA: BC; US: AK, CA; MX: BN, BS, SO)
- A. (Emplenota) obscurella* Gravenhorst 1806: 159 NAO (ES; FR; GB: England, Scotland; IE; NO), Irish Sea (GB: England, Wales; IE), North Sea (BE; DE; DK; GB: England, Scotland; NL; NO), Baltic Sea (DE; DK; PL; SE), Irish Sea (GB; IE)  
= *algarum* Fauvel 1862b: 92
- A. (Emplenota) pacifica* (Casey) 1894: 290 NPO (CA: BC; US: CA, WA; MX: BN)
- A. (Emplenota) phycophila* Allen 1937: 219 - NAO (GB)
- A. (Emplenota) puetzi* (Assing) 1995: 225 - East Sea (KR; RU: Sakhalin, Kamchatka)
- A. (Polystomota) grisea* Kraatz 1856: 96 - NAO (FR; GB: England, Scotland; IE; MA; NO; PT), Irish Sea (GB: Wales), North Sea (BE; DE; DK; GB: England, Scotland; NL; NO), Baltic Sea (DE; DK; FI; PL; SE), Mediterranean Sea (DZ; ES; FR; IT; YU)
- A. (Polystomota) punctatella* Motschulsky 1858b: 240 - NAO (England, Scotland; IE), Irish Sea (GB: England, Wales), North Sea (BE; GB: England, Scotland; NL), NAO (FR)
- A. (Triochara) nubis* (Assing) 1995: 232 - East Sea (RU: Sakhalin, Kamchatka)
- A. (Triochara) trisulcata* Weise 1877: 88 - NPO, East Sea (CN: Hong Kong; JP: HN; KR)
- A. (Triochara) zerchei* (Assing) 1995: 231 - East Sea (KR; RU: Primorie, Sakhalin)

## ATHETINI

*Acticola* Cameron 1944b: 618

*A. falkandica* Cameron 1944b: 619 - SAO (Falkland Islands)

*Adota* Casey 1910: 67 [rev Gusarov 2003a,b]

= *Panalota* Casey 1910: 71

= *Phyconoma* Easton 1971: 24

= *Halostiba* Yosii and Sawada 1976: 86

*A. colpophila* Gusarov 2003b: 16 - Gulf of California (MX: SO)

*A. gnyptoides* (Casey) 1910: 69 - NPO (US: AK, CA)

*A. madida* (Bernhauer) 1907: 400 - NPO (JP: HN, KY)

*A. magnipennis* (Bernhauer) 1943: 184 - NPO, East Sea (JP: HN, KY; KR)

*A. maritima* (Mannerheim) 1843: 224 - NPO (CA: BC; US: AK, CA), NAO (GB: England), North Sea (GB: England, Scotland)

= *massettensis* Casey 1910: 68

= *subintima* Casey 1910: 68

= *setosetarsis* Casey 1910: 71

= *insons* Casey 1911: 125

- = *scolopacina* Casey 1911: 124
- = *scortea* Casey 1911: 124
- = *immigrans* (Easton) 1971: 25 - North Sea (GB), SPO (NZ)
- A. ushio* (Sawada) 1971a: 304 - NPO, East Sea (JP: HN, KY)
- Atheta* Thomson 1858: 36
  - A. (Actophylla) varendorffiana* Bernhauer and Scheerpeltz [in Scheerpeltz 1934: 1637] - North Sea (DE)
  - = *varendorffi* Bernhauer 1908a: 334 [preoccupied]
  - A. (Badura) ririkoae* Sawada 1989b: 285 - NPO, East Sea (JP: HN; KR)
  - A. (Badura) tokiokai* (Sawada) 1971a: 306 - NPO, East Sea (JP: HN, KY; KR)
  - A. (Datomicra) acadensis* Klimaszewski and Majka 2007: 49 NAO (CA: NB, NS, PE, QC)
  - A. (Sipalatheta) algarum* Pace 1999b: 680 - South China Sea (CN: Hong Kong)
  - A. novaescotiae* Klimaszewski and Majka [in Klimaszewski et al. 2006: 68] NAO (CA: NB, NS, NL, Sable Island; St Pierre and Miquelon: Miquelon)
- Brundinia* Tottenham 1949: 78
  - B. marina* (Mulsant and Rey) 1853: 39 - NAO (FR; GB: England), North Sea (DE; GB: England, Scotland; NL), Baltic Sea (DE; SE), Irish Sea (GB: England, Wales; IE), Mediterranean Sea (FR; IT)
  - = *imbecilla* (G. Waterhouse) 1858: 6074 (and 1859: 16)
  - = *thinobia* Thomson 1861: 73
  - B. meridionalis* (Mulsant and Rey) 1853: 38 - NAO (FR; GB: England), Irish Sea England, Wales), North Sea (BE; GB: England), Mediterranean Sea (FR; IT)
- Halobrecta* Thomson 1858: 35 [rev Gusarov 2004, mainly Nearctic]
  - H. algae* (Hardy) 1851: 78 - Baltic Sea (DK; EE; FI; RU: Karelia; SE), North Sea (BE; GB: England, Scotland), NAO (FR; NO), SPO (AU)
  - = *puncticeps* (Thomson) 1852: 134
  - = *anthracina* Fairmaire 1852: 687 [synonymy based on statement by Fairmaire 1856: 424 but not otherwise verified]
  - H. algophila* (Fenyès) 1909: 419 - NPO (US: CA), SPO (NZ; CL: Palena), SAO (Tristan da Cunha: Inaccessible Island), NAO (GB), Mediterranean Sea (FR: Corsica)
  - = *barbarae* (Casey) 1910: 18
  - = *importuna* (Casey) 1911: 111
  - H. cingulata* (Cameron) 1920: 266 - South China Sea (SG)
  - = *consors* (Cameron) 1920: 266
  - H. discipula* Pace 1999a: 171 - SPO (AU; CL: Valparaíso)
  - H. flavipes* Thomson 1861: 50 - NAO (NO), Baltic Sea (DK; EE; FI; SE), Mediterranean Sea (IT), North Sea (BE; DE; GB: England, Scotland), NAO (CA: NB; US: NY, VA), SPO (AU; CL: Llanquihue)
  - = *maritima* (G. Waterhouse) 1863: 137
  - = *halobrectha* (Sharp) 1869: 139

- = *pubes* (Mulsant and Rey) 1873a: 660
- = *puncticeps* sensu Mulsant and Rey 1875: 12
- = *pocahontas* (Casey) 1910: 19
- = *vaticina* (Casey) 1910: 19
- = *incertula* (Casey) 1910: 84
- H. halensis* Mulsant and Rey 1873b: 173 - Mediterranean Sea (FR)
- H. princeps* (Sharp) 1869: 142 - NAO (GB: England)
- Hydrosmeeta* Thomson 1858: 33
- H. subalgarum* Pace 1999b: 672 - South China Sea (CN: Hong Kong)
- Iotarphia* Cameron 1943: 352
- I. australis* Cameron 1943: 352 - Tasman Sea (AU: New South Wales)
- Osakatheta* Maruyama, Klimaszewski and Gusarov 2008: 40
- O. yasukoae* Maruyama, Klimaszewski and Gusarov 2008: 41 - NPO (JP: HN)
- Pontomalota* Casey 1885: 296 [rev Ahn and Ashe 1992]
- P. opaca* (LeConte) 1863: 28 - NPO (CA: BC; US: CA, OR, WA; MX: BN)
  - = *californica* Casey 1885: 298
  - = *nigriceps* Casey 1885: 299
  - = *luctuosa* Casey 1911: 164
  - = *bakeri* Bernhauer 1912b: 170
- P. terminalia* Ahn and Ashe 1992: 356 - NPO (US: CA)
- Psammopora* Pace 2003: 154
- P. delittlei* Pace 2003: 157 - Tasman Sea (AU: Tasmania)
- Psammotiba* Yosii and Sawada 1976: 82 [rev Gusarov 2003b]
- P. comparabilis* (Mäklin) 1853: 181 - NPO (CA: BC, US: AK, CA)
- P. hilleri* (Weise) 1877: 90 - NPO, East Sea (JP: HN, KY)
  - = *multipunctata* (Sawada) 1971a: 301
- P. jessoensis* (Brundin) 1943: 22 - NPO (JP: HK, HN; RU: Maritime Territory)
- P. kamtschatica* (Brundin) 1943: 21 - NPO (JP: HK; RU: Kamchatka, Kuril Islands)
- P. kenaii* Gusarov 2003b: 28 - NPO (CA: BC; US: AK, CA)
- Tarphiota* Casey 1894: 332 [rev Ahn 1996b, Ahn 1999, Klimaszewski et al. 2006]
- T. densa* (Moore) 1978a: 115 - NPO (MX: BS, SO)
  - = *hirsutula* Casey 1910: 75
- T. fucicola* (Mäklin) 1852: 306 - NPO (CA: BC), Gulf of California (MX: BC, SO)
  - = *debilicollis* Casey 1910: 75
  - = *pallidipes* Casey 1894: 333 - NPO (US: CA)
- T. geniculata* (Mäklin) 1852: 308 - NPO (CA: BC; US: AK, CA, OR)
  - = *iota* Casey 1910: 76
  - = *insolita* Casey 1910: 76
  - = *litorina* Casey 1910: 75
  - = *seditiosa* Casey 1910: 76

*Thinusa* Casey 1894: 371 [rev Ahn 1997b]

*T. fletcheri* Casey 1906: 353 - NPO (CA: BC; US: AK, CA, OR, WA)

= *divergens* Casey 1911: 213

= *nigra* Casey 1911: 214

= *robustula* Casey 1911: 215

*T. maritima* (Casey) 1885: 312 - NPO (CA: BC; US: CA, OR, WA; MX: BN)

= *obscura* Casey 1906: 354

#### DIGLOTTINI [key Pace 1986]

*Diglotta* Champion 1887: 228 (repeated 1899: 265) [rev Haghebaert 1991]

= *Diglossa* Haliday 1837: 252 [preoccupied]

*D. brasiliensis* Caron and Ribeiro-Costa 2008: 53 - SAO (BR: Paraná)

*D. legneri* Moore and Orth 1979a: 339 - NPO (US: CA)

*D. littoralis* (Horn) 1871: 331 - NAO (US: NJ)

*D. maritima* Lea 1927: 277 - SPO (FJ: Levuka)

*D. mersa* (Haliday) 1837: 252 North Sea (BE; DE; DK; FR; GB: England; NL), Irish Sea (GB: N. Ireland, Wales), Mediterranean Sea (DZ; IT), NAO (CA: NB; GB: England, Scotland; IE)

= *D. submarina* Fairmaire 1856: 468

*D. pacifica* Fenyès 1921: 17 - NPO (US: CA, OR; MX: BN)

*D. secqi* Pace 1992: 180 - Red Sea (DJ)

*D. sinuaticollis* (Mulsant and Rey) 1870: 176 NAO (GB: England), Irish Sea (GB: England, Wales; IE)

= *D. crassa* (Mulsant and Rey) 1870: 180

#### FALAGRIINI [rev Hoebeke 1985, phy Ahn and Ashe 1995]

*Bryobiota* Casey 1894: 367 [rev Ahn and Ashe 1995]

*B. bicolor* (Casey) 1885: 311 - NPO (CA: BC; US: WA, OR, CA; MX: BN)

= *californica* (Scheerpeltz) 1965: 49

*B. giulianii* (Moore) 1978a: 113 - NPO (US: CA, WA)

*Myrmecopora* Saulcy 1864: 429 [rev Assing 1997a,b, Palaearctic]

*M. (Lamproxenus) algarum* (Sharp) 1874: 12 - NPO, East Sea (JP: HN, Tokara Island)

= *miyamotoi* (Sawada) 1955: 85

*M. (Lamproxenus) chinensis* Cameron 1944c: 158 South China Sea (CN: Hong Kong)

*M. (Lamproxenus) reticulata* Assing 1997b: 344 - NPO (RU: Far East; KP)

*M. (Lamproxenus) rufescens* (Sharp) 1874: 11 - NPO, East Sea (JP: HN, KY)

*M. (Paraxenus) laesa* (Erichson) 1839b: 73 Mediterranean Sea (IT; PT; FR; DZ; TN; HR; ES: Balearic Islands), NAO (ES: Canary Islands)

= *tenuicornis* (Küster) 1854: no. 3

*M. (Xenus) anatolica* (Fagel) 1969: 117 - Mediterranean Sea (CY; TR)

*M. (Xenus) bernhaueri* Koch 1936: 210 - Red Sea (EG; IL)



- M. (Xenusa) boehmi* Bernhauer 1910: 259 Mediterranean Sea (FR: Corsica; IT; GR; TN; MA; CY)  
 = *sydowi* Bernhauer 1927b: 97  
 = *mediterranea* Fagel 1970: 152
- M. (Xenusa) brevipes* Butler 1909: 29 NAO (FR; GB: England), Irish Sea (GB: Wales; IE), North Sea (GB: England) [but see Hammond 2000: 257]  
 = *oweni* Assing 1997a: 114 [*fide* Hammond 2000: 257]
- M. (Xenusa) maritima* (Wollaston) 1860: 51 NAO (ES: Canary Islands; PT: Madeira)
- M. (Xenusa) minima* Bernhauer 1901a: 537 Black Sea (BG; RO) Mediterranean Sea (GR; MA)  
 = *buressi* Rambousek 1910: 19 [in Czech] and 21 [in French]  
 = *pamphylica* (Fagel) 1969: 120
- M. (Xenusa) simillima* (Wollaston) 1864: 534 NAO (ES: Canary Islands; GB: England, Scotland; IE; NO), Baltic Sea (DK; DE), North Sea (GB: England, Scotland), Mediterranean Sea (FR; ES; PT; TN; DZ; EG), NAO (PT: Azores)  
 = *lobmanderi* Bernhauer 1927a: 167
- M. (Xenusa) sulcata* (Kiesenwetter) 1850: 218 Mediterranean Sea (AL; GR; FR: Corsica; IT: Sardinia, Sicily; HR), North Sea (GB), Black Sea (RO; BG; UA)  
 = *carica* Fagel 1970: 155
- M. (Xenusa) uwida* (Erichson) 1840: 916 NAO (GB: England), Mediterranean Sea (AL; BG; CY; ES; GR; TN; IT; HR; YU), North Sea (BE; DE; GB: England; NL), NAO (FR), Black Sea (UA; GE)  
 = *meridiogallica* Scheerpeltz 1972: 101

## HOMALOTINI

### *Cameronium* Koch 1936: 202

- C. flavipenne* Cameron 1944a: 318 - INO (SO; TZ: Zanzibar)  
*C. gomyi* Pace 1985: 622 - INO (Comoros)  
*C. lamuense* Pace 1994: 155 - INO (KE: Lamu)  
*C. obockianus* (Fauvel) 1905: 146 - Red Sea [DJ; ET; YE: Barim (= Perim) Island]  
*C. sonorensis* Moore 1964a: 175 - Gulf of California (MX: SO)

### *Heterota* Mulsant and Rey 1874: 194 [rev Park et al. 2008]

- H. arenaria* Cameron 1920: 251 - South China Sea (SG)  
*H. brevicollis* (Bernhauer) 1929: 187 - Red Sea [YE: Barim (= Perim) Island]  
*H. gomyi* Jarrige 1973: 257 - INO (MG)  
*H. obscura* Cameron 1938: 174 - INO (RE)  
*H. pamphylica* Fagel 1969: 123 - Mediterranean Sea (TR)  
*H. pictipennis* (Fauvel) 1905: 142 - Red Sea (DJ; ET), INO (SO)



- H. plumbea* (G. Waterhouse) 1858: 6074 (and 1859: 15) - Mediterranean Sea (Europe), NAO (ES: Canary Islands; GB: England; US: FL), Irish Sea (GB: Wales), Caribbean Sea (JM; MX: QR)  
 = *fairmairii* (Brisout) 1859: ccxviii  
 = *godelinaisei* (Fauvel) 1862b: 92  
 = *trogophloeoides* (Wollaston) 1864: 536  
 = *impressa* (Mulsant and Rey) 1875: 459  
*H. rougemonti* Pace 1993: 137 - Bali Sea (ID: Bali)  
*H. sunjaei* Park and Ahn [in Park et al.] 2008: 111 - NPO (KR)  
*H. vinsoni* Cameron 1947a: 118 - INO (MG; MU; RE, Comoros)  
*Linoglossa* Kraatz 1859a: 10  
*L. murphyi* Sawada 1991: 142 - South China Sea (SG)  
*Paractocharis* Cameron 1917c: 154  
*P. deharvengi* Pace 1990: 81 - Luzon Sea (PH: Mindoro)  
*P. fucicola* Cameron 1917c: 155 - South China Sea (SG)  
*P. orousseti* Pace 1990: 79 - Luzon Sea (PH: Mindoro)  
*Pseudopasilia* Ganglbauer 1895: 211  
*P. testacea* (Brisout) 1863: 16 NAO (FR; GB: England), North Sea (BE), Mediterranean Sea (FR: mainland, Corsica; HR; IT; TN)  
*Thinobiosus* Moore and Legner 1977: 468  
*T. salinus* Moore and Legner 1977: 469 - Gulf of California (MX: SO)

## LIPAROCEPHALINI [phy Ahn and Ashe 1996b, Ahn et al. 2010]

- Amblopusa* Casey 1894: 355 [rev Ahn and Ashe 1996a, Zerche 1998]  
 = *Boreorhadinus* Sawada 1991: 147  
*A. alaskana* Ahn and Ashe 1996a: 143 - NPO (US: AK)  
*A. brevipes* Casey 1894: 356 - NPO (CA: BC; US: AK, CA)  
 = *pallida* Casey 1911: 212  
*A. hokkaidona* Ahn and Ashe 1996a: 142 - NPO (JP: HK)  
*A. magna* Zerche 1998: 106 - NPO (JP: HK; RU: Kuril Islands)  
*A. pacifica* (Sawada) 1991: 147 - NPO (JP: HK)  
*Baeostethus* Broun 1909a: 96 [rev Leschen et al. 2002]  
*B. chiltoni* Broun 1909a: 97 - SPO (NZ: Campbell, Auckland, Antipodes Island)  
*Diaulota* Casey 1894: 354 [rev Ahn 1996a, Zerche 1998]  
 = *Genoplectes* Sawada 1955: 81  
*D. alaskana* Ahn 1996a: 278 - NPO (US: AK)  
*D. aokii* Sawada 1971b: 104 - NPO (JP: HK, HN; KR; US: AK)  
*D. densissima* Casey 1894: 354 - NPO (CA: BC; US: CA, OR, WA)  
 = *D. insolita* Casey 1894: 355  
*D. fulviventris* Moore 1956a: 120 - NPO (US: CA; MX: BN)  
*D. harteri* Moore 1956a: 123 - NPO (US: CA; MX: BN)  
 = *D. megacephala* Moore 1956a: 124

- D. pacifica* Sawada 1971b: 101 - NPO (JP: HN; KR)  
*D. uenoi* (Sawada) 1955: 82 - NPO (JP: HN, RY; KR)  
*D. vandykei* Moore 1956a: 125 - NPO (US: CA)  
*Halorhadinus* Sawada 1971b: 92 [rev Ahn 2001]  
*H. aequalis* Sawada 1971b: 92 - NPO, East Sea (JP: HN; KR)  
*H. inaequalis* Sawada 1971b: 95 - NPO, East Sea (JP: HN; KR)  
*H. sawadai* Maruyama and Hayashi 2009: 72 - East Sea (JP: HN)  
*Ianmoorea* Ahn 2006: 36  
     = *Moorea* Ahn 2004: 255  
*I. zealandica* (Ahn) 2004: 258 - SPO (NZ: North Island, South Island)  
*Liparocephalus* Mäklin 1853: 191 [rev Ahn 1997a, Maruyama and Ahn 2000b]  
*L. brevipennis* (Mäklin) 1853: 192 - NPO (US: AK)  
*L. cordicollis* LeConte 1880: 177 - NPO (CA: BC; US: AK, CA, OR, WA)  
*L. litoralis* Kirschenblatt 1938: 532 - NPO (RU: Kuril Islands; JP: HK)  
*L. tokunagai* Sakaguti 1944: 20 - NPO (JP: SH, KY)  
*Paramblopusa* Ahn and Ashe 1996a: 148 [rev Ahn and Ashe 1996a, Maruyama and Ahn 2000a]  
*P. borealis* (Casey) 1906: 354 - NPO (CA: BC; US: AK, OR, WA; JP: HK)  
*P. eoa* Ahn and Maruyama 2000: 359 - NPO (RU: Kuril Islands)

#### MYLLAENINI [phy Ahn and Ashe 2004]

- Brachypronomaea* Sawada 1956: 197 [rev Ahn et al. 2003]  
     = *Thalassopora* Jarrige 1959: 63  
*B. esakii* Sawada 1956: 197 - NPO (JP: RY)  
*B. marchemarchadi* (Jarrige) 1959: 64 - South China Sea (VN)  
*B. nosybiana* (Jarrige) 1959: 65 - INO (MG)  
*B. sawadai* Jarrige 1964: 178 - SPO (NC)  
*Bryothinusa* Casey 1904: 312 [rev Pace 1986, Haghebaert 1995, Ashe 2005]  
     = *Halesthenus* Sawada 1955: 83  
*B. algarum* Sawada 1971b: 90 - NPO (JP: HN, KY)  
*B. cameroni* (Fauvel) 1904a: 74 - Red Sea [ER; YE: Kameron (= Cameran) Island, Barim (= Perim) Island]  
     = *B. microphthalma* (Bernhauer) 1929: 187  
*B. catalinae* Casey 1904: 313 - NPO (US: CA)  
*B. celebensis* (Fauvel) 1878a: 301 - Celebes Sea (ID: Sulawesi)  
*B. chani* Moore and Legner 1971: 107 - South China Sea (CN: Hong Kong)  
*B. chengae* Ahn 1998: 335 - SPO (Caroline Island: Palau)  
*B. fluenta* Moore and Legner 1975: 111 - South China Sea (CN: Hong Kong)  
*B. gangjinensis* Ahn and Jeon 2004: 29 - NPO (KR)  
*B. grootaerti* Haghebaert 1995: 29 - Bismarck Sea (PG: Laing Island)  
*B. hauseri* Ashe 2005: 582 - South China Sea (MY: Malaya)  
*B. hongkongensis* Moore, Legner and Chan 1973: 77 South China Sea (CN: Hong Kong)

- B. koreana* Ahn and Jeon 2004: 31 - NPO (KR)  
*B. madecassa* Pace 2008: 568 - INO (MG)  
*B. minuta* (Sawada) 1955: 83 - NPO (JP: HN, RY; KR)  
*B. nakanei* (Sawada) 1955: 85 - NPO (JP: RY; KR)  
*B. neoguineensis* Pace 2000a: 115 - Bismarck Sea (PG: Laing Island)  
*B. orousseti* Pace 1990: 66 - Luzon Sea (PH: Mindoro)  
*B. papuensis* Haghebaert 1995: 31 - Bismarck Sea (PG: Cape Vogel Peninsula)  
*B. parvula* Haghebaert 1995: 27 - Bismarck Sea (PG: Laing Island)  
*B. perexilis* Pace 1994: 132 - INO (SO: Sar Uanle)  
*B. peyerimhoffi* (Fauvel) 1904a: 73 Red Sea, Gulf of Akkaba, Mediterranean Sea (IL)  
*B. rothi* Moore and Legner 1975: 110 - Gulf of California (MX: SO)  
*B. sakishimana* Sawada 1991: 144 - NPO (JP: RY)  
*B. samoensis* Pace 1984: 67 - SPO (WS: Upolu Island)  
*B. sawadai* Moore, Legner and Chan 1973: 75 - South China Sea (CN: Hong Kong)  
*B. sinensis* Moore, Legner and Chan 1973: 76 - South China Sea (CN: Hong Kong)  
*B. subtilissima* (Cameron) 1904: 157 Red Sea [YE: Barim (= Perim) Island; SO: Hartan Peninsula]  
*B. testacea* (Cameron) 1904: 157 - Red Sea [YE: Barim (= Perim) Island]  
*B. testaceipennis* (Cameron) 1919: 245 - South China Sea (SG)  
*B. tsutsuii* (Sawada) 1955: 84 - NPO (JP: HN, RY)  
     = *B. serpentis* (Sawada) 1955: 84  
*Corallia* Fauvel 1878a: 212  
     *C. polyporum* Fauvel 1878a: 213 Arafura Sea [ID: Kepulauan Aru (Aru and Wokam), Kepulauan Kai (= Kei or Ke Island)]  
*Lautaea* Sawada 1989a: 83  
     *L. murphyi* Sawada 1989a: 85 - South China Sea (SG)  
*Myllaena* Erichson 1837: 382  
     *M. insipiens* Casey 1911: 237 - NAO, Gulf of Mexico (US: AL, FL, LA, NJ, PA)  
*Polypea* Fauvel 1878a: 301  
     *P. coralli* Fauvel 1878a: 302 - Arafura Sea (ID: Kepulauan Aru)  
*Rothium* Moore and Legner 1977: 460 [rev Ahn and Ashe 1996c]  
     *R. ashlocki* Ahn and Ashe 1996c: 247 - SPO (EC: Galapagos)  
     *R. evansi* Ahn and Ashe 1996c: 248 - SPO (EC: Esmeraldas, Guayas; PE: Piura)  
     *R. giulianii* Moore 1978b: 155 - NPO (MX: GU, SI)  
     *R. littoralis* Klimaszewski and Peck 1998: 228 - SPO (EC: Galapagos)  
     *R. pallidus* Ahn and Ashe 1996c: 247 - NPO (MX: GU)  
     *R. sonorensis* Moore and Legner 1977: 462 - Gulf of California (MX: SO)

## OXYPODINI

*Chilodera* Cameron 1944b: 619

*C. falklandica* Cameron 1944b: 620 - SAO (Falkland Islands)

*Dasydera* Cameron 1948: 731

= *Calonotus* Cameron 1945c: 171 [preoccupied]

= *Mecrona* Blackwelder 1952: 232

*D. algophila* (Broun) 1886: 941 - SPO (NZ: Mokohinau Island)

*Gyronotus* Cameron 1948: 731

= *Eurynotus* Cameron 1945c: 170 [preoccupied]

= *Marecon* Blackwelder 1952: 230

*G. rufipennis* (Broun) 1880: 92 - SPO (NZ: North Island)

*Oreuryalea* Assing and Maruyama 2002: 210

*O. watanabei* Assing and Maruyama 2002: 217 - NPO, East Sea (RU: Primorie, Sakhalin; JP: HK)

## PHYTOSINI

*Actocharis* Sharp 1870: 279

*A. readingii* Sharp 1870: 279 - NAO (FR; GB: England), Mediterranean Sea (DZ; FR: Corsica; HR; IT: mainland, Sardinia, Sicily; MT)

= *marina* Fauvel 1871: 159

*A. cassandrensis* Assing 1992: 45 - Mediterranean Sea (GR)

*Arena* Fauvel 1862c: 292

*A. fultoni* Cameron 1945c: 162 - SPO (NZ)

*A. tabida* (Kiesenwetter) 1850: 219 - NAO (FR; GB: England), North Sea (DE; DK; GB: England, Scotland; NL)

= *octavii* Fauvel 1862c: 292

*Euphytosus* Bernhauer and Scheerpeltz 1926: 552

[*Pseudophytosus* Haghebaert 1993: 161 is not a valid name]

= *Paraphytosus* Bernhauer 1922a: 236 [preoccupied]

*E. schenklengi* (Bernhauer) 1922a: 236 - South China Sea (TW)

*Phytosus* Curtis 1838: 718

*P. (Actosus) andalusiaensis* Haghebaert 1993: 161 Mediterranean Sea (ES: Andalusia)

*P. (Actosus) balticus* Kraatz 1859b: 52 - NAO (ES: Canary Islands; ES; FR; GB: England; IE; NO; PT), Irish Sea (GB: England), North Sea (BE; DE; GB: England, Scotland; NL), Baltic Sea (DE; DK; SE), Mediterranean Sea (DZ; FR; IT; MA; TN)

*P. (Actosus) holtzi* Bernhauer 1935: 48 - Mediterranean Sea (GR: Crete)

*P. (Actosus) nigriventris* (Chevrolat) 1843: 42 - NAO (ES; FR; GB: England; PT), Irish Sea (GB: England), North Sea (BE; NL), Mediterranean Sea (IT; MA)

= *minyops* Wollaston 1864: 531

*P. (Actosus) schatzmayri* Bernhauer 1941: 95 - NAO (PT: Azores)

*P. (s. str.) caribeanus* Haghebaert 1993: 163 - Caribbean Sea (GP)

*P. (s. str.) fenyesi* (Bernhauer) 1915a: 315 - NAO (SN)

= *senegalensis* Wendeler 1930: 252

*P. (s. str.) spinifer* Curtis 1838: 718 - NAO (ES: Canary Islands; FR; GB: England; IE; MA; PT), Mediterranean Sea (DZ; EG; ES; FR; GR; IT; MA; TN), Black Sea (BG; RO; TR), Baltic Sea (DE; DK; FI; SE), North Sea (BE; DE; DK; GB: England; NL)

= *dimidiatus* Wollaston 1865: 453

#### Incertae sedis

*Salinamexus* Moore and Legner 1977: 463 [rev Jeon and Ahn 2007]

= *Biophytosus* Moore and Legner 1977: 465

*S. browni* Moore and Legner 1977: 464 - Gulf of California (MX: SO)

*S. koreanus* Jeon and Ahn 2007: 193 - NPO (KR)

*S. reticulatus* (Moore and Legner) 1977: 466 - Gulf of California (MX: SO)

### OXYTELINAE

#### OXYTELINI

*Anotylus* Thomson 1859: 44

*A. maritimus* Thomson 1861: 131 - NAO (FR; GB: England, Scotland; NO), Baltic Sea (SE), North Sea (BE; DK; GB: England; NL), Mediterranean Sea (IT: Sicily; TN)

= *perrisii* Fauvel 1862a: xxiv

= *oceanus* Fauvel 1862c: 292

*Blediotrogus* Sharp 1900: 234

*B. cordicollis* (Broun) 1907: 57 - SPO (NZ: Chatham Islands)

*B. cribricollis* Fauvel 1900: 184 - SPO (NZ)

*B. fauveli* (Bernhauer and Schubert) 1911: 129 - SPO (AU)

*B. guttiger* Sharp 1900: 234 - SPO (NZ)

*Pareiobledius* Bernhauer 1934: 495

*P. alutellus* (Bernhauer) 1934: 495 - SAO (ZA)

*P. madegassa* Scheerpeltz 1969: 127 - INO (MG)

*P. pruinus* (Bernhauer) 1912a: 178 - SAO (ZA)

*Sartallus* Sharp 1871b: 217

*S. signatus* Sharp 1871b: 217 - SPO (AU: South Australia)

#### THINOBIINI

*Bledius* Leach 1819: 174

*aequatorialis* gp

*B. aequatorialis* Mutchler 1925: 225 - SPO (EC: mainland, Galapagos Islands)

*B. ceratus* Blackwelder 1943: 118 - Caribbean Sea (BS; CU; DO; HT; JM; US: FL)

*B. susae* Herman 1983: 98 - Gulf of Mexico (US: TX, South Padre Island)

## armatus gp

*B. fenyesi* Bernhauer and Schubert 1911: 129 - NPO (US: CA; MX: BN, BS)  
= *lecontei* Bernhauer 1905: 14 [preoccupied]

*B. monstratus* Casey 1889b: 46 - NPO (CA: BC; US: OR, CA)

## basalis gp

*B. cordatus* (Say) 1834: 461 - NAO and Gulf of Mexico (US: NY-FL, MS, TX)

*B. doderoi* Bondroit 1912: 66 - Mediterranean Sea (GR; IT)

*B. fergussoni* Joy 1912: 44 NAO (GB: England, N. Ireland, Scotland; IE),  
Baltic Sea (DE; EE; FI; PL; SE; RU: Karelia), North Sea (BE; FR; GB:  
England, Scotland; NO), Irish Sea (GB: England, Wales), Black Sea (RO),  
Mediterranean Sea (FR; MA; TN)

*B. gradensis* Bernhauer 1929: 183 - Mediterranean Sea (IT)

*B. minor* Mulsant and Rey 1878: 634 - Mediterranean Sea (AL; FR; IT)

*B. neglectus* Casey 1889b: 69 - NAO (CA: NL, NB, NS; US: ME-GA)

*B. subniger* Schneider 1898: 62 NAO (GB: England, Scotland; IE), Irish Sea  
(GB: England, Wales, IE), North Sea (DE; GB: England, Scotland; NL),  
Mediterranean Sea (ES; TN)

*B. thinopus* Herman 1976: 86 - Gulf of Mexico (US: FL, AL, TX)

*B. turbulentus* Casey 1889b: 70 - Gulf of Mexico (MX: QR, YU; US: FL, MS)

## bonariensis gp

*B. bonariensis* Bernhauer 1912a: 168 - SAO (AR; BR; UY)

## forcipatus gp

*B. actitus* (Herman) 1972: 127 - Gulf of Mexico (US: TX)

*B. litoreus* (Herman) 1972: 129 - Gulf of Mexico (US: FL)

## fratellus gp

*B. fratellus* Eppelsheim 1885: 144 - SAO (GH; NG; SN)

## furcatus gp

*B. maritimus* Bernhauer 1923: 176 - Red Sea (SU)

## gigantulus gp

*B. marinus* Bernhauer 1922b: 168 - INO (SC: Aldabra Islands)

*B. philippinus* Bernhauer 1912d: 248 - South China Sea (PH: Luzon)

*B. yezoensis* Nakane 1963: 21 - NPO (JP; KR)

## infans gp

*B. helferi* Fauvel 1904b: 112 - INO (IN; MM)

*B. infans* Rottenberg 1870: 36 - Mediterranean Sea (DZ; IT; LY; TN), Red  
Sea (YE)

*B. renominatus* Cameron 1914: 203 - INO (ET; SO)

= *bernhaueri* Cameron 1912: 28 [preoccupied]

## lamelliceps gp

*B. hasticeps* Bernhauer 1937: 583 - INO (MG; TZ)

## pulchellus gp

*B. pulchellus* Kraatz 1859a: 169 - INO (IN; LK; Chagos Archipelago)

## punctatissimus gp

- B. albomarginatus* Bernhauer 1922a: 225 - South China Sea (TW)  
*B. amplicollis* Fauvel 1900: 185 - SPO (NZ)  
*B. bidentifrons* Broun 1912: 401 - SPO (NZ)  
*B. buehleri* Scheerpeltz 1957b: 226 - Timor Sea (ID: Sumba)  
*B. buettikeri* Coiffait 1981a: 241 - Red Sea (SA)  
*B. capensis* Cameron 1945a: 708 - SAO (ZA)  
*B. caribbeanus* Blackwelder 1943: 113 - Caribbean Sea (JM; DO-TT)  
*B. caroli* Blackburn 1888: 13 - SPO (AU)  
*B. exiguus* Scheerpeltz 1933: 1114 - SPO (AU)  
     = *minor* Bernhauer 1920: 6 [preoccupied]  
*B. fernandezi* Bernhauer 1939: 234 - SAO (UY)  
*B. fossiventris* Fauvel 1889: 252 - SPO (NC)  
*B. injucundus* Blackburn 1888: 14 - SPO (AU)  
*B. maindroni* Fauvel 1903: 151 - INO (IN)  
*B. michaelsoni* Bernhauer 1915b: 313 - SAO (NA)  
*B. microcephalus* Fauvel 1901: 72 - NPO (CO), Caribbean Sea (TT: Trinidad)  
*B. orientalis* Bernhauer and Schubert 1911: 133 - Red Sea (DJ)  
     = *lividipes* Fairmaire 1892: 90 [preoccupied]  
*B. pontilis* Blackburn 1902: 22 - SPO (AU)  
*B. punctatissimus* LeConte 1877: 226 NAO (US: MDFL), Gulf of Mexico (US; MX: VC), Caribbean Sea (JM; PR), Gulf of California (MX), NPO (CO; EC)  
*B. salinus* Cameron 1947b: 704 - SPO (NZ)  
*B. tristis* Aubé 1843: 92 Mediterranean Sea (AL; DZ; ES; FR; IT; TN), Red Sea (SN)

## rugosicollis gp

- B. bituberculatus* Cameron 1940: 183 - Andaman Sea (MY)

## verres gp

- B. albopubescens* Cameron 1941: 434 - South China Sea (PH: Luzon)  
*B. arenicola* Fauvel 1904b: 112 - Arabian Sea [IN: Malabar, Mahé (= Mayyazhi)]  
*B. fraterculus* Cameron 1936: 40 - Andaman Sea (MY)  
*B. jacobsoni* Cameron 1928: 106 - INO (ID: Sumatra)  
*B. madagascariensis* Bernhauer 1901b: 169 - INO (MG)  
*B. marginalis* Cameron 1945a: 707 - SAO (ZA: Cape Province)  
*B. parens* Cameron 1941: 434 - South China Sea (PH: Luzon)  
*B. perrieri* Fauvel 1904c: 305 - INO (MG)  
*B. petzi* Bernhauer 1908b: 104 - INO (TZ)

*Carpelimus* Leach 1819: 174

- C. lucidus* (Cameron) 1944a: 312 - INO (TZ: Zanzibar)

*Teropalpus* Solier 1849: 330

- = *Trogolinus* Sharp 1900: 231  
*T. coloratus* (Sharp) 1900: 231 - SPO (NZ)



- T. lithocharinus* (LeConte) 1877: 245 - NPO (US: CA, WA)  
*T. luniger* (Fauvel) 1868: 40 - SPO (CL)  
*T. maritimus* (Broun) 1903: 615 - SPO (NZ)  
*T. pictipes* (Lea) 1910: 126 - SPO (AU: Tasmania)  
*T. senex* (Fauvel) 1868: 40 - SPO (CL)  
*T. skottsbergii* (Bernhauer) 1921: 41 - SPO (CL: Juan Fernandez Island)  
*T. suturalis* Solier 1849: 331 - SPO (CL)  
*T. unicolor* (Sharp) 1900: 232 - SPO (AU, NZ), INO (ZA), NAO (GB: England)  
     = *anglicanus* (Sharp) 1900: 232  
*Thinobius* Kiesenwetter 1844: 355  
     = *Yosiityphlus* Sawada 1971c: 327  
*T. frizzelli* Hatch 1957: 94 - NPO (CA: BC; US: CA, WA; MX: BN)  
*T. marinus* Cameron 1917d: 155 - South China Sea (SG)  
*T. kuroshio* (Sawada) 1971c: 327 - NPO (JP: HN)

## SCYDMAENINAE

### CEPHENNIINI

- Cephennodes* Reitter 1884: 420  
     = *Chelonoidum* Strand 1935: 285  
*C. araiorum* (Jałoszyński) 2003: 226 - NPO (JP: HN)

## PAEDERINAE

### PAEDERINI

- Chetocephalus* Cameron 1944a: 314  
*C. maritimus* Cameron 1944a: 314 - INO (MU)  
*Medon* Stephens 1833: 273  
*M. marinus* Cameron 1944a: 313 - INO (MU)  
*M. pocoferus* (Peyron) 1857: 718 - Mediterranean Sea (DZ; FR; IT; TN), NAO (FR; GB: England)  
     = *maritimus* Aubé 1863: 36  
*M. prolixus* (Sharp) 1874: 65 - NPO, East Sea (JP: HN)  
*M. rubeculus* Sharp 1889: 264 - NPO (JP: HN), South China Sea (CN: Hong Kong)  
*Ophioomma* Notman 1920: 704  
*O. rufa* Notman 1920: 705 - Gulf of Mexico (US: FL)  
*Sunius* Stephens 1829: 24  
*S. ferrugineus* (Bierig) 1934b: 326 - Caribbean Sea (CU; JM)  
*S. minutus* (Casey) 1905: 180 - NAO (US: FL)

## STAPHYLININAE

### STAPHYLININI

### PHILONTHINA

- Bisnius* Stephens 1829: 23



- B. macies* (Sharp) 1874: 41 - NPO (JP; KR)
- Cafius* Stephens 1829: 23
- C. aguayoi* Bierig 1934a: 66 - NAO (US: CT, MA)
- C. algarum* (Sharp) 1874: 38 - NPO, East Sea (JP: HN; KR), South China Sea (CN: Hong Kong)
- C. algophilus* Broun 1894: 419 - SPO (NZ)
- C. andamanensis* Coiffait 1981b: 337 - INO (Andaman Island)
- C. australis* (Redtenbacher) 1867: 28 - SPO (AU: New South Wales, Victoria)  
= *areolatus* Fauvel 1877: 251
- C. bistriatus* (Erichson) 1840: 502  
= *bilineatus* (Erichson) 1840: 503  
ssp. *bistriatus* (Erichson) 1840: 503 - NAO (CA: NB, NL, NS, QC; US: MA, MD, ME, NJ, NY, RI, VA, FL; BM, BS), Gulf of Mexico (MX: CA, VC; US: FL, TX), Caribbean Sea (AG; BB; CU; DO; GD; GP; JM; KN; LC; MS; PR; TT; VE; VI; MX: QR)  
ssp. *fulgens* Frank [in Frank et al. 1986]: 153 - NPO (US: CA; MX: BS), Gulf of California (MX: BN, BS, SO)
- C. bisulcatus* (Solier) 1849: 314 - SPO (CL)
- C. bryanti* Cameron 1943: 343 - SPO (AU)
- C. canescens* (Mäklin) 1852: 313 - NPO (US)
- C. caribeus* Bierig 1934a: 68 - Caribbean Sea (AG; CU; DM; GD; GP; JM; PR; VI; VE; US: FL)
- C. catenatus* Fauvel 1877: 256 - SPO (AU: New South Wales)  
= *velutinus* Fauvel 1877: 256
- C. caviceps* Broun 1886: 942- SPO (NZ)  
= *puncticeps* White 1846: 6
- C. ceylonicus* Bernhauer 1902: 29 - INO (LK)
- C. cicatricosus* (Erichson) 1840: 454 Mediterranean Sea (southern Europe; IT), NAO (FR; GB: England)  
= *sculticeps* (Motschulsky) 1858a: 649
- C. decipiens* (LeConte) 1863: 40 - NPO (US; MX)
- C. flicki* Vauloger 1897: 238 - Mediterranean Sea (IT; LY; TN)
- C. fonticola* (Erichson) 1840: 501 - Red Sea (EG), INO (SO)
- C. fucicola* Curtis 1830: pl. 323 - NAO (GB: England; IE), Irish Sea (GB: N. Ireland, Wales; IE), North Sea (FR; GB)
- C. gigas* Lea 1929: 204 - SPO (AU: Lord Howe Island)
- C. histrio* (Sharp) 1874: 37 - NPO, East Sea (CN: Hong Kong; JP: HN; KP; KR)
- C. lithocharinus* (LeConte) 1863: 38 - NPO (CA; US; MX)
- C. litoreus* (Broun) 1880: 108 - SPO (NZ)
- C. luteipennis* Horn 1884: 237 - NPO (CA; US; MX)
- C. maritimus* (Broun) 1880: 109 - SPO (NZ)

- C. martini* Cameron 1927: 251 - Red Sea (SA; YE)  
 = *arrowi* Bernhauer 1931b: 234
- C. mimulus* (Sharp) 1874: 38 - NPO, East Sea (JP: HN; KR)
- C. mutatus* Gemminger and Harold 1868: 590 - NPO (US; CA)  
 = *femoralis* (Mäklin) 1853: 189
- C. nasutus* Fauvel 1877: 257 [and 1879: 84] - SPO (FJ)
- C. nauticus* (Fairmaire) 1849: 288 - NPO (CN; JP; TW; US: HI), SPO (AU; FR: Tahiti; NC), INO (SO; LK; MU), Red Sea [YE: Barim (= Perim) Island]  
 = *longipennis* (Walker) 1858: 205  
 = *puncticollis* (Boheman) 1858: 31  
 = *parallelus* (Kraatz) 1859a: 99  
 = *densiventris* Fauvel 1877: 258
- C. opacus* (LeConte) 1863: 40 - NPO (western US; CA; MX)  
 = *dubius* (LeConte) 1863: 39
- C. pacificus* (Erichson) 1840: 501 - SPO (AU: New South Wales, Queensland, Tasmania, Victoria)  
 = *littoralis* Fauvel 1877: 254
- C. quadriimpressus* (White) 1846: 6 - SPO (NZ)  
 = *expuncticollis* Koch 1936: 180
- C. ragazzii* Gestro 1889: 32 - Red Sea, INO (SO)
- C. rufescens* Sharp 1889: 44 - NPO, East Sea (JP: HN; KR), South China Sea (CN: Hong Kong)
- C. rufifrons* Bierig 1934a: 68 - Caribbean Sea (CU), NAO (US: FL)
- C. sabulosus* Fauvel 1877: 253 - SPO (AU: New South Wales, Queensland)  
 = *postseriatulus* Koch 1936: 179
- C. seminitens* Horn 1884: 236 - NPO (CA; US; MX)
- C. seriatus* Fauvel 1877: 255 - SPO (AU)
- C. subtilis* Cameron 1922: 121 - Caribbean Sea (AG; CU; DM; GP; JM; KN; MS; PR; TT; VI), Gulf of Mexico (US: FL), NAO (US: FL; BM)
- C. sulcicollis* (LeConte) 1863: 40 - NPO (US; MX)
- C. velutinus* Fauvel 1877: 256 - SPO (AU)
- C. vestitus* (Sharp) 1874: 37 - NPO, East Sea (JP: HN; KP; KR)
- C. xantholoma* (Gravenhorst) 1806: 41 - NAO (ES: Canary Islands; FR; GB: England, Scotland; IE; IS; PT), Irish Sea (GB: England, Wales, N. Ireland; IE), North Sea (BE; DE; FR; GB: England, Scotland; NL; NO), Baltic Sea (DE; DK; EE; FI; PL; RU: Karelia; SE), Mediterranean Sea (DZ; EG; ES; FR; GR; IT; MA; TN; TR), Black Sea (UA)  
 = *lateralis* Stephens 1833: 246  
 = *littoralis* Stephens 1833: 247  
 = *tessellatus* Stephens 1833: 247  
 = *variegatus* (Erichson) 1840: 453  
 = *variolosus* (Sharp) 1871a: 181

- = *keysianus* Donisthorpe 1930: 97
- = *heroopoliticus* Koch 1936: 169
- C. zealandicus* Cameron 1947b: 705 - SPO (NZ)
- Gabronthus* Tottenham 1955: 178
  - G. maritimus* (Motschulsky) 1858a: 661 - NAO (ES: Canary Islands), Mediterranean Sea (CY; DZ; EG; FR; GR; IL; IT; LB; LY; MA; TR), Red Sea (DJ; ET; SA), INO (MU; RE), South China Sea (ID; MY; SG; VN), NPO (JP; TW)
  - = *mimulus* (Rottenberg) 1870: 30
- Orthidus* Mulsant and Rey 1876: 339
  - O. cribratus* (Erichson) 1840: 431 - Mediterranean Sea, NAO
    - ssp. *cribratus* (Erichson) 1840: 431 - Mediterranean Sea (IT)
    - ssp. *atlanticus* Coiffait 1956: 221 - NAO (ES; FR; PT; MA)
- Philonthus* Stephens 1829: 23
  - P. nudus* Sharp 1874: 36 - NPO (CA: BC; US: WA; RU: Kuril Islands), NPO, East Sea (KR; JP: HK, HN, KY)
- Phucobius* Sharp 1874: 35
  - P. africanus* Bernhauer 1937: 617 - INO (TZ: western Usambara)
  - P. congruus* (Walker) 1858: 205 - INO (LK)
    - = *punctilinea* (Walker) 1858: 205
    - = *horni* (Bernhauer) 1902: 28
  - P. cupreipennis* Cameron 1918: 89 - Java Sea (MY; SG)
  - P. densipennis* Bernhauer 1931a: 131 - NPO (JP: RY)
  - P. pectoralis* (Boheman) 1858: 31 - ?sea (CN: province not stated)
  - P. semiaereus* Cameron 1934: 22 - SPO (New Hebrides)
  - P. simulator* Sharp 1874: 35 - NPO, East Sea (KR; JP: HN, KY; RU: Primorie)
  - P. tricolor* Bernhauer 1917: 125 - South China Sea (CN: Hong Kong; TW)
- Remus* Holme 1837: 64
  - R. corallicola* (Fairmaire) 1849: 289 SPO (AU; FJ; NC), INO (LK; MG; MU; SC; SO), Red Sea (DJ; YE), South China Sea (CN: Hong Kong), Java Sea (ID; MY; SG)
    - = *occidentalis* Blackburn 1888: 48
  - R. filum* Kiesenwetter 1849: 19 Mediterranean Sea (IT; LY; YU), Black Sea, NAO (FR; DE; BG; RO; TR; EG; LY; HR), INO (SO)
  - R. pruinus* (Erichson) 1840: 510 - NAO (ES: Canary Islands; PT; NL; FR; RU), North Sea (BE; FR; NL), Mediterranean Sea (IT; TR)
  - R. sericeus* (Holme) 1837: 64 - NAO (ES: Canary Islands; FR; GB: England; PT: Madeira; NO), Baltic Sea (DE; DK; SE), North Sea (DE; DK; GB: England; NL), Irish Sea (GB: Wales), Mediterranean Sea (EG; ES; FR; GR; IT; LY; TR), Black Sea (BG), INO (MU), SPO (AU: South Australia, Tasmania, Victoria, Western Australia)
    - = *aegyptiacus* Motschulsky 1858a: 665
    - = *obscuricornis* Koch 1936: 170

*Thinocafius* Steel 1949: 309

*T. insularis* Steel 1949: 309 - SPO (NZ: Chatham Islands)

## QUEDIINA

*Heterothops* Stephens 1829: 23

*H. asperatus* Smetana 1971: 34 - NPO (CA: BC; US: CA)

*Quediocafus* Cameron 1945b: 791

*Q. hudsoni* Cameron 1945b: 791 - SPO (NZ)

*Q. insolitus* (Sharp) 1886: 379 - SPO (NZ)

*Q. taieriensis* (Broun) 1894: 424 - SPO (NZ)

## STAPHYLININA

*Hadropinus* Sharp 1889: 115

*H. fossor* Sharp 1889: 116 - NPO, East Sea (JP: HK; RU: Sakhalin)

*Hadrotes* Mäklin 1852: 313

*H. crassus* (Mannerheim) 1846: 509 - NPO (CA: BC; US: AK, CA, OR, WA; MX: BN)

*H. wakefieldi* Cameron 1945b: 786 - SPO (NZ)

*Liusus* Sharp 1889: 116

*L. hilleri* (Weise) 1877: 93 - NPO, East Sea (CN: Manchuria; JP: HN; KR; RU: Sakhalin)

*L. humeralis* (Matsumura) 1911: 113 - NPO, East Sea (CN; JP; KR; RU: Sakhalin)

*Thinopinus* LeConte 1852: 215

*T. pictus* LeConte 1852: 216 - NPO (CA: BC; US: AK, CA, OR, WA; MX: BN)

## Habits, habitats, and classificatory notes

### MICROSILPHINAE

*Microsilpha* includes four species although others are recognized but are undescribed. Three are South American and not coastal, but the one New Zealand species, *M. litorea* Broun, is known only from seashores (Klimaszewski and Watt 1997). Only this genus is included within the subfamily.

### OMALIINAE

Worldwide, there are about 117 genera of Omaliinae placed in seven tribes (Thayer 2005). Only the genera *Crymus*, *Omaliomimus*, *Macralymma*, and *Giulianium* seem entirely restricted to seashores. Some species within the genera *Micralymma* and *Oma-*

*lium* seem restricted to seashores. All of the coastal species thus far described occur at high latitudes and not in the tropics.

## APHAENOSTEMMINI

*Giulianium* includes three species, all of them found under debris below the high tide mark, of beaches of the North Pacific Ocean. Larvae are unknown and there is no known association with seaweed (Ahn and Ashe 1999).

## OMALIINI

*Crymus* (= *Arpediomimus*) includes two species, *C. antarcticus* and *C. kronii*, both associated with seaweed on seashores (Steel 1964). The larva of *C. kronii* was described by Steel (1964). Hughes et al. (2004) pointed out the existence of this species in the intertidal zone, not just on the South Island of New Zealand, but also on the Antipodes, Auckland, and Campbell islands of New Zealand, and found that it is one of the hosts of *Cucujomyces phycophilus* Weir and Rossi (Ascomycetes: Laboulbeniales).

*Macralymma* includes only one species, *M. punctiventre* Cameron, and it is pre-cinctive ("endemic") to New Zealand. Cameron (1945c) found the specimen in the Broun collection, labeled 'Taieri Beach.' It is widespread on the South Island of New Zealand and also occurs on Chatham Island. It is found under rotting kelp on sandy beaches (Emberson 1998). Hughes et al. (2004) pointed out the existence of this species in the intertidal zone of the Antipodes Islands of New Zealand and found that it is one of the hosts of *Cucujomyces phycophilus*.

*Micralymma marinum* adults and larvae inhabit cracks in rocks in the intertidal zone of rocky coastlines. Adults and larvae are predacious, but their prey range is uncertain (Thayer 1985), probably including Collembola. In Britain, adults overwinter and larvae develop during the summer months (Steel 1970), or adults and larvae overwinter (King et al. 1979). In the northeastern USA some of the beetles may overwinter as larvae (Thayer 1985). These apterous beetles tolerate immersion in seawater (Elliott et al. 1983). A second species, *M. brevilingue* Schiødte (1845: 377, syn. *M. dicksoni* Mäklin 1878: 24) has similar habits but is not entirely restricted to seashores, being also found in damp moss near coasts (Steel 1958) so we do not list it. A third species, *M. laticolle*, was admitted by Motschulsky (1860), its describer, not to belong to the genus *Micralymma*, but has not yet been assigned to another genus (see also Steel 1958). A fourth species, *M. caucasicum* Melichar, is not coastal (Steel 1962).

*Omalimimus* occurs only on seashores, and some species may be abundant in rotting seaweed (Steel 1964). The larva of *O. venator* was described by Steel (1964); Hughes et al. (2004) pointed out the existence of this species in the intertidal zone not just on the New Zealand mainland, but also on the Antipodes, Auckland, Campbell, Macquarie and Snares islands of New Zealand, and found that it is one of the hosts

of *Cucujomyces phycophilus* Weir and Rossi (Ascomycetes: Laboulbeniales). Emberson (1998) pointed out the existence of undescribed species of *Omalium* from the Chatham Islands of New Zealand.

*Omalium* species occupy various habitats, and only some occur on seashores associated with drifted seaweed. Seashore species include the European *O. laeviusculum*, *O. riparium*, and *O. rugulipenne*, as well as the North American *O. algarum*. Backlund (1945) found *O. laeviusculum* exclusively in deep layers of seaweed beds, but *O. riparium* was also found in carrion. Mjöberg (1906) described the pupa of *O. riparium*, and noted that it took seven days to develop to the adult. Steel (1970) found that larvae of *O. laeviusculum* and *O. riparium* occur in the summer months in Britain, whereas Larsson and Gigha (1959) had reported larvae of the latter in the winter in Iceland. Populations of *O. riparium* inhabiting Mediterranean shores have smaller adults than do those from northern Europe and have been considered a distinct subspecies (*O. riparium impar* Mulsant & Rey). *Omalium littorale* Kraatz seems to be strictly a seashore species in northern Europe, but in southern Europe it has been reported from high altitudes far from the sea (Zanetti 1987), so is not included in the checklist.

## PSELAPHINAE

Worldwide there are 1200 genera of Pselaphinae, and all are predacious (Thayer 2005). Seashores are a very minor part of their habitat range. The genus *Physoplectus* is known only from saline coastal habitats.

## BATRISITAE

### BATRISINI

*Arthromelus quadratus*, *Batriscenites celer*, *B. humicola*, and *Batriscenus foveiterminalis* adults were all found in mangrove forests in Singapore, where most of them were associated with mounds of *Thalassia* (Hydrocharitaceae) turtlegrass (Tanokuchi 1989).

## GONIACERITAE

### BRACHYGLUTINI

*Berlara bella* adults were collected in a mangrove forest in Singapore, but not in parts inundated by the tide (Tanokuchi 1989).

*Brachygluta* has at least 43 species, but just six of them, known from America north of Mexico, appear to be restricted to coastal habitats, all these on the Atlantic coast (Chandler 1997).

*Briaraxis depressa*, the only representative of this genus, is known only from “under rubbish or logs on the beach” in the circum-Caribbean regions (Chandler 1992, 2002).

*Mangalobythus furcifer*, *M. acutifolius*, and *M. murphyi* adults were all collected in mangrove forests in Singapore or Thailand, where adults were seen to be active on open ground at low tide (Tanokuchi 1989).

The genus *Nisaxis* appears to be entirely coastal. *Nisaxis maritima* is known only from coastal habitats in United States Gulf Coast States; other species have been found in coastal saline habitats and also in inland saline habitats (Chandler 1997).

*Pedisinops regulus* adults were collected in the intertidal zone and on a coral reef in Japan's Ryukyu Islands (Sawada 1991).

The genus *Physoplectus* is entirely coastal. *Physoplectus vinsoni* is known only from coastlines in Mauritius in the Indian Ocean, whereas *P. reikoeae*, *P. miyakei*, and *P. irritans* are from Pacific coastlines.

*Prosthecarthron sauteri*, originally described from Taiwan, was found to be widespread on patches of halophilous grasses close to the sea in North Korea, on patches of a reed on mud in river estuaries in the Japanese mainland, in mangrove habitats in the Ryukyu Islands, and under stones on the muddy ground of a mangrove seashore in Vietnam (Nomura et al. 2006).

## ALEOCHARINAE

Worldwide, over 1,151 genera of this subfamily have been described, but its true diversity is without doubt much greater (Thayer 2005). It now contains some 12,851 species, but Hammond (1975) postulated that it might contain as many as 100,000 species. The genera are currently distributed among 51 tribes whose relationships require much study. It contains many specialist seashore inhabitants, not only at the level of genus, but even (as currently defined) at the level of tribe. Nowhere else among the Staphylinidae have entire tribes specialized to inhabit coastal habitats.

Prepupae of Aleocharinae spin a silken cocoon in which they pupate; some earlier authors incorrectly supposed the dorsal abdominal gland to be the source of the silk, although that gland produces defensive chemicals; the cocoon is not a special adaptation to immersion in water (Frank and Thomas 1984a).

## ALEOCHARINI

*Aleochara*, the type genus of this tribe, appears to be the only one with coastal representatives, and thus far 16 are known. Its larvae develop as ectoparasitoids of cyclorrhaphous dipteran pupae within the dipteran puparium. On sea beaches, such dipteran puparia are typically found in piles of decaying seaweed, but also in carrion. All eight members of the subgenus *Emplenota* are found associated with such materials on sea beaches in Europe, north Africa, Korea, Japan, and both coasts of North America (Klimaszewski 1984). Additionally, *A. (Coprochara) sulcicollis* is also found in such habitats on the Pacific coast of North America and Chile in the south Pacific;



*A. (Coprochara) squalithorax* is found on the Pacific shores of Japan and Korea; and *A. (Coprochara) salsipotens* is found on the African shores of the Indian Ocean and the African shores of the south Atlantic. Other members of the subgenus *Coprochara* do not, or seldom, occupy beaches. Two species of the subgenus *Polystomota* (*A. punctatella* and *A. grisea*) occupy European shores of the North Atlantic. *Aleochara punctatella* has long been confused with *A. grisea*, so old records of *A. grisea*, doubtless including some in the checklist, are in doubt and need re-evaluation. Three members of the subgenus *Triochara* occupy Pacific shores of eastern Asia, no others are known, and the habitat of all three appears to be seaweed. Parasitoidism by *A. (Emplenota) obscurella* (as *A. algarum*) of a dipterous puparium [*Orygma luctuosum* Meigen (Diptera: Coelopidae)] was reported by Scott (1916), and then in greater detail by Scott (1920) who noted that hosts were *Coelopa pilipes* Haliday and *C. frigida* F. [as *Fucomyia gravis* Haliday (Diptera: Coelopidae)]. Lesne and Mercier (1922) were able to rear a 2<sup>nd</sup> instar of *A. obscurella* (as *A. algarum*) from *Coelopa* puparia and described and illustrated it. Paulian (1938b) reared *A. obscurella* (as *A. algarum*) from '*Fucella fucorum* Haliday', which we suspect referred to *Fucellia fucorum* Fallén (Diptera: Anthomyiidae), and added further descriptions and sketches of parts of the larvae; Paulian (1941) also described the larvae without specifying the host. Cals (1964) encountered *A. obscurella* (as *A. algarum*) as a parasitoid of *Coelopa frigida* and illustrated the pharate adult within its host puparium. Because parasitoidism is the only known way of life among larvae of at least 20 species of *Aleochara*, it is thought that all species have this habit (Peschke and Fuldner 1977; Klimaszewski 1984). Adult *Aleochara* are predacious.

## ATHETINI

*Acticola falkandica* is the sole representative of this genus. The type specimen was collected in December 1914 in seaweed at Port Stanley, Falkland Islands (Cameron 1944b).

*Adota* has three Nearctic representatives (*A. colpophila*, *A. gnypetoides*, and *A. maritima*) reported from seaweed stranded on Pacific shores, and three Palearctic species (*A. madida*, *A. magnipennis*, and *A. ushio*) from eastern Asia (Gusarov 2003b). In the British Isles, *A. maritima* was reported by Easton (1971, under the synonym of *Atheta immigrans*) as an adventive species. Thus all six species are known only as seashore inhabitants.

*Atheta*, at the time of the Coleopterorum Catalogus (Scheerpeltz 1934), was a generic name applied to many confused and disparate groups. Over the subsequent years monophyletic groups have been split off from it as distinct genera, but the task is not yet complete. The following six seashore species are still assigned to it, within several subgenera. *Atheta novaescotiae*, not assigned to a subgenus, is a salt-tolerant, coastal, beach-drift species known from Atlantic Canada (Klimaszewski et al. 2006). *Atheta (Actophylla) varendorffiana* is known from the North Sea coast of Germany. *Atheta (Badura) ririkoae* dwells on the coasts of Korea and Honshu, Japan, as does *A.*

(*Badura*) *tokiokai* (but this latter also on the coasts of Kyushu). *Atheta* (*Datomicra*) *acadiensis*, from Canadian Maritime Provinces, is typically found in dry beach-drift material at the top of the littoral zone, the material consisting of dead *Ascophyllum nodosum* (L.) and *Fucus vesiculosus* L. (Klimaszewski and Majka 2007). Finally, *A.* (*Sipalatheta*) *algarum* is found in seaweed on the coast of Hong Kong in the South China Sea (Pace 1999b).

*Brundinia* was initially described as *Homalota meridionalis* with “variety” *marina* by Mulsant and Rey (1853). The “variety” was later recognized as a valid species, and the generic name *Brundinia* was introduced later. This, or these, species (now *B. meridionalis* and *B. marina*) were initially reported from plant debris in a salt marsh at Hyères, on the Mediterranean coast of France. It or they (no others have been recognized) were later reported from similar habitats on the coasts of the North Atlantic, Irish, North, and Baltic seas. On the German shores of the Wadden Sea (separated by barrier islands from the North Sea), *B. marina* is a common species in the lower salt marsh, and it shows time-varying abundance within the elevational gradient (Irmeler and Heller 2002).

*Halobrecta* has about seven described species which in the past have been much confused by entomologists leading to much synonymy. The species occurring on Mediterranean shores need review (Gusarov 2004). For this reason, *H. halensis* Mulsant and Rey is here listed as a distinct species. The specimens of *Halobrecta flavipes* reported by Pace (2000b) from Chaiten, Chile belong to *H. algophila* (Gusarov 2004). The specimens of *Halobrecta flavipes* reported from Inaccessible Island in the South Atlantic Ocean by Klimaszewski et al. (2002) belong to *H. algophila* (Gusarov 2004). Adults of *Exatheta cingulata* and *E. consors* were described by Cameron (1920) and stated to have been collected in fungi, but Sawada (1985) transferred *E. cingulata* to *Halobrecta*, and Sawada (1987) synonymized *E. consors*. Because fungi are rarely found on seashores, and because all other species of *Halobrecta* occur on seashores, not in fungi, Cameron’s (1920) specimens may have been mislabeled. *Halobrecta discipula* was reported as a new species in Chile by Pace (1999a) who stated that it was found “in decaying vegetables (lettuces, onions)” near Valparaíso; if those vegetables had been dumped on a sea beach (not stated) near the port city of Valparaíso, they could have been used by the beetles as surrogates for drifted seaweed; see also comments under *Myrmecopora uvida*. Illustrations of the habitus and diagnostic structures of adult *H. flavipes* together with details about its habitat in New Brunswick are provided by Klimaszewski et al. (2008).

The synonymy of *Halobrecta flavipes* in the checklist follows Pope (1977) and Lott (2008) and differs from that given by Klimaszewski et al. (2002) following Bernhauer and Scheerpeltz (1926). Bernhauer and Scheerpeltz (1926) listed ‘*Aleochara elongatula* Stephens’ (1832) as a synonym of *H. flavipes*, but Stephens (1832) referred to *Aleochara elongatula* Gravenhorst which, if Stephens was correct in his identification, refers to what is now called *Atheta* (*Philhygra*) *elongatula* (Gravenhorst) or *Philhygra elongatula*, a species which is not closely related. Furthermore, the habitat specified by Stephens (1832) for this species [“not common: found occasion-

ally within the metropolitan district” (of London)] does not seem a likely habitat for a species of *Halobrecta*. If ‘*Aleochara elongatula* Stephens’ really is a synonym, it is the senior name and should be listed first among synonyms; however, this is likely to be a misidentification by Stephens. Also, according to Bernhauer and Scheerpeltz (1926), *Halobrecta atricilla* (Scriba) 1866: 290 *nec* Erichson is a synonym of *H. flavipes* and is so ranked in the checklist. For unexplained reasons, Bernhauer and Scheerpeltz (1926) did not accept Scriba’s (1866) redescription of *H. atricilla* (Erichson) as pertaining to that species.

*Hydrosmeeta subalgarum* is the only species assigned to this genus and is reported to inhabit seashores. Four specimens, collected under seaweed on sand at Tai Long, Hong Kong, China, are the only known collection of this species (Pace 1999b).

*Iotarphia* is monotypic. Its single species, *I. australis*, is known from adult specimens collected in a “maritime habitat” from Rockdale near Sydney and from Illawarra, both in New South Wales, Australia (Cameron 1943).

*Osakatheta* is monotypic. The single species, *O. yasukoae*, is known only from adults collected under stones on tidal flats at river mouths on the eastern coast of Honshu, Japan’s largest island, where their habitat is threatened by industrial development (Maruyama et al. 2008).

*Pontomalota* contains two species, *P. opaca* and *P. terminalis*, both of them known from the Pacific coast of North America. They inhabit the mid to upper littoral zone of fine-grained sandy beaches, covered by tides only once or twice each month and containing stranded seaweed. Adults appear to be most active at night or on heavily cloudy days. They may occasionally be very abundant with hundreds of individuals per m<sup>2</sup>. Whereas *P. terminalis* is known only from California, *P. opaca* is distributed from Alaska to Baja California and shows clinal variation in color with specimens black in the north to light brown in the south. Their immature stages are unknown (Ahn and Ashe 1992). Adults spend the daylight hours beneath driftwood or piles of stranded seaweed (Kincaid 1961b).

*Psammopora* contains only one species, *P. delittlei*; adults were detected on a sandy beach in Tasmania, Australia (Pace 2003).

*Psammotiba* is a genus containing five species, all from seashores in the northern Pacific, with two species in the Nearctic (*P. comparabilis* and *P. kenaii*) and three in the Palearctic (*P. hilleri*, *P. jessoensis*, and *P. kamtschatica*). Adults (the immature stages have not been reported) have been found in drifted seaweed (Gusarov 2003b).

*Tarphiota* has three species (*T. geniculata*, *T. fucicola*, and *T. densa*) confined to fine-grained sandy seashores of Pacific North America. They live in the mid- to upper-intertidal zone containing decaying seaweed and covered by only one or two high tides monthly. Larvae are unknown (Ahn 1996b, 1999).

*Thinusa* contains only two species (*T. fletcheri* and *T. maritima*) that live in the intertidal zone of sandy beaches immersed daily by tides on the Pacific coast of North America (Moore and Legner 1977; Topp and Ring 1988a; Ahn 1997b). Adults are active at night (Kincaid 1961b).

## DIGLOTTINI

Various genera have been removed to other tribes, leaving only the genera *Diglotta* and *Paradiglotta*.

*Diglotta* contains eight species, three (*D. mersa*, *D. sinuaticollis*, and *D. littoralis*) from the North Atlantic and adjacent seas (North, Irish, and Mediterranean), one (*D. brasiliensis*) from the South Atlantic, one (*D. secqi*) from the Red Sea, two (*D. legneri* and *D. pacifica*) from the North Pacific, and one (*D. maritima*) from the South Pacific. Lohse (1985), followed by Haghebaert (1991) confused the two European species of *Diglotta*; although this issue was resolved by Good (1998), distribution records of these two species are doubtless still confused. A larva, collected on the west coast of Denmark in June 1917 in association with adults of *D. mersa*, was described and illustrated by Kemner (1925). Of the two European species, *D. sinuaticollis* (as *D. mersa*) occupies the intertidal and supralittoral zone of sandy beaches, whereas *D. mersa* (as *D. submarina*) occurs mostly in salt and mud-marshes (Haghebaert 1991). The type locality of *D. brasiliensis* is the intertidal zone of an estuarine beach in the state of Paraná (Caron and Ribeiro-Costa 2008). The North American species occur on sandy beaches (Moore and Orth 1979a). The adults of the western Palaearctic and Brazilian species have 4 posterior tarsomeres whereas the North American and Pacific species have 5 tarsomeres (Haghebaert 1991; Caron and Ribeiro-Costa 2008; Klimaszewski et al. 2008). Illustrations of the habitus and diagnostic structures of adult *D. mersa* together with details about its habitat in New Brunswick are provided by Klimaszewski et al. (2008).

*Paradiglotta*, with its single species *P. nunni* Ashe and Ahn (2005) from New Zealand is not included in the present list because it has not been found on seashores, only inland.

## FALAGRIINI

Falagriini are a tribe of about 30 genera worldwide, only two of which have coastal representatives. The species of America north of Mexico were revised by Hoebeke (1985). A cladistic analysis of the genera of America north of Mexico by Ahn and Ashe (1995) showed monophyly of *Bryobiota* together with *Myrmecopora*, setting them apart from other genera in this tribe, which needs further study including more terrestrial falagriines.

The two species of *Bryobiota* (*B. bicolor* and *B. giulianii*) inhabit sandy beaches of the Pacific coast of North America. They live in the upper littoral zone, which is covered by tides only once or twice each month and contains buried decaying seaweed (Topp and Ring 1988a; Ahn and Ashe 1995). Larvae and diet are unreported.

*Myrmecopora* is currently divided into three subgenera (*Lamproxenusa*, *Paraxenusa*, and *Xenusa*). Fourteen species of *Myrmecopora* have been reported from seashores. Four species assigned to subgenus *Lamproxenusa* (*M. algarum*, *M. chinensis*, *M. reticulata*,

and *M. rufescens*) dwell on the shores of eastern Asia in seaweed on sandy beaches (Assing 1997b). One species assigned to *Paraxenus* (*M. laesa*) is known from the Mediterranean Sea and the Canary Islands of the north Atlantic. Ten species assigned to *Xenus* range from the shores of the Red, Black, and Mediterranean seas to the north Atlantic with its adjacent seas (Irish, North, and Baltic); one of these (*M. maritima*) is known only from Madeira and the Canary Islands, and one (*M. bernhaueri*) only from the Red Sea; additional species occupy other habitats. *Myrmecopora tenuicornis* (Küster) was treated as a synonym of *M. laesa* (Erichson) by Fauvel (1902) and subsequent authors. Assing (1997a) expressed misgivings about this synonymy but explained that the location of the type specimen of *M. tenuicornis* is unknown, so the synonymy cannot be resolved. *Myrmecopora (Xenus) uvida*, a species widespread in the western Palaearctic, was newly reported as an adventive species in Chile by Pace (1999a) who stated that it was found “under decaying vegetable products (cauliflowers, cabbages, onions, etc.)” near Antofagasta; if those vegetables had been dumped on a sea beach (not stated) near the port city of Antofagasta, they could have been used by the beetles as surrogates for drifted seaweed; see also comments under *Halobrecta discipula*.

## HOMALOTINI

Most genera and species of this tribe are not seashore inhabitants. Four genera, *Heterota*, *Paractocharis*, *Pseudopasilia* and *Thinobiosus*, are exclusively seashore species. Just one species of *Linoglossa* has been reported from coastal habitats. Two species of subgenus *Halmaeus* Kiesenwetter (1877) [= *Antarctophytosus* Enderlein (1909) = *Paraphytosus* Cameron (1917e) = *Austromalota* Brèthes (1925)] of the genus *Leptusa* Kraatz (1856), namely *L. atriceps* (C.O. Waterhouse 1875: 54 and 1879: 230) from South Georgia and Kerguelen Island, and *L. darwinii* (F.H. Waterhouse 1879: 531) from the Falkland Islands and South Georgia (= *rufomixtus* Brèthes 1925: 171) are omitted from the checklist because they are not strictly seashore species (Steel 1964).

*Cameronium* has two species in the Indian Ocean (*C. flavipenne* and *C. gomyi*), one in the Red Sea (*C. obockianus*), one in North Africa (*C. liebmanni* Scheerpeltz 1957a) and, remarkably, one in the Gulf of California (*C. sonorensis*). If the last is correctly assigned to genus, the dispersal of its ancestors to that locale would be surprising and not easily explicable; consequently, its placement should be reinvestigated. *Cameronium liebmanni* is known only from an inland freshwater lake in Algeria so we do not include it in the present compilation. In Somalia, *C. flavipenne* adults were trapped more abundantly in the wet season than in the dry season (Chelazzi et al. 1983).

*Heterota* is represented by ten species distributed from the East Sea (*H. sunjaei*), South China Sea (*H. arenaria*), and Bali Sea (*H. rougemonti*) through the Mascarene Islands of the Indian Ocean (*H. gomyi*, *H. obscura*, and *H. vinsoni*), the Red Sea (*H. brevicollis* and *H. pictipennis*), to the Mediterranean (*H. pamphylica* and *H. plumbea*). The distribution of the last of these extends from the Mediterranean to the west coasts of Europe and the Canary Islands, and more recently was discovered in southern and

northwestern Florida (USA), Jamaica, and the Caribbean coast of Mexico (Frank and Thomas 1984b). There is no apparent reason to believe that movement of *H. plumbea* from Europe to the Canary Islands and then to the Caribbean was assisted by humans. Small, salt-tolerant, winged insects are among the most likely candidates to disperse westward naturally with assistance of trade winds at tropical and subtropical latitudes. An annotated catalog of the species is provided by Park et al. (2008). On a sandy beach in Somalia, *H. pictipennis* adults were trapped more abundantly in the wet season than in the dry season (Chelazzi et al. 1983).

*Linoglossa murphyi* adults were collected in a mangrove forest in Singapore (Sawada 1991), but there is no evidence that other species of the genus are halobionts.

*Paractocharis* is represented by three species. The first described was *P. fucicola*, from sandy beaches under seaweed at Changi, Singapore, in the South China Sea (Cameron 1917c). The others, *P. deharvengi* and *P. orousseti*, were both found at Puerto Galera on Mindoro, Philippines, in the Luzon Sea, in "lavages" (washings) on the beach (Pace 1990).

*Pseudopasilia* is represented only by *P. testacea*, found on the coasts of the North Atlantic (England and France), North Sea (Belgium and southeastern England), and western Mediterranean (southern France, Italy including Sardinia, Tunisia, and probably Croatia). For many years entomologists confused *Arena tabida* with *P. testacea* (Tronquet 2003). Tronquet (2003) also noted adults found under small stones in the littoral zone.

*Thinobiosus salinus* is known only from the shores of Sonora in the Gulf of Mexico, where specimens were found in seaweed on the edge of a tidal pool (Moore and Legner 1977). No other species have been assigned to this genus.

## LIPAROCEPHALINI

All members of this tribe appear restricted to coastal habitats. It appears to be monophyletic, with relationships of its seven genera suggested as ((*Baeostethus*, *Ianmoorea*) (*Paramblopusa* ((*Amblopusa*, *Halorhadinus*) (*Liparocephalus*, *Diaulota*)))) by Ahn et al. (2010). Five of its genera appear restricted to the north Pacific Ocean (with adjoining Gulf of California and East Sea), and two to the south Pacific.

*Amblopusa* contains five species, all in the North Pacific, two of them (*A. alaskana* and *A. brevipes*) in North America, two (*A. pacifica* and *A. hokkaidona*) on Hokkaido in northern Japan, and one (*A. magna*) in the Far East of Russia. All of them inhabit the mid-littoral zone. A late instar of *A. alaskana* was described in detail by Ahn and Ashe (1996a). *Amblopusa pacifica* adults were collected in wrack at Akeshi, eastern Hokkaido, in August 1990 (Sawada 1991). *Amblopusa magna* adults were collected in piles of drifted seaweed on fine sand beaches at Ryazonovka near Slavyanka, Russia in June 1993.

*Baeostethus* has one species (*B. chiltoni*) that occupies rocky intertidal areas in the Antarctic islands of New Zealand. Adults, as is typical of Liparocephalini, are flightless; a late instar was described by Steel (1964) and by Leschen et al. (2002).



*Diaulota densissima* adults were found to be plentiful along the rocky alga-covered shores of Departure Bay (British Columbia, Canada), and larvae were found and illustrated; adults are flightless (Saunders 1928). Adults and larvae of *D. vandykei*, and adults and a few suspected larvae of *D. densissima* were obtained from cracks of rocks and under algae at Moss Beach (San Mateo County, California, USA) in January; brief descriptions and an illustration of the apex of the larval abdomen of both was provided by Chamberlin and Ferris (1929). Moore (1956b) added some diagnostic notes on the structures of larvae of *D. densissima*, *D. fulviventris*, *D. vandykei*, and *D. harteri* collected from beaches in California and Baja California (Mexico) and provided a key to identify them. Meyerdirk (1969) studied a population of *D. fulviventris* at La Jolla (California, USA) and found adults and larvae in June, August, September, and December 1968 with the number of beetles averaging 0.70 per cm<sup>2</sup>, increasing to 1.1 per cm<sup>2</sup> in March 1969. The range of the population extended from 0.76 to 1.83 m above mean low tide, exclusively in association with the acorn barnacle *Chthamalus fissus* Darwin (Thoracica: Chthamalidae). The barnacle provided refuge from wave force and currents. Beetle activity was highest at low tide, and ceased when the substrate became saturated with sea water. Laboratory observations showed no evidence of a circadian or tidal rhythm. Pupae were formed in silk-like cocoons within empty barnacle tests and cracks in rocks. Adults respired under water by means of cutaneous respiration, expanding their abdomen to expose more membranous area (between sclerotized plates) when they were submerged.

*Halorhadinus* is represented by three species inhabiting the littoral zone of boulder shores and sandy beaches where seaweed is stranded in Korea and Japan. Maruyama and Hayashi (2009) discussed the habitat of *Halorhadinus* in detail. Their immature stages are unknown (Ahn 2001).

*Ianmoorea* is represented by a single species, *I. zealandica*, detected under fine gravels in the intertidal region of Breaker Bay, Wellington, New Zealand (Ahn 2004, 2006).

*Liparocephalus* is represented by four species. *Liparocephalus cordicollis* adults (misidentified as those of *L. brevipennis*) were found to be plentiful along the rocky alga-covered shores of Departure Bay (British Columbia, Canada), and larvae were found and illustrated. Adults are flightless. Placed on the surface of a dish containing water and algae, adults descended and moved among the plants under water. They were thought to carry a film of air trapped “in the hairs of the body.” Oviposition and pupation were not obtained in the laboratory. In the laboratory, larvae ate pupae of the chironomid *Telmatogeton* Schiner, and some ate dead larvae of that genus and of *Camptocladius* Wulp. Adults were not observed to feed. The disappearance of algae from the rocks during the summer may account for disappearance of the beetles and their subsequent reappearance during the winter. The foregoing account by Saunders (1928) is confusing in that it is drawn from observations on both *L. cordicollis* and *Diaulota densissima* and sometimes makes no distinction between the two. Topp and Ring (1988b) reinvestigated, and found *L. cordicollis* between 0.2 and 2.0 m above lowest spring tide on rocky shores of Vancouver Island (British Columbia, Canada).



Adults and larvae can withstand submergence in seawater for > 2 weeks at 10°C, and thus can withstand continuous inundation from one spring tide to the next, although submerged larvae cease feeding and growing and do not pupate. Eggs were deposited in rock crevices or between the thalli of algae. Prepupae of *L. cordicollis* spin silken cocoons incorporating plant fragments in which to pupate. *Liparocephalus cordicollis* adults, and a few larvae and pupae, were found at Moss Beach (San Mateo County, California, USA) in November and December (Chamberlin and Ferris 1929). The final instar was described and illustrated, and the pupa was stated to be formed within a cocoon. A brief redescription by Moore (1956b) of the larva failed to clarify which of the species (*L. brevipennis* or *L. cordicollis*) he was describing. *Liparocephalus tokunagai* adults and larvae were found abundantly on rocky shores between high and low tide-marks in the spring of the year. The adults were seen to devour amphipod crustaceans of the genus *Gammarus* Fabricius (Sakaguti 1944). The final instar of *L. cordicollis* was described and illustrated by Ahn (1997a) with diagnostic characters for larvae of *L. brevipennis* and *L. tokunagai* together with a key for identification of larvae.

*Paramblopusa* was named by Ahn and Ashe (1996a) as a new genus to contain *P. borealis*, transferred from *Amblopusa* and known from the Pacific coast of North America where specimens have been collected under rocks on tidal flats. Later, *P. eoa* was described from the Kuril Islands of Russia, on rocks below a cliff at Negodnaya Bay (Ahn and Maruyama 2000).

## MYLLAENINI

Concepts of the tribe Myllaenini have changed radically in the past few years, and genera such as *Bryothinusa* and *Rothium* have been transferred to it from other tribes.

*Brachypronomaesa esakii* adults were found on a coral reef five km off the coast of Ishigaki Island, one of the Ryukyu Islands of southern Japan. This reef is submerged for all but about two hours daily. Adults are apterous, and the food and immature stages are unknown (Esaki 1956). Later collections from Okinawa in March–May 2002 showed that the beetles occupied coral reefs having large populations of Collembola suggesting that the beetles are predatory on the springtails (Ahn et al. 2003). The second species assigned to this genus is *B. sawadai*, described from the Bay of St. Vincent in New Caledonia by Jarrige (1964) without further information on its habitat. The nomenclature is not totally straightforward because Jarrige (1964) equated *Thalassopora* [a genus he described (Jarrige 1959) to include *T. nosybiana* from Madagascar, and *T. marchemarchadi* (the type species) from what is now Vietnam] with *Brachypronomaesa* in a heading “*Brachypronomaesa* Sawada, 1956 (*Thalassopora* Jarr., 1959)”. Nowhere in the following text did he mention this was a newly proposed synonymy. However, the text included description of a new species, *B. sawadai*, together with mention of characters of “*B. Marche-Marchadi*” and “*B. Esakii*”. Jarrige’s (1964) intent was clearly to synonymize *Thalassopora* with *Brachypronomaesa*. We believe that Jarrige (1964), despite lack of discussion, made a formal synonymy of *Thalassopora* with *Brachypronomaesa* thereby

transferring both the species he had originally placed in *Thalassopora*. A revision of the four species (*B. esakii*, *B. marchemarchadi*, *B. nosybiana*, and *B. sawadai*) is necessary. The habitat of *Thalassopora* (now *Brachypronomaea*) *nosybiana* was described as in sand and rocks near the oceanographical station of Nosy Bé, Madagascar (Paulian 1959). Its final instar (recognized because it was with adults, not reared) was described and illustrated, its digestive tube being packed with unicellular algae (Paulian 1959).

*Bryothinusa*, with 30 described species, restricted to seashores and with no species from other habitats, has the greatest diversity of all genera of the coastal Aleocharinae. Twenty-four live on shores of the Pacific Ocean and its surrounding seas, two (*B. madecassa* and *B. perexilis*) on the shores of the Indian Ocean, and four (*B. cameroni*, *B. peyerimhoffi*, *B. subtilissima*, and *B. testacea*) on the shores of the Red Sea (one of these also found in the Mediterranean). None has been reported from the Atlantic Ocean. Revisions have been published by Pace (1986), Haghebaert (1995), and Ashe (2005). Adults of *B. sakishimana* were collected in a mangrove forest (Sawada 1991), and those of *B. chengae* at light (Ahn 1998). The larva of the Californian *B. catalinae* was described by Moore and Orth (1979b) who described its habitat as “a shallow reef which is exposed at low-water and under water at high tide ... largely a field of boulders two or three feet across with smaller stones and gravel in sand ... adults and larvae were found beneath and on the stones in an association with dense worm tubes, chitons, limpets, small abalones, flatworms, small crabs and brittle stars.” Four species (*B. gangjinensis*, *B. koreana*, *B. minuta*, and *B. nakanei*) have been found on Korean shores (Ahn and Jeon 2004).

Of the *Bryothinusa* species found in Hong Kong, the habitats were distinguished as follows by Moore and Legner (1971) and Moore et al. (1973): (*B. chani*): “At low water they wander on the surface of the mud flat and at high tide they are under seawater by holding tightly to rocks and dead shells”; (*B. sawadai*): “Found among rock crevices and shells of barnacles and oysters. When burrowing in the sand never more than 1 cm deep. Can be found throughout the year.”; (*B. sinensis*): “Found among rocks, wandering on sand or burrowing in the sand as deep as 36 cm. Seems to prefer a sandy beach. Can be found throughout the year.”; (*B. hongkongensis*): “Found wandering on sand or burrowing in sand as deep as 36 cm from mid-tidal zone to low tidal zone. Specimens were found from February to April” (Moore and Legner 1971; Moore et al. 1973). Descriptions do not suggest that members of the genus are alike in sharing one kind of habitat. The habitat of *B. fluenta* was even more unusual: it was “beneath the surface of a fresh water stream a short distance from the seashore...[it] seems to have invaded a fresh water habitat directly from salt water” (Moore and Legner 1975). Adults of four of the Hong Kong *Bryothinusa* species (*B. chani*, *B. hongkongensis*, *B. sawadai*, and *B. sinensis*) were apparently able to breathe by plastron respiration, to feed on decaying microcrustacea, to occupy the mid-intertidal zone (0.6–1.5 m above mean tide level), to spend their time mainly at 0–15 cm below the sand surface, to mate on the sand surface, to occupy sand of low organic content, and to be macropterous except for *B. chani* (Wong and Chan 1977). Larvae of these four remained below the sand surface. The larva of *B. koreana* was described by Jeon and Ahn (2009), making it only the second species of the genus so treated (with the larva identified by mitochondrial DNA).

*Corallis polyporum* adults were stated by Fauvel (1878a) to be “sous-marines” (submarine or aquatic in the sea) under coral polyps in March. They occur together with those of *Polypea*, both genera being monotypic (Fauvel 1878a).

*Lautaea murphyi* adults were collected under mangroves in intertidal mud flats. Their digestive tracts were filled with fragments of harpacticoid copepods (Sawada 1989a). The genus is monotypic.

*Myllaena* is a genus distributed worldwide whose members are typically found in freshwater habitats, such as on the banks of lakes and streams, and in swamps. However, the North American *M. insipiens* Casey seems to have been collected only at or very close to coasts of the Atlantic and Gulf of Mexico of the USA (Klimaszewski 1982a), and is provisionally included in this list. *Myllaena leleupi* Pace (1985) is excluded from the checklist because its type locality in the Galapagos Islands is 4 km from the coast, despite a later record in a coastal habitat (Klimaszewski and Peck 1998).

*Polypea corallis* adults were said to have been found in the ocean under coral polyps, in March, to occur with those of *Corallis*, and to be submarine (Fauvel 1878a). Klimaszewski (1982b) argued that *Polypea* seems to lack any structural modification for a submarine existence.

*Rothium* was first included in Myllaenini by Ahn and Ashe (1996c). Its six described species inhabit the Pacific (including Gulf of California) coasts of North and South America. The first to be described, *R. sonorensis*, was collected in the intertidal zone “from algae covered pitted ryolite” and from “a tide pool” (Moore and Legner 1977). Discovery of two further Mexican species, and three from parts of Ecuador (including the Galapagos Islands) and Peru yielded little more information about their habitats. The larva of *M. sonorensis* was described by Moore (1977) and subsequently redescribed in detail by Ahn and Ashe (1996c).

## OXYPODINI

Seashore representatives of this large tribe belong to seem to be only four genera, all monotypic: *Chilodera*, *Dasydera*, *Gyronotus*, and *Oreuryalea* (Assing and Maruyama 2002).

*Chilodera falklandica* is known only from seaweed at Port Stanley in the Falkland Islands in the South Atlantic, where specimens were collected in December 1914 (Cameron 1944b).

*Dasydera algophila* was found among seaweeds on Mokohinau Island, New Zealand (Broun 1886).

*Gyronotus rufipennis* is known only from North Island, New Zealand (Broun 1880).

*Oreuryalea watanabei* is known from the Russian Far East and northern Japan, where it was detected among drifted seaweed and other debris on beaches. Four female specimens collected in July and September each had a single mature egg in its ovaries (Assing and Maruyama 2002).

## PHYTOSINI

The relationships of the tribe are unclear, and it is not yet certain that the tribe is monophyletic. The four genera remaining in this tribe, after removal of others to Myllaenini etc., are *Actocharis*, *Arena*, *Euphytosus*, and *Phytosus*.

*Actocharis* contains only two European species, *A. readingii* and *A. cassandrensis*. Both occur on Mediterranean coasts, and the first also occurs on North Atlantic coasts of England and France.

*Arena* includes one European species (*A. tabida*) and another (*A. fultoni*) from New Zealand, a remarkable distribution if these placements are actually correct (*A. fultoni* needs re-evaluation). *Arena tabida* was for years confused by entomologists with *Pseudopasilia testacea* (Homalotini) but it has a narrower distribution on the Atlantic coast of northwestern France, with the coasts of England and Wales, Irish Sea coasts of England and Wales, and North Sea coasts of England, Scotland, Netherlands, Germany, and Denmark, and is associated with drifted *Fucus* (Tronquet 2003).

Bernhauer (1922a) described specimens collected by Hans Sauter at Alikang in Taiwan under the name *Phytosus* (*Paraphytosus*) *schenklingi*. No information about its habitat was provided. However, the name *Paraphytosus* Bernhauer was preoccupied (by *Paraphytosus* Cameron 1917e). In 1926, the Coleopterorum Catalogus designated *Euphytosus* Bernhauer and Scheerpeltz as a replacement name (through objective synonymy) for *Paraphytosus* Bernhauer, with *P. (Euphytosus) schenklingi* as its type and only species.

Haghebaert (1993) excluded *Euphytosus* from the genus *Phytosus* with the consequence that it was automatically raised to generic rank; in that same work, Haghebaert also suggested (he wrote that he planned a later publication on the subject) an unnecessary new name (*Pseudophytosus*) for *Euphytosus*, so *Pseudophytosus* is not a valid name.

*Phytosus* is currently divided into two subgenera. The typical subgenus has three species; *P. spinifer* in the North Atlantic and Baltic, Black, North, and Mediterranean seas; *P. fenyesei* on the Atlantic coast of Senegal; and *P. caribeanus* on the shores of Guadeloupe in the West Indies. All five members of subgenus *Actosus* dwell on coasts of the North Atlantic and/or adjacent seas (Mediterranean, Baltic, or North). The present checklist does not mention *Phytosus atriceps* C.O. Waterhouse (1875), from Kerguelen Island, redescribed and illustrated by him (1879). The species was made the type of a new genus, *Antarctophytosus* by Enderlein (1909). Unaware of that assignment, Cameron (1917b) reported its finding in seaweed on a sandy sea-shore in the Falkland Islands and made it the type of a new genus, *Paraphytosus*. Cameron (1917e) admitted that he had misidentified the specimens, and also that the generic name *Antarctophytosus* Enderlein had precedence over *Paraphytosus*. Steel (1964) considered the genera *Antarctophytosus* Enderlein (1909), *Paraphytosus* Cameron (1917b), and *Austromalota* Brèthes (1925) all to be synonyms of *Halmaeus* Kiesenwetter (1877). Steel (1964) also found that none of the species of *Halmaeus* is restricted to seashores.

## **Incertae sedis**

*Salinamexus* has two species (*S. browni* and *S. reticulatus*) that occur on the shores of the Gulf of California and one (*S. koreanus*) in Korea. Adults are able to fly. All three appear to occur exclusively under boulders or seaweed on seashores; their immature stages are unknown (Jeon and Ahn 2008). Ahn et al. (2010) showed that the genus was not a member of the Liparocephalini. Its phylogenetic position is uncertain.

## **OXYTELINAE**

Worldwide, there are 47 known genera in Oxytelinae (Thayer 2005).

### **OXYTELINI**

This tribe has about 14 genera worldwide (Newton et al. 2001), including *Anotylus*, *Blediotrogus*, *Pareiobledius*, and *Sartallus*.

*Anotylus* has about 350 species worldwide, most associated with forest leaf litter, decomposing organic matter such as dung and carrion, and in mammal or ant nests (Newton et al. 2001). However, the European *A. maritimus* appears to occupy only seashore habitats, where it lives among seaweed and other drifted debris. *Anotylus speculifrons* (Kraatz 1857: 862) is often found in coastal habitats in western Europe, but occurs also in eastern Europe far from seashores; consequently we do not include it.

*Blediotrogus* has about five known species in Australia, New Zealand, and the Chatham Islands, and they are frequently found on seashores, under moist high-tide beach wrack (Makranczy 2006).

*Pareiobledius* has three known species, from the Afrotropical region, whose characteristic habitat is on seashores, under kelp (Makranczy 2006).

*Sartallus* has only one species, the Australian *S. signatus*, with winged adults. That species “is common on our [?South Australian] sandy beaches, where it [the adult?] hides under the seaweed and rubbish and feeds chiefly upon dead barnacles” (Froggatt 1907).

### **THINOBIINI**

This is the largest oxyteline tribe with about 20 genera worldwide (Newton et al. 2001). *Carpelimus*, with several hundred species, worldwide, is one of the two largest genera. Most *Carpelimus* species occupy freshwater habitats such as the shores of rivers and lakes, and in swamps. Moore and Legner (1974b) described the larva and pupa of *C. debilis* (Casey 1889a: 374) and noted that it occurs often in stranded seaweed on the Pacific shores of North America, but occupies other habitats as well (so we do not

list it). On the shores of Zanzibar, *C. lucidus* was found in seaweed and there seems to be no other habitat information, so we list it (Cameron 1944a).

*Bledius*, with over 450 species worldwide, is the other large genus within the Thiniini. Herman (1986) classifies it according to species groups (an ordering we follow in our checklist) rather than formal subgenera. Only some *Bledius* species dwell on seashores, and some of these also occur inland in saline habitats. We attempt to list only those species that do not occur also in inland saline habitats, resulting in the inclusion of 57 species names. We refer the reader to Herman (1986: 11–72) for an excellent overview of the natural history of the genus [including its predators and parasites (partially reviewed in Frank 1982, 1985b)]. Consequently herein we mention only a few other studies.

Several *Bledius* species found on coasts have been studied in some detail and have been noted as coastal species. However, evidence suggests that these are not strictly coastal, so they are not included in the present checklist. We mention them in part lest the reader should think we overlooked them and in part because the references cited provide larval and/or behavioral descriptions that may prove useful in comparison with future larval descriptions of coastal species. Four *Bledius* species [*B. tricornis* (Herbst), *B. hinnulus* Erichson), *B. gallicus* Gravenhorst] as *B. fracticornis* (Paykull), and *B. palipes* (Gravenhorst)] from Denmark have fine larval illustrations by Schiødte (1864). Five *Bledius* species [*B. arenarius* (Paykull), *B. talpa* (Gyllenhal), *B. subterraneus* (Erichson), *B. fuscipes* (Rey), and *B. opacus* (Block)] from Finland have fine larval descriptions and illustrations by Krogerus (1925). Four *Bledius* species [*B. spectabilis* Kraatz, *B. unicornis* (Germar), *B. furcatus* (Olivier), and *B. fuscicornis* Cameron] from France have larval descriptions by Paulian (1938a, 1941). The larva of *B. albonotatus* Mäklin from California was described (as *B. ornatus* LeConte) by Moore and Legner (1974a). Notes on behavior and development of *B. spectabilis* were published by Paulian (1942). Detailed studies by Bro Larsen (1936, 1951, 1952) of beetles of some Danish salt marsh and dune habitats dealt with several *Bledius* species particularly *B. spectabilis*. More recently, the distribution, behavior, and physiology of *B. spectabilis* was studied in Norfolk, England (Evans et al. 1971; Wyatt 1982, 1986; Wyatt and Foster 1988). The latter show the remarkable development of subsocial behavior among *Bledius* in which adults care for their brood. They also illustrate the form of the tunnels in which the entrance is narrowed to a bottle-neck so that it is not easily flooded by the tide (and can be blocked by the adults within a few minutes) to trap oxygen inside. Included is information on oxygen consumption and the minute algae that are the food of adults and larvae.

*Bledius fenyesi* and *B. monstratus* dwell on the Pacific coasts of North America in decaying, sand-covered piles of seaweed (Herman 1986). Evans's (1980) claim that adults of *B. monstratus* are predacious requires verification given that other members of the genus feed on algae.

*Bledius subniger* dwells on the sandy shores of Ostvoorn (Netherlands) just below the high tide mark, tunneling within the topmost 5 cm of sand, and occurring at densities up to 500 per m<sup>2</sup>. It feeds on Chlorophyta, Cyanophyta, and Chrysophyta (dia-



toms). It is very active at the end of June and the beginning of July, with individuals walking on the sand surface, followed by swarming for mating or dispersal (Hollander and van Etten 1974; Hollander 1983). These authors also discuss the habits of *Bledius arenarius*, which dwells in dunes above the beach and is thus excluded from our list of seashore staphylinids.

Griffiths and Griffiths (1983) found that *Bledius punctatissimus* attained densities of up to 2,260 adults per m<sup>2</sup> on a sheltered marine beach at Pawley's Island (South Carolina, USA). Adults and larvae below the sand surface are regularly immersed by the tide. At low tide they emerged to form galleries just below the sand surface and fed on diatoms. This activity lasted as long as 11 hours at higher shoreline elevations or 7 hours at elevations nearer to the lower limits of its distribution. At other times, adults and larger larvae singly occupied individual deeper burrows, whereas some females occupied maternal burrows together with their eggs and small larvae in side galleries; such burrows retained air during immersion by the tide. Griffiths and Griffiths' (1983) account is accompanied by drawings of egg, larva, pupa, and adult.

*Teropalpus* contains nine species, all with seashore distribution (Newton et al. 2001). Just one (*T. lithocharinus*) is native to the northern hemisphere, the remaining eight to the southern hemisphere, but one of the latter (*T. unicolor*), native to New Zealand, Australia, and South Africa, is adventive in Britain and arrived there before 1900. Specimens are found on seashores under driftwood and algae (Makranczy 2006).

*Thinobius* includes nearly 100 species worldwide (Newton et al. 2001). Most appear to live on banks of streams and ponds and other freshwater habitats, but a few live on intertidal mudflats. Among the latter, are *T. frizzelli* on the Pacific coast of North America, *T. marinus* in Singapore, and *T. kuroshio* on Honshu, the largest island of Japan. Dense populations of *T. frizzelli* were reported from felted growth of an alga [*Lyngbya semiplana* (C. Agardh) J. Agardh (Oscillatoriaceae)] in the intertidal zone of Willapa Bay in southwestern Washington State, where the individuals retreated into burrows among the algae when the tide was high; adults are winged, but the wings are of reduced size; adults are thought to eat "detritus"; larvae received a cursory description from Kincaid (1961a). In contrast, *T. kuroshio* adults were found in decaying drifted seaweed on a pebbly beach (Sawada 1971c), and *T. marinus* adults were found at Changi, Singapore, on sandy beaches under seaweed (Cameron 1917d). The synonymy of *Yosiityphlus* with *Thinobius* was documented by Gusarov and Makranczy (2004).

## SCYDMAENINAE CEPHENNIINI

Worldwide, over 4600 species in some 82 genera of this subfamily have been recorded (O'Keefe 2005). *Cephennodes araiorum* was found under stones on a stony/ sandy beach on the Pacific coast of Honshu, central Japan. It is the only member of the genus known from seashores, the others having been found in leaf litter and rotten deciduous wood at inland sites (Jałoszyński 2003).



## PAEDERINAE

About 221 genera of Paederinae occur worldwide; the tribal classification is under revision by L.H. Herman, so subsequent classification may be changed, and the placement of coastal species in the tribe Paederini (below) is uncertain (Thayer 2005).

### PAEDERINI

About 200 genera of Paederini occur worldwide. The monotypic genus *Chetocephalus* is based on *C. maritimus*, which was found in seaweed on Mauritius (Cameron 1944a).

*Medon* contains about 350 species worldwide, few of them occurring on seashores. Two exceptions are *M. marinus*, which was found in seaweed on Mauritius, and *M. pocoferus* found on the shores of the Mediterranean Sea and the North Atlantic Ocean. Two others are *M. prolixus* and *M. rubeculus*, both initially described from Japan, the former first reported from seaweed at Iwosima and Amakusa (Sharp 1874), but the latter was subsequently detected in Hong Kong where it appears to be restricted to the seashore (Rougemont 2001).

*Sunius* has about 125 species worldwide, but few of these are typically found on seashores. These exceptions are *S. ferrugineus* from the Caribbean Sea and *Sunius minutus* from Florida. Both live in drifted seaweed, and they may represent only a single species.

*Ophioomma*, a genus only recently transferred to Paederini, has only one species, *O. rufa*. Specimens were initially collected on the Gulf of Mexico coast of Florida, by sifting debris on the beach of Charlotte Harbor.

## STAPHYLININAE

This subfamily contains 320 genera worldwide (Thayer 2005). Adults and larvae are predacious.

### STAPHYLININI

This tribe contains 200 genera worldwide (Newton et al. 2001). Those with coastal species are *Thinopinus*, *Hadrotes*, *Hadropinus*, *Liusus*, *Thinocafius*, *Cafius*, *Remus*, *Phucobius*, and *Orthidus*, which do not form a monophyletic group.

**PHILONTHINA** are a subtribe with five genera (*Thinocafius*, *Cafius*, *Phucobius*, *Remus*, and *Orthidus*) that appear to be restricted to seashores. Additionally one species each of *Bisnius*, *Gabronthus* and *Philonthus* are also confined to coastal habitats. The systematic position of *Thinocafius* is unclear and requires study in relation to the phylogeny of *Cafius*.

*Bisnius* is moderately large genus with global distribution. Only one species (*B. macies*) appears to typically be found in drifted seaweed. It was originally described from Japan and was recently detected in Korea (Cho 2008).

*Cafius*, in its early sense included also species now assigned to *Remus*. An attempt by Koch (1936) to divide it into subgenera was not completed for all species known at that time, and perhaps for that reason was not followed by some later authors. Later, Coiffait (1963, 1974) elevated one subgenus (*Remus*) to generic rank and proposed a new subgeneric name (*Suborthidus*) to include one species within the remaining species of *Cafius*. Coiffait's effort to subdivide *Cafius* applied only to European, North African and Middle Eastern species, and was similarly not followed by some later authors. A global revision of the genus is necessary. In the present study 44 species are listed within *Cafius* (sensu stricto), making this the most species-rich genus of coastal staphylinids. One seemingly unusual species (*C. splendoris* Last 1987) is reported from Mt. Amigwiwa 1000–2300 m in Papua New Guinea and is excluded from our list. It may be misassigned to genus. Four species are attributed to *Remus* (see below). The zoogeographic distribution of species is as follows: Pacific Ocean 29, Atlantic Ocean 8, Indian Ocean 5, Pacific plus Atlantic 1, Pacific plus Indian 1.

The larva of the European *C. xantholoma* was described by Paulian (1941). Larvae and pupae of the Pacific species, *C. canescens*, *C. lithocharinus*, *C. luteipennis*, and *C. seminitens*, were described by James et al. (1971). Subsequently Moore (1975) described the larva of *C. sulcicollis*, although Orth and Moore (1980) stated that it was actually the larva of *C. bistriatus*. More recently, Jeon and Ahn (2002, 2007) used DNA sequence data to associate field-collected larvae with identified adults, and were able thus to identify and describe larvae of *C. histrio*, *C. mimulus*, *C. rufescens*, *C. fucicola*, *C. nauticus*, and *C. vestitus* from the Pacific. The European *C. xantholoma* is the best investigated species. Backlund (1945) compared the attractiveness of sterile leaves of various flowering plants versus sterile *Fucus* fronds and found that 26 adult specimens chose *Fucus* versus only four that chose other leaves. *Cafius xantholoma* is predacious as adult and larva, and is stenotopic in deep layers of seaweed beds (Backlund 1945). Observations of *C. xantholoma* larvae throughout the year (Backlund 1945; Egglisshaw 1965) suggest that the species is multivoltine. Adults of *C. xantholoma* are highly resistant to wetting and may take flight directly from the water surface (Backlund 1944), and adults of *C. bistriatus* show the same ability to avoid wetting and drowning (Frank et al. 1986). On the Pacific coast, mass flights of beetles including *C. luteipennis* have been observed and generally are in a direction parallel to the shore (Leech and Moore 1971; Evans 1980). We are unaware of reports of brachyptery among *Cafius* species. The distribution of *Cafius* species along the coast of the United States from New Jersey south to Florida and the Gulf of Mexico is determined by the abundance of drifted algae (not drifted seagrasses or marshgrasses), and some algae, for example *Sargassum fluitans* Borgessen, appears not to provide suitable habitat (Frank et al. 1986). The wider distribution of species originally described from the West Indies is still not determined although Frank (1985a) reported *C. caribeanus* from the South American mainland, and *C. subtilis* from Florida (USA). There are also questions about the relationship between *C. subtilis* and *C. aguayoi* (described from Massachusetts, USA), and of these

to the European species now known as *Remus sericeus*. A report of *C. xantholoma* from Chile was later shown to be based on misidentification. Most species of *Cafius* are found on the Pacific Ocean shores, and only one species (*C. bistriatus*) occurs both there and on the Atlantic. It is believed to have colonized Atlantic coasts from the Pacific, and the two populations have been distinguished morphologically at the subspecific level (Frank et al. 1986). On a sandy beach in Somalia, adults of three *Cafius* species (*C. fonticola*, *C. nauticus*, and *C. raggazzii*), were trapped more abundantly in the dry season than in the wet, and were more active at night in the dry season than in the wet (Chelazzi et al. 1983).

*Gabronthus* was recognized as a genus distinct from *Philonthus* in 1955. It contains over 30 species; only one of these, *G. maritimus*, found in the Mediterranean, the Red Sea, the Indian Ocean, and the South China Sea, appears to be restricted to coastlines. A report of *G. maritimus* from Cuba is highly questionable and without voucher specimens so has not been accepted in this study.

*Orthidus* contains only one species (*O. cribratus*) found on the Mediterranean coasts of southern Europe and northern Africa, and Atlantic coasts from Brittany (France) south to Morocco. It dwells under rocks and in piles of drifted algae on seashores (Coiffait 1974).

*Philonthus nudus* was for many years misattributed to *Cafius*, perhaps because of its atypical (for the genus *Philonthus*) coastal habitat. Its populations are distributed on the coasts of Korea, Japan, and the Kuril Islands of Russia, to the Pacific coasts of Canada and the United States, typically in drifted seaweed. *Philonthus* is a genus of over 1,000 species.

*Phucobius* contains eight species, of which seven are found under seaweed on beaches. The eighth (*P. africanus*) was reported from a single specimen collected at 1600 m in western Usambara, Tanzania. The collection locality has been confused, or perhaps the species belongs to another genus. Six species inhabit shorelines of the western Pacific (including the East Sea, South China Sea, and Java Sea, but not farther north than Japan), and one is found the shores of the Indian Ocean. Sharp (1874) recorded *P. simulator* as common under seaweed at Amakusa and Iwosima, Japan.

*Remus* contains four species, all found in drifted seaweed on coastlines. One species (*R. sericeus*) occurs on coasts of the North Atlantic, Baltic, Irish, North, Mediterranean, and Black seas and, remarkably, in Australia. Another species (*R. filum*) is confined to the coasts of the Mediterranean and Black seas. Another species (*R. pruinus*) is known from coasts of the North Atlantic, North, and Mediterranean seas. Another species (*R. corallicola*) is known from the South Pacific, South China, and Java seas, the Indian Ocean, and the Red Sea. The larva and pupa of *R. sericeus* were described and illustrated by Paulian (1941). Bierig (1934a) did not record *R. pruinus* from Cuba as listed by Peck (2005). On a sandy beach in Somalia, adults of two *Remus* species (*R. corallicola* and *R. filum*), were trapped more abundantly in the dry season than in the wet (Chelazzi et al. 1983).

*Thinocafius* contains only one known species, *T. insularis*, which is known only from the Chatham Islands east of New Zealand.

**QUEDIINA** are a subtribe containing one genus (*Quediocafius*) that appears to be restricted to seashores. One or more members of two other genera (*Heterothops* and *Quedius*) may or may not be restricted to seashores.

*Heterothops* is a widespread genus. Only one species, the North American *H. asperatus*, appears to be confined to Pacific seashores. The European *H. binotatus* (Gravenhorst) is often but not exclusively found on seashores and so is not included.

*Quediocafus* is known only from New Zealand, and all three species appear confined to sea beaches. Marris (2000) mentioned that *Q. insolitus* was found "under tussock mats and among coastal vegetation".

*Quedius* is a large and widespread genus apparently without any coastal specialists. *Quedius simplicifrons* Fairmaire (1861), which has several synonyms and is widely distributed at and near the coasts of western Europe and adjacent islands, does not meet our definition of a strictly coastal species. *Quedius umbrinus* Erichson (1839a), which Mjöberg (1906) found to be the most abundant insect in drifted seaweed at Bohuslän in southwestern Sweden, is not included because it also occurs in inland habitats.

**STAPHYLININA** are a subtribe containing the strictly coastal genera *Thinopinus*, *Hadrotus*, *Hadropinus*, and *Liusus*, as well as genera that are not confined to seashores.

*Thinopinus pictus*, occupying sandy Pacific beaches of North America from Alaska south to Baja California, has adults that are mottled dark and pale ('melanic') on dark sand beaches, but pale on pale sand beaches (Malkin 1958). Malkin (1958) pointed out that adults are nocturnal, spending the day concealed, and was perplexed as to why there should be two color forms. On beaches near Goleta (California, USA), adults and larvae are present throughout the year; dissected females contained 2–3 eggs, which are laid singly beneath the sand and hatch in ~ 14 days (Craig 1970). These eggs are white and ~ 3.0 mm long and 2.2 mm wide. Larval development time was not determined, but larvae were observed at night running on the sand surface like the adults (Craig 1970). On the western shore of Vancouver Island (British Columbia, Canada), the main prey is *Orchestia californiana* (Brandt), an amphipod which spends the day in temporary burrows on the upper part of the beach and is ambushed at night by the adult beetles (Richards 1982). The activity pattern of the amphipod is adjusted to weather and tidal patterns, and on some nights it is active after midnight, whereas beetles are active soon after dark; thus, beetles often foraged when amphipods were inactive (Richards 1983). Beetles attacked 0.147 amphipods per minute and captured 9.1% of prey attacked (Richards 1984). Color photographs of a larva (15 mm long) and adults of the two color forms (16–18 mm long) were provided by Evans (1980).

*Hadrotus crassus* lives on the Pacific coasts of North America, and is found from Alaska south to Baja California. Its larva was described by Moore (1964b) and color photographs of adult (11–17 mm) and larva (14–16 mm) are provided by Evans (1980); adults and larva are nocturnal predators of crustaceans and insects. A species described from New Zealand as *Hadrotus wakefeldi* may belong to another, possibly undescribed, genus (Klimaszewski et al. 1996).

*Hadropinus fossor* is an old-world representative of the subtribe, inhabiting the shores of northern Japan and of Sakhalin Island (Russia). It makes burrows in sand under seaweed (Sharp 1889).

## Discussion

Many kinds of insects can be found, some of them dead or dying in sea drift on shorelines. Many are winged insects that have alighted on the sea surface and have subsequently been washed up on the shore. Non-coastal species may exploit such concentrations of food. The senior author observed a pile of lawn grass clippings on a beach in Guanacaste, Costa Rica, that produced an abundance of staphylinids, none of them coastal species. Thus, would-be collectors of coastal staphylinids may be misled by the mere presence of staphylinids on seashores — a knowledge of non-coastal genera is essential. Coastal species represent only about 0.7% all Staphylinidae, although almost 400 species of them are known. Some representative species are shown in Figures 6–8.

Aleocharinae are the most species rich of the eight subfamilies with coastal representatives (Fig. 9). The Pacific Ocean has many more species of coastal staphylinids than do the other oceans (Fig. 10). The United States has more species of coastal staphylinids than do eight other countries (Fig 11).

## Polyphyly of coastal Staphylinidae

Lawrence and Newton (2000) and Newton et al. (2001) recognized four lineages among the Staphylinidae, and included among them the Pselaphinae and Scydmaeninae (Fig. 1). All four lineages contain genera and species that are restricted to coastal habitats. Coastal genera and species are found in eight subfamilies which contain mainly non-coastal genera. No subfamily contains only coastal genera. It is not until the level of tribe that there are taxa—all of them in the Aleocharinae—that include exclusively coastal genera.


The evolution of coastal genera in the Staphylinidae is polyphyletic. They have arisen from non-coastal ancestors among eight subfamilies. Furthermore, many coastal species belong to genera that include non-coastal species. Thus, we should expect a diversity of structural, physiological, and behavioral adaptations among them.

## The Pacific ocean as the main evolutionary center

Selecting only those genera that are exclusively coastal and separating them into three groups according to their provenance, yields the following:


- Coastal genera from the Pacific only: 40
- From the Pacific and elsewhere: 14
- From anywhere except the Pacific: 10

The Pacific Ocean (including its various seas such as the East Sea and South China Sea) is clearly the major cradle of coastal Staphylinidae.




**Figure 6.** Habitus photographs. **A** *Bryobiota bicolor*, 2.8 mm **B** *Bryothinusa koreana*, 3.2 mm **C** *Diaulota aokii*, 2.6 mm **D** *Diglossa sinuaticollis*, 3.0 mm **E** *Heterota sunjaei*, 2.4 mm **F** *Liparocephalus cordicollis*, 4.3 mm **G** *Myrmecopora simillima*, 3.1 mm **H** *Oreuryalea watanabei*, 4.9 mm **I** *Phytosus balticus*, 2.7 mm **J** *Pontomalota opaca*, 3.6 mm **K** *Psammotibia hilleri*, 5.3 mm **L** *Tarphiota geniculata*, 2.3 mm.






**Figure 7.** Habitus photographs. **A** *Omalium laeviusculum*, 4.3 mm **B** *Giulianium alaskanum*, 2.6 mm **C** *Prosthecarthron sauteri*, 1.9 mm **D** *Bledius fenyesi*, 4.7 mm **E** *Medon prolixus*, 4.6 mm **F** *Bisnius macies*, 8.3 mm **G** *Orthidus cribratus*, 12.2 mm **H** *Philonthus nudus*, 8.1 mm **I** *Hadropinus fossor*, 21.8 mm **J** *Hadrotetes crassus*, 13.6 mm **K** *Liusus hilleri*, 15.4 mm **L** *Liusus humeralis*, 14.0 mm **M** *Phucobius simulator*, 9.8 mm **N** *Remus corallicola*, 5.5 mm **O** *Remus sericeus*, 6.8 mm.



**Figure 8.** Habitus photographs. **A** *Cafius australis*, 15.4 mm **B** *C. bistratus*, 7.7 mm **C** *C. bistrio*, 8.5 mm **D** *C. lithocharinus*, male, 9.6 mm **E** *C. lithocharinus*, female, 9.3 mm **F** *C. litoreus*, 13.9 mm **G** *C. luteipennis*, 9.2 mm **H** *C. mimulus*, 8.0 mm **I** *C. pacificus*, 9.7 mm **J** *C. quadriimpressus*, 17.9 mm **K** *C. rufescens*, 6.2 mm **L** *C. seminitens*, 13.6 mm **M** *C. sulcicollis*, 7.5 mm **N** *C. xantholoma*, 7.8 mm **O** *Thinnocafius insularis*, 14.3 mm.




**Figure 9.** The number of coastal Staphylinidae species within each subfamily.

Coastal staphylinids have arisen within all four major lineages of the family, and even within these lineages they are polyphyletic. The simplest explanation is that the Pacific is oldest ocean (having arisen from the ancient ocean called Panthalassa). An alternative explanation might be that the shores of the Pacific are longer than the shores of other oceans, and somehow the species/area relationship (which has generally been supported for the staphylinid fauna of land masses) would suggest greater diversity on Pacific shores. A second alternative might be that non-coastal staphylinid faunas in lands surrounding the Pacific are more diverse than elsewhere and somehow had greater genetic plasticity. A third alternative might be that somehow the Pacific coasts have or had a special abundance of suitable habitat.

The ten genera not represented in the Pacific are: European (*Actocharis*, *Brundinia*, *Pseudopasilia*, and *Phytosus*), far South Atlantic (*Chilodera* and *Acticola*), Caribbean (*Briaraxis*), Gulf of Mexico (*Ophioomma*), and South Atlantic and Indian Ocean (*Pareiobledius* and *Chetocephalus*). *Briaraxis*, *Chilodera*, *Acticola* and *Ophioomma* are monotypic and because there are few specimens with habitat information, it is not certain whether the latter three are truly coastal. Europe, with four genera in the North Atlantic Ocean (and/or Mediterranean, Black, North, Irish, and Baltic seas) falls second to the Pacific Ocean in terms of precinctive genera.

*Actocharis* and *Phytosus* are currently placed in the tribe Phytosini. This tribe includes two other genera that have representative species in the Pacific. Unfortunately, this tribe requires revision and its monophyly is uncertain. Therefore, it is not clear whether *Actocharis* and *Phytosus* evolved from coastal ancestors that immigrated from the Pacific, or whether they evolved from non-coastal European ancestors. The discovery of a species of *Phytosus* in the Caribbean is discussed later.




**Figure 10.** The number of coastal Staphylinidae species found in various oceans and seas.

Apart from whether *Chilodera* and *Acticola* are truly coastal, their attribution to the South Atlantic Ocean needs qualification because they are known only from the Falkland Islands. Could they have a yet undetected distribution elsewhere? Another coastal genus, *Crymus*, has two species, distributed in the South Atlantic Ocean (Falkland Islands and South Georgia) and South Pacific Ocean (the Campbell and Auckland Islands of New Zealand) thus having a sub-Antarctic distribution. None of these genera extends farther north. Antarctic seas do not seem to have provided a conduit for migration of less cold-hardy coastal staphylinids around Cape Horn or the Cape of Good Hope. Forty genera remain confined to the Pacific and have not been detected in the South Atlantic. Among the 14 genera that are known from the Pacific and elsewhere, only two (*Halobrecta* and *Diglossa*) have species in the South Atlantic: *H. algophila*, which may have arrived adventively via shipping activities, and *D. brasiliensis*.

Even the Arctic Ocean has a partially coastal species (*Micralymma brevilinque*) whose congener, *M. marina*, is truly coastal and extends south through the Atlantic Ocean, east to the North and Baltic seas, and west to New England and Atlantic Canada. A third species (*M. laticolle*), also from the Arctic Ocean, does not belong to *Micralymma* (see Habits, Habitats, and Classificatory Notes), and should be transferred to another genus. A fourth species, in Eurasia, is not coastal.

The foregoing discussion suggests that almost all coastal genera of Staphylinidae evolved from non-coastal ancestors on the shores of the Pacific Ocean. Some apparently dispersed by migrating to the shores of other oceans. Geological constraints are outlined below.

The primordial ocean surrounding Pangaea in the Permian was Panthalassa, generally accepted as the direct antecedent of the Pacific Ocean (Scotese 2010). The breakup



**Figure 11.** The number of coastal Staphylinidae species found in various countries.

of Pangaea into Laurasia and Gondwanaland began by formation of the Tethys Sea which was the antecedent of the Indian Ocean and the Red Sea. Earlier authors (1960s) included the Mediterranean Sea, but recent authors credit the Mediterranean as being a separate entity, perhaps part of the Neotethys ocean basin, formed ca. 200 mya. The separation of Laurasia continued during the Triassic with the beginning of the North Atlantic Ocean. In the Jurassic, the North Atlantic Ocean widened and the Tethys Sea began to narrow. Not until the Cretaceous did the South Atlantic Ocean began to widen. In the Eocene, the South Atlantic Ocean became connected to the South Pacific Ocean by opening of the Drake Passage between Antarctica and South America some 41 mya (Scher and Martin 2006). The Mediterranean Sea lost its connection to the Indian Ocean some 15 mya. About 7 mya, the Mediterranean lost its narrow connections to the North Atlantic, its waters evaporated and it became a salt desert. At 4.86–4.6 mya, the Strait of Gibraltar opened, and water poured in from the North Atlantic in an immense flood (Garcia-Castellanos et al. 2009). The North Atlantic Ocean lost a major connection to the Pacific Ocean by completion of the Central American isthmus 3.5–3.1 mya (Bartoli et al. 2005). This is the geological background against which the current distribution of coastal Staphylinidae may be understood. It suggests that the Tethys Sea (and its extension, the Neotethys ocean basin), could have offered an early conduit from the Pacific via the Mediterranean to the Atlantic until the connection between the Mediterranean and Indian Ocean was lost, some 15 mya. However, between 7 and 4.8 mya, the Mediterranean became a salt desert and surely lost those parts of its coastal fauna that could not tolerate extreme salinity and aridity; the coastal Staphylinidae that had dispersed to the Atlantic could have survived, but other populations would have become extinct within the Mediterranean. [This ac-

count would to some extent be altered if recent ideas of a closed Pacific Ocean in the upper Triassic to lower Jurassic (McCarthy 2003) become generally accepted, although those ideas may make current distribution of Pacific Staphylinidae easier to explain].

### **The width of the north Pacific ocean has not been a severe barrier**

Liparocephalini (Aleocharinae) include seven genera containing 24 species, and all species are coastal. Five genera inhabit the North Pacific Ocean only, and four of them have representatives on both sides of it (i.e., in Asia and in North America). However, only two of the 24 species, *Diaulota aokii* and *Paramblopusa borealis*, occur on the shores of both continents. This suggests that the width of the North Pacific Ocean has not been a severe barrier to dispersal, and that most dispersal occurred in the remote past, allowing subsequent speciation.

Ahn et al. (2010) showed that the distribution of Liparocephalini was congruent with geological history, including the splitting of Pangaea and the isolation of the Palearctic from the Nearctic. They hypothesized that the ancestor of the tribe was distributed along the Panthalassan Ocean and repeated dispersals occurred into the Nearctic from the Palearctic.

Although these insects, presumably with favorable currents and winds, may cross oceans, we must suspect that barriers are formed by intervening land masses. This suspicion is heightened by dearth of records from inland localities, with just a few from California's Salton Sea (which once was connected to the Gulf of California). Thus, the biogeography of coastal staphylinids differs greatly from that of non-coastal species.

### **Westward dispersal across the Atlantic to the Caribbean**

*Phytosus* (Aleocharinae: Phytosini) has five European species, the range of two of which includes the Canary Islands; their presence on these islands suggests an ability to disperse. Three other species are known: one from the Azores, one from Senegal, and one from Guadeloupe in the Caribbean. Their ancestors are postulated here to have dispersed from Europe in the remote past. The westward dispersal of the putative ancestor of *P. caribeanus* across the Atlantic at the latitude of the Caribbean could have been assisted by trade winds.

*Heterota* (Aleocharinae: Homalotini) contains three species in the Pacific, three in the Indian Ocean, two in the Red Sea, and one in the Mediterranean. A tenth species (*H. plumbea*) inhabits the Atlantic coasts of Europe, the Mediterranean, and the Canary Islands. It has also been found in the Caribbean (Jamaica and the Yucatan peninsula of Mexico) and Florida (Frank and Thomas 1984b). Small winged insects adapted to a seashore existence could be prime candidates for transoceanic dispersal by the prevailing winds such as from southern Europe or northern Africa to the Canary Islands and thence to the Caribbean.



## Northward dispersal from the Red sea to the Mediterranean

The Red Sea may have been a conduit for dispersal from the Indian Ocean to the Mediterranean. If such dispersal did occur, a subsidiary question is whether it occurred recently, after the building of the Suez Canal, whether it occurred long before that and east or west of the Sinai Peninsula, or whether it occurred in the remote past before sea connections between the Red and Mediterranean seas closed ~15 mya. In this context it is worth noting that there is no evidence for dispersal of coastal staphylinids from the Indian Ocean via the South Atlantic to the North Atlantic.

*Bryothinusa peyerimhoffi* (Aleocharinae: Myllaenini) is reported from the Red and Mediterranean seas.

*Heterota* (Aleocharinae: Homalotini) has three species in the Pacific, three in the Indian Ocean, two in the Red Sea, and one in the Mediterranean (and one, previously mentioned, distributed from the Mediterranean to the Caribbean). Current distribution suggests that the Red Sea may have been an ancient conduit to the Mediterranean.

*Cameronium* (Aleocharinae: Homalotini) contains three species in the Indian Ocean, one in the Red Sea, and one (*C. liebmanni*, not listed) at inland localities in Algeria, adjacent to the Mediterranean. Current distribution suggests that the Red Sea may have been an ancient conduit to the Mediterranean. Could *C. liebmanni* have found refuge at inland locations as the Mediterranean Sea became a salt desert ca. 7 mya? A species from the Gulf of California was attributed to this genus by Moore (1964a), but this assignment needs reassessment.

## Eastward dispersal from the Pacific to the Atlantic

*Cafius* subgenus *Euremus* contains 10 species in the Pacific and a few others elsewhere. Because *C. bistriatus* has one subspecies (*C. b. bistriatus*) on the Caribbean and Atlantic coasts of North America, and one subspecies (*C. b. fulgens*) in the Gulf of California and on the Pacific shores of California and Baja California, it has been posited that it and the ancestor of *C. rufifrons* arrived in the Caribbean from the Pacific (Frank et al. 1986). If Central America is now an insuperable barrier to such dispersal, then the date of dispersal was probably before the connection of Central America to South America 3.5–3.1 mya.

*Aleochara litoralis* (Aleocharinae: Aleocharini) occurs on the Atlantic coast of North America from Florida north to Newfoundland and Quebec, as well as on the Pacific coast from California north to Alaska. There has been no report of specimens from the Canadian Arctic that could explain a natural dispersal.

## The distribution of coastal species of non-coastal genera

Selecting the coastal species belonging to non-coastal genera and placing them into the same categories as in preceding list yields the following:

Coastal species of non-coastal genera from the Pacific only: 10 genera, 39 species  
From the Pacific and elsewhere: 3 genera, 4 species  
From anywhere except the Pacific: 13 genera, 35 species  
The genus *Bledius* is excluded from this compendium.

Here, it can be seen that the numbers of genera and species from the Pacific only are almost identical to those from anywhere except the Pacific. The species included are such as *Philonthus nudus* and *Heterothops asperatus* (Pacific Ocean), *Medon pociferus* (Mediterranean), *M. marinus* (Indian Ocean), *M. prolixus* (Japan), *Sunius ferrugineus* (Caribbean), and *S. minutus* (United States). They belong to genera that have many non-coastal species. Although they have evolved to occupy coastal habitats, their anatomical characters do not set them apart as belonging to genera that are distinct from the non-coastal ancestors. In other words, their adaptation to coastal habitats is probably more recent than in genera that consist only of coastal species.

### **The genus *Bledius* is a special case**

We do not include the genus *Bledius* in the above accounts because we believe it is a special case, with possibilities of overland dispersal during its evolution.

### **Effects caused by humans**

There can be few species that have not been affected by human activity, and coastal staphylinids are no exception. *Teropalpus unicolor* is native to New Zealand and now occurs adventively in Australia, South Africa, on the Pacific coast of North America, and the south and southwest coasts of Britain (Hammond 2000). *Adota maritima* (syn. *Atheta immigrans*) is known from the Pacific coast of North America and adventively from Britain (Easton 1971, Hammond 2000). The European *Halobrecta flavipes* is another likely candidate for a list of species dispersed by shipping because it has been detected on the coasts of Chile, Australia, and eastern Canada. Another European species recently detected on the Atlantic coast of Canada is *Diglossa mersa* (Klimaszewski et al. 2008).

The building of seawalls for construction of marinas and docks, and for protection of buildings from wave action, damages beaches and in some places may harm coastal staphylinid populations. The senior author remembers that drifted seaweed accumulated on a beach by the west end of a bridge at Ft. Pierce, Florida in the early 1970s and harbored *Cafius* species. Twenty years later, the area had become a marina with seawalls and without seaweed accumulations; very many undocumented changes are likely to have been caused by similar constructions elsewhere. Human enjoyment of beaches to some extent depends upon the cleanliness of the beaches, and humans have employed beach-sweeping machines to remove not only garbage but also 'un-

sightly' drifted marine algae, which are the habitat and basis for food chains for some coastal staphylinids (Frank et al. 1986). An unusual sign at a public beach at Santa Barbara, California in summer 2009 (seen by the senior author) explained that seaweed was not being cleared there because it was the habitat of invertebrate animals that serve as food for birds. Oil spills on beaches are surely harmful to staphylinid populations, although we have seen no documentation. The compaction of beaches by human activity (vehicular as well as crowds of people) may be detrimental to Staphylinidae. Recently much-discussed threats to coasts and oceans are a rise in sea level and acidification, but their potential effect on coastal staphylinid populations have not been discussed in print if at all.

## Summary

1. Not all Staphylinidae found on seashores are coastal.
2. Coastal Staphylinidae comprise <1% of all species of Staphylinidae.
3. Coastal Staphylinidae have arisen in eight of the subfamilies, are polyphyletic, and have diverse adaptations of structure, physiology, and behavior that allow them to exist in such habitats.
4. There are far more strictly coastal genera in the Pacific than elsewhere and the Pacific appears to be the most important evolutionary center for this group.
5. There are approximately as many coastal species (belonging to otherwise non-coastal genera) in the rest of the world as there are in the Pacific.

## Acknowledgments

When JHF began a literature search, he already had originals or copies of most of the cited publications in his personal library. In early summer 2001, he began to request interlibrary loans through the University of Florida libraries (thanks especially to Janice Kaehler) for items he did not have and were not held by the University of Florida libraries. He thanks several friends for providing copies of pages of articles and books that the Interlibrary Loan Service of the University of Florida was unable to provide: Don Chandler (University of New Hampshire, Durham), Lanna Cheng (Scripps Institution of Oceanography, La Jolla), Terry Erwin (Smithsonian Institution, Washington, DC), Peter Hodge (editor, *The Coleopterist*), Berit Pedersen (library of the Royal Entomological Society, London) and Lizbeth Gale (library of the Royal Botanic Gardens, Kew). He thanks Beverly Pope (library of the Division of Plant Industry, Florida Department of Agriculture and Consumer Services, Gainesville), for help in tracking correct citations. After 2001, compilation became very much easier for some subfamilies due to the publication of Herman's (2002) catalog, and because all of Zoological Record became searchable online. Meanwhile, KJA had accumulated references on the taxonomic groups that he had studied. He thanks S. Nomura (National Museum of

Nature and Science, Tokyo), H. Hoshina (Fukui University, Fukui) and P. Jałoszyński (Poland) for providing valuable references. He is also grateful to members of the Chungnam National University Insect Collection (Daejeon) for taking habitus photographs and preparing figure plates. M.-J. Jeon (National Institute of Biological Resources, Incheon) provided us with many excellent habitus photographs of *Cafius* and related genera. Comments by Jan Klimaszewski (Natural Resources Canada, Québec) and Don Chandler significantly improved the manuscript, as did comments by an anonymous reviewer. KJA dedicates this paper to the late Steve Ashe, who provided the inspiration of studying the systematics and evolution of the seashore inhabiting staphylinid beetles. This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2009-0073111).

## References

- Ahn KJ (1996a) A review of *Diaulota* Casey (Coleoptera: Staphylinidae: Aleocharinae), with description of a new species and known larvae. *Coleopterists Bulletin* 50: 270–290.
- Ahn KJ (1996b) Revision of the intertidal aleocharine genus *Tarphiota* (Coleoptera: Staphylinidae). *Entomological News* 107: 177–185.
- Ahn KJ (1997a) A review of *Liparocephalus* Mäklin (Coleoptera: Staphylinidae). *Pan-Pacific Entomologist* 73: 79–92.
- Ahn KJ (1997b) Revision and systematic position of the intertidal genus *Thinusa* Casey (Coleoptera: Staphylinidae: Aleocharinae). *Entomologica Scandinavica* 28: 75–81.
- Ahn KJ (1998) *Bryothinus chengae*, a new species of the intertidal genus *Bryothinus* Casey from the Caroline Islands. *Coleopterists Bulletin* 52: 333–338.
- Ahn KJ (1999) *Tarphiota densus* (Moore), a new combination and key to the species of the genus *Tarphiota* Casey (Coleoptera: Staphylinidae: Aleocharinae). *Journal of the Kansas Entomological Society* 71: 191–193.
- Ahn KJ (2001) Phylogenetic relationships of the intertidal genus *Halorhadinus* Sawada and key to the genera of the Liparocephalini (Coleoptera: Staphylinidae: Aleocharinae). *Insect Systematics and Evolution* 32: 123–132. doi:10.1163/187631201X00074
- Ahn KJ (2004) *Moorea zealandica* new genus and species from New Zealand with a discussion of its phylogenetic relationships (Coleoptera: Staphylinidae: Aleocharinae). *New Zealand Journal of Zoology* 31: 255–261. doi:10.1080/03014223.2004.9518378
- Ahn KJ (2006) Replacement name for the genus *Moorea* Ahn (Coleoptera: Staphylinidae). *Coleopterists Bulletin* 60: 36. doi:10.1649/878.1
- Ahn KJ, Ashe JS (1992) Revision of the intertidal aleocharine genus *Pontomalota* (Coleoptera: Staphylinidae) with a discussion of its phylogenetic relationships. *Entomologica Scandinavica* 23: 347–359.
- Ahn KJ, Ashe JS (1995) Systematic position of the intertidal genus *Bryobiota* Casey and a revised phylogeny of the falagrine genera of America north of Mexico (Coleoptera: Staphylinidae: Aleocharinae). *Annals of the Entomological Society of America* 88: 143–154.

- Ahn KJ, Ashe JS (1996a) Revision of the intertidal aleocharine genus *Amblopusa* Casey and description of the new genus *Paramblopusa* (Coleoptera: Staphylinidae). *Journal of the New York Entomological Society* 103: 138–154.
- Ahn KJ, Ashe JS (1996b) Phylogeny of the intertidal aleocharine tribe Liparocephalini (Coleoptera: Staphylinidae). *Systematic Entomology* 21: 99–114. doi:10.1046/j.1365-3113.1996.d01-8.x
- Ahn KJ, Ashe JS (1996c) A revision of *Rothium* Moore and Legner (Coleoptera: Staphylinidae: Aleocharinae) with a discussion of its phylogenetic relationships. *Journal of the Kansas Entomological Society* 69: 234–256.
- Ahn KJ, Ashe JS (1999) Two new species of *Giulianium* Moore from the Pacific coasts of Alaska and California (Coleoptera: Staphylinidae: Aleocharinae). *Pan-Pacific Entomologist* 75: 159–184.
- Ahn KJ, Ashe JS (2004) Phylogeny of the Myllaenini and related taxa (Coleoptera: Staphylinidae: Aleocharinae). *Cladistics* 20: 123–138. doi:10.1111/j.1096-0031.2004.00012.x
- Ahn KJ, Jeon MJ (2004) The Korean species of the genus *Bryothinusa* Casey with descriptions of two new species (Coleoptera: Staphylinidae: Aleocharinae). *Coleopterists Bulletin* 58: 29–35. doi:10.1649/590
- Ahn KJ, Jeon MJ, Branham MA (2010) Phylogeny, biogeography, and the stepwise evolutionary colonization of intertidal habitat in the Liparocephalini based on morphological and molecular characters (Coleoptera: Staphylinidae: Aleocharinae). *Cladistics* 26: 344–358.
- Ahn KJ, Maruyama M (2000) *Paramblopusa eoa*, a new intertidal species (Coleoptera: Staphylinidae: Aleocharinae) from Kuril Islands. *Coleopterists Bulletin* 54: 359–364. doi:10.1649/0010-065X(2000)054[0359:PEANIS]2.0.CO;2
- Ahn KJ, Maruyama M, Jeon MJ (2003) Redescription of the intertidal genus *Brachypronomaea* Sawada (Coleoptera: Staphylinidae: Aleocharinae) from the Ryukyu Islands, Japan with a discussion of its phylogenetic relationships. *Journal of the Kansas Entomological Society* 76: 622–629.
- Allen AA (1937) A new species of *Aleochara* (Col., Staphylinidae). *Entomologist's Monthly Magazine* 73: 218–220.
- Ashe JS (2005) A new species of the intertidal staphylinid genus *Bryothinusa* Casey from Malaysia, with an overview of geographic distribution and an associated catalog of *Bryothinusa* (Staphylinidae: Aleocharinae: Myllaenini). *Coleopterists Bulletin* 58 (2004): 581–597.
- Ashe JS, Ahn KJ (2005) *Paradiglotta nunni*, a remarkable new genus and species in the aleocharine tribe Diglottini (Coleoptera: Staphylinidae: Aleocharinae) from New Zealand. *Journal of the New York Entomological Society* 112 (2004): 111–120.
- Assing V (1992) A new species of the genus *Actocharis* Fauvel 1869 from Greece: *Actocharis cassandrensis* n. sp. (Col.; Staphylinidae). *Entomologische Blätter für Biologie und Systematik der Käfer* 88: 43–48.
- Assing V (1995) The Palaearctic species of *Emplenota* Casey, *Polystomota* Casey, *Triochara* Bernhauer and *Skenochara* Bernhauer & Scheerpeltz, with descriptions of three new species (Coleoptera, Staphylinidae, Aleocharinae). *Beiträge zur Entomologie* 45: 217–237.

- Assing V (1997a) A revision of the Western Palaearctic species of *Myrmecopora* Saulcy, 1864, sensu lato and *Eccoptyoglossa* Luze, 1904 (Coleoptera: Staphylinidae, Aleocharinae, Falagriini). *Beiträge zur Entomologie* 47: 69–151.
- Assing V (1997b) A revision of the Eastern Palaearctic species of *Myrmecopora* Saulcy, 1864 with notes on some species from the Oriental region (Coleoptera: Staphylinidae, Aleocharinae, Falagriini). *Beiträge zur Entomologie* 47: 337–352.
- Assing V, Maruyama M (2002) A new genus and species of intertidal Oxypodini (Coleoptera, Staphylinidae, Aleocharinae) from the eastern Palearctic Region. *Special Bulletin of the Japanese Society of Coleopterology*, Tokyo 5: 209–220.
- Aubé C (1833) *Pselaphiorum monographia cum synonymia extricata*. *Magasin de Zoologie* 3: 1–72, pls. 78–94.
- Aubé C (1843) Note sur deux espèces nouvelles de brachélytres. *Annales de la Société Entomologique de France* (sér. 2) 1: 89–92.
- Aubé C (1863) *Lithocharis maritima*. In: Allard E, Aubé C, Brisout de Barneville C, Chevrolat A, Fairmaire L, Fauvel A, Grenier A, Kraatz G, Linder J, Reiche L, de Saulcy F *Matériaux pour la faune française*: 36. In: Grenier A, *Catalogue des coléoptères de France et Matériaux pour servir à la faune des coléoptères français*. Grenier, Paris, 1–130. [note: these pages, 1–130, form a separately-paged appendix to the catalogue].
- Audisio P, Taglianti AV (2010) *Insecta Coleoptera*. *Biologia Marina Mediterranea* 17 (suppl. 1): 547–571.
- Backlund HO (1944) The ability of the shore beetle, *Cafius xantholoma* Grav. to evade seadrift. *Kungliga Fysiografiska Sällskapetets Lund Förhandlingar* 14: 14.
- Backlund HO (1945) Wrack fauna of Sweden and Finland. Ecology and chorology. *Opuscula Entomologica* (supplementum): 1–237, pls. 1–6.
- Barrett JH, Yonge CM (1958) *Collins pocket guide to the sea shore*. Collins, London, 272 pp. + 40 colored pls. + 40 black and white pls.
- Bartoli G, Sarnthein M, Weinelt M, Erlenkeuser H, Garbo-Schönberg D, Lee D (2005) Final closure of Panama and the onset of northern hemisphere glaciation. *Earth and Planetary Science Letters* 237: 33–44. doi:10.1016/j.epsl.2005.06.020
- Bernhauer M (1901a) Neunte Folge neuer Staphyliniden aus Europa, nebst Bemerkungen. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien* 50 (1900): 532–541. [note: volumes 8–67 of this journal include the letters k-k before the word Zoologisch—these are an abbreviation for kaiserlich-königlichen (imperial-royal)]
- Bernhauer M (1901b) Zur Staphylinidenfauna von Madagascar. *Deutsche Entomologische Zeitschrift* (1901): 161–176.
- Bernhauer M (1902) Zur Staphyliniden-Fauna von Ceylon. *Deutsche Entomologische Zeitschrift* (1902): 17–45.
- Bernhauer M (1905) Neue exotische Staphyliniden. *Deutsche Entomologische Zeitschrift* (1905): 9–21.
- Bernhauer M (1907) Zur Staphylinidenfauna von Japan. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien* 57: 371–414.
- Bernhauer M (1908a) Beiträge zur Kenntnis der paläarktischen Staphyliniden-Fauna. *Münchener Kolopterologische Zeitschrift* 3: 32–335.



- Bernhauer M (1908b) Neue Staphylinidae aus dem subtropischen und tropischen Afrika. In: Schultze L, Zoologische und anthropologische Ergebnisse einer Forschungsreise im westlichen und zentralen Südafrika. Vol. 1. Systematik und Tiergeographie. Denkschriften der Medicinisch-Naturwissenschaftlichen Gesellschaft zu Jena 13: 101–118.
- Bernhauer M (1910) Beitrag zur Staphylinidenfauna des palaearktischen Gebietes. Entomologische Blätter für Biologie und Systematik der Käfer 6: 256–260.
- Bernhauer M (1912a) Zur Staphylinidenfauna von Südamerika. Entomologische Blätter für Biologie und Systematik der Käfer 8: 167–179.
- Bernhauer M (1912b) Staphylinidae, 170–172. In: Baker CF, Some Coleoptera of the beach at Laguna with descriptions of new species by Dr. M. Bernhauer. Annual Report of the Laguna Marine Laboratory 1: 170–173.
- Bernhauer M (1912c) Beitrag zur Staphyliniden-fauna von Afrika (Col.). Entomologische Mitteilungen 1: 203–209.
- Bernhauer M (1912d) Neue Staphyliniden der Philippinen. Philippine Journal of Science 7: 245–254.
- Bernhauer M (1915a) Neue Staphyliniden des tropischen Afrika. Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien 65: 287–321.
- Bernhauer M (1915b) Coleoptera. VII. Staphylinidae. In: Michaelsen W (Ed) Beiträge zur Kenntnis der Land- und Süßwasserfauna Deutsch-Südwestafrikas. Ergebnisse der Hamburger deutsch-südwestafrikanischen Studienreise 1911, L. Friederichsen, Hamburg, 31–321.
- Bernhauer M (1917) Ein neuer *Phucobius* aus China. Wiener Entomologische Zeitung 36: 125.
- Bernhauer M (1920) Results of Dr. E. Mjöberg's Swedish scientific expeditions to Australia 1910–1913. 22. Staphylinidae. Arkiv för Zoologi 13 (8): 1–27.
- Bernhauer M (1921) Coleoptera-Staphylinidae von den Juan Fernandez-Inseln und der Osterinsel. In: Skottsberg C (Ed) Natural history of Juan Fernandez and Easter Island. Vol. 3. Zoology. Almqvist and Wiksells, Uppsala, 41–44.
- Bernhauer M (1922a) Sauter's Formosa-Ausbeute: Staphylinidae. 1. Teil. Archiv für Naturgeschichte (A) 88: 220–237.
- Bernhauer M (1922b) No. II. Coleoptera, Staphylinidae. In: Reports of the Percy Sladen Trust Expedition to the Indian Ocean in 1905, under the leadership of Mr. J. Stanley Gardiner, M. A. vol. 7. Transactions of the Linnean Society of London (ser. 2) Zoology 18: 165–186.
- Bernhauer M (1923) Staphylinidae. In: Ebner R, Wissenschaftliche Ergebnisse der mit Unterstützung der Akademie der Wissenschaften in Wien aus der Erbschaft treitl von F. Werner unternommen Zoologischen nach dem Anglo-Aegyptischen Sudan (Kordofan) 1914. IX. Coleoptera A.: 176–178. Akademie der Wissenschaften in Wien, Mathematisch-Naturwissenschaftliche Klasse, Denkschriften 98: 165–199.
- Bernhauer M (1927a) Zwei neue Kurzflügler aus Schweden. Entomologisk Tidskrift 57: 167–168.
- Bernhauer M (1927b) Neue Staphyliniden des paläarktischen Faunengebietes. Koleopterologische Rundschau 13: 90–99.
- Bernhauer M (1929) Neue Kurzflügler des paläarktischen Gebietes. Koleopterologische Rundschau 14: 177–195.

- Bernhauer M (1931a) Zur Staphylinidenfauna des chinesischen Reiches. Wiener Entomologische Zeitung 48: 125–132.
- Bernhauer M (1931b) Neuheiten der paläarktischen Staphylinidenfauna. Koleopterologische Rundschau 17: 232–245.
- Bernhauer M (1934) The staphylinid fauna of South Africa. Annals of the South African Museum 30 (4): 481–509, pl. xiv.
- Bernhauer M (1935) Neuheiten der paläarktischen Staphylinidenfauna. Koleopterologische Rundschau 21: 39–48.
- Bernhauer M (1937) Neuheiten der ostafrikanischen Staphylinidenfauna. Festschrift zum 60. Geburtstage von Professor Dr. Embrik Strand. [no publisher named on titlepage] Riga 2: 577–619.
- Bernhauer M (1939) Zur Staphylinidenfauna Argentinien und Brasiliens (Col.). Revista de Entomologia (Rio de Janeiro) 10: 231–249.
- Bernhauer M (1941) Eine neue Art der Gattung *Phytosus* Curt. (Staph.), mit einer Übersicht der Arten der Untergattung *Actosus* Muls. et Rey. Koleopterologische Rundschau 26: 95–96.
- Bernhauer M (1943) Neuheiten der paläarktischen Staphylinidenfauna. Mitteilungen der Münchener Entomologischen Gesellschaft 33: 169–188.
- Bernhauer M, Scheerpeltz O (1926) Staphylinidae VI. In: Schenkling S (Ed) Coleopterorum Catalogus. W. Junk, Berlin, pars 82, 499–988.
- Bernhauer M, Schubert K (1911) Staphylinidae II. In: Schenkling S (Ed) Coleopterorum Catalogus. W. Junk, Berlin, pars 29: 87–190.
- Bierig A (1934a) Neues aus der Staphyliniden-Gattung *Cafius* (Col.), nebst Beschreibung neuer Arten aus Kuba und Nordamerika. Revista de Entomologia (Rio de Janeiro) 4: 65–70.
- Bierig A (1934b) Un género y especies nuevas de los Paederinae (Col. Staph.) de Cuba. Memorias de la Sociedad Cubana de Historia Natural „Felipe Poey“ 8: 325–330.
- Blackburn T (1888) Notes on Australian Coleoptera, with descriptions of new species. Transactions of the Royal Society of South Australia 10: 12–30, 36–51.
- Blackburn T (1902) Further notes on Australian Coleoptera, with descriptions of new genera and species. XXX. Transactions of the Royal Society of South Australia 26: 16–30.
- Blackwelder RE (1943) Monograph of the West Indian beetles of the family Staphylinidae. US National Museum Bulletin 182: i–viii, 1–658.
- Blackwelder RE (1952) The generic names of the beetle family Staphylinidae with an essay on genotypy. US National Museum Bulletin 200: i–iv, 1–483.
- Boheman CH (1858) Coleoptera. Species novas descripsit. In: Kongliga Svenska Fregatten Eugénies resa omkring jorden under befäl af C.A. Virgin, Åren 1851–1853. Zoologi 1, Insekter. Kongliga Svenska Vetenskapsakademien, Stockholm, 1–112 + 1 pl.
- Bondroit J (1912) Notes sur les *Bledius* Mann. (Staphylinidae). Annales de la Société Entomologique de Belgique 56: 63–68.
- Brendel E (1865a) On some new species of Pselaphidae. Proceedings of the Entomological Society of Philadelphia 5: 28–32.
- Brendel E (1865b) New species and corrections in the family Pselaphidae. Proceedings of the Entomological Society of Philadelphia 5: 255–260.

- Brendel E (1866) Descriptions of some new species of Pselaphidae. Proceedings of the Entomological Society of Philadelphia 6: 189–194.
- Brendel E (1894) On some Pselaphidae. Entomological News 5: 158–160, pl. 5.
- Brèthes J (1925) Un coléoptère et un diptère nouveaux de la Georgie du Sud. Comunicaciones del Museo Nacional de Historia Natural «Bernardo Rivadavia» 2 (16): 169–173.
- Brisout C (1859) Diagnoses de plusieurs espèces nouvelles du genre *Homalota* propres à la France. Bulletin de la Société Entomologique de France (1859): ccxvii–ccxix.
- Brisout C (1863) *Leptusa testacea*. In: Allard E, Aubé C, Brisout de Barneville C, Chevrolat A, Fairmaire L, Fauvel A, Grenier A, Kraatz G, Linder J, Reiche L, de Saulcy F, Matériaux pour la faune française: 16. In: Grenier A, Catalogue des coléoptères de France et Matériaux pour servir à la faune des coléoptères français. Grenier, Paris, 1–130. [note: these pages, 1–130, form a separately-paged appendix to the book]
- Bro Larsen E (1936) Biologiske Studien über tunnelgrabende Käfer auf Skallingen. Videnskabelige Meddelelser fra Dansk Naturhistorisk Forening i København 100: 1–232, pls. 1–2.
- Bro Larsen E (1951) Studies on the soil fauna of Skallingen. Qualitative and quantitative studies on alterations in the beetle fauna during five years' natural development of some sand and salt-marsh biotopes. Oikos 3: 166–192.
- Bro Larsen E (1952) On subsocial beetles from the salt-marsh, their care of progeny and adaptation to salt and tide. Transactions of the 9<sup>th</sup> International Congress of Entomology (Amsterdam, 1951): 502–506.
- Broun T (1880) Manual of the New Zealand Coleoptera. Colonial Museum and Geological Survey Department, Wellington, Part 1, xix + 651 pp.
- Broun T (1886) Manual of the New Zealand Coleoptera. Colonial Museum and Geological Survey Department, Wellington, Parts 3–4, i–viii, 745–973.
- Broun T (1893) Manual of the New Zealand Coleoptera. New Zealand Institute, Wellington, Parts 5, 6, and 7. v–xvii, 975–1504.
- Broun T (1894) Descriptions of new Coleoptera from New Zealand. Annals and Magazine of Natural History (ser. 6) 14: 302–312, 379–386, 419–428.
- Broun T (1903) Descriptions of new genera and species of New Zealand Coleoptera. Annals and Magazine of Natural History (ser. 7) 11: 602–618.
- Broun T (1907) Descriptions of six new species of Coleoptera from New Zealand. Annals and Magazine of Natural History (ser. 7) 19: 56–63.
- Broun T (1909a) Descriptions of Coleoptera from the subantarctic islands of New Zealand; with remarks on the affinities of the genera, etc. In: Chilton C, The subantarctic islands of New Zealand. Reports on the geo-physics, geology, zoology, and botany of the islands lying to the south of New Zealand. Philosophical Institute of Canterbury, Christchurch, Vol. 1, 78–123.
- Broun T (1909b) Descriptions of new genera and species of New-Zealand Coleoptera. Annals and Magazine of Natural History (ser. 8) 3: 223–233.
- Broun T (1910) Descriptions of new genera and species of Coleoptera. New Zealand Institute Bulletin 1: 1–78.
- Broun T (1911) Additions to the coleopterous fauna of the Chatham Islands. Transactions of the New Zealand Institute 43: 92–115.

- Broun T (1912) Descriptions of new genera and species of Coleoptera. Transactions of the New Zealand Institute 44: 379–440.
- Broun T (1914) Descriptions of new genera and species of Coleoptera. Part II. New Zealand Institute Bulletin 1: 79–142.
- Brundin L (1943) Zur Kenntnis einiger in die *Atheta*-Untergattung *Metaxya* M. & R. gestellten Arten (Col. Staphylinidae). Lunds Universitets Årsskrift N.F., Avd. 2, 39: 3–37, pls. 1–7.
- Butler EA (1909) *Myrmecopora brevipes*, a new species allied to *M. uvida*, Er., and an addition to the British fauna. Entomologist's Monthly Magazine 45: 29–31, 63.
- Cals P (1964) Modification de la mode de vie du parasite *Aleochara algarum* Fauvel (Coleoptera Staphylinidae) en fonction d'une variation de coincidence phénologique avec son hôte *Coelopa frigida* Fallén (Diptera Coelopidae). Annales de la Société Entomologique de France 89: 760–766.
- Cameron M (1904) Description of two new species of *Diglossa* (*Diglotta*) from the island of Perim. Entomologist's Monthly Magazine 40: 157–158.
- Cameron M (1912) Description of a new species of *Bledius* (subgen. *Belidus*) from the Red Sea. Entomologist's Monthly Magazine 48: 28–29.
- Cameron M (1914) *Bledius bernhaueri*, Cam.; change of name. Entomologist's Monthly Magazine 50: 203.
- Cameron M (1917a) On a new group of Staphylinidae. Entomologist's Monthly Magazine 53: 123–125.
- Cameron M (1917b) Description of a new genus of Staphylinidae. Entomologist's Monthly Magazine 53: 125.
- Cameron M (1917c) Description of a new genus of Staphylinidae. Entomologist's Monthly Magazine 53: 154–155.
- Cameron M (1917d) Description of a new species of *Thinobius*. Entomologist's Monthly Magazine 53: 155.
- Cameron M (1917e) The genus *Paraphytosus* mihi: synonymical note. Entomologist's Monthly Magazine 53: 233–234.
- Cameron M (1917f) Synonymic note on the group Arpediopsini. Entomologist's Monthly Magazine 53: 277.
- Cameron M (1918) New species of Staphylinidae from Singapore. Part I. Transactions of the Royal Entomological Society of London (1918): 58–90.
- Cameron M (1919) New species of Staphylinidae from Singapore. Part II. Transactions of the Royal Entomological Society of London (1919): 231–246.
- Cameron M (1920) New species of Staphylinidae from Singapore. Part III. Transactions of the Royal Entomological Society of London (1920): 212–284.
- Cameron M (1922) Descriptions of new species of Staphylinidae from the West Indies. Part II. Annals and Magazine of Natural History (ser. 9) 9: 113–128.
- Cameron M (1927) Description of a new species of *Cafius* (Col. Staphylinidae) from the Red Sea. Entomologist's Monthly Magazine 63: 251.
- Cameron M (1928) Fauna sumatrensis. Beitrag No. 54. Staphylinidae (Col.) Entomologische Mitteilungen 17: 90–110.
- Cameron M (1934) The staphylinid fauna of the New Hebrides. Stylops 3: 20–24.

- Cameron M (1936) New species of Staphylinidae (Col.) from the Malay Peninsula. Journal of the Federated Malay States Museums 18: 40–53.
- Cameron M (1938) New species of Staphylinidae from Mauritius and Reunion, collected by Mr. J. Vinson. Entomologist's Monthly Magazine 74: 173–174.
- Cameron M (1940) New species of Oriental Staphylinidae. Entomologist's Monthly Magazine 76: 181–184.
- Cameron M (1941) New species of Staphylinidae (Coleoptera) from the Philippines. Annals and Magazine of Natural History (ser. 11) 7: 430–447.
- Cameron M (1943) New species of Staphylinidae (Col.) from Australia and New Guinea. Annals and Magazine of Natural History (ser. 11) 10: 336–354.
- Cameron M (1944a) New oriental Staphylinidae. Annals and Magazine of Natural History (ser. 11) 11: 312–322.
- Cameron M (1944b) New species of Staphylinidae (Col.) from the Falkland Islands. Annals and Magazine of Natural History (ser. 11) 11: 618–621.
- Cameron M (1944c) New species of Staphylinidae (Col.) from China and Japan. Entomologist's Monthly Magazine 80: 158–159.
- Cameron M (1945a) New species of South African Staphylinidae (Col.). Annals and Magazine of Natural History (ser. 11) 11: 705–730.
- Cameron M (1945b) Some observations on the Staphylinidae of the Broun collection in the British Museum, with descriptions of new genera and species. Annals and Magazine of Natural History (ser. 11) 11: 779–793.
- Cameron M (1945c) Some observations on the Broun Collection of Coleoptera in the British Museum, with descriptions of new genera and species. Annals and Magazine of Natural History (ser. 11) 12: 158–180.
- Cameron M (1947a) New species of Staphylinidae (Col.) from Mauritius. Proceedings of the Royal Entomological Society of London (ser. B) 16: 114–119.
- Cameron M (1947b) New species of Staphylinidae (Col.) from New Zealand. Annals and Magazine of Natural History (ser. 11) 13: 701–710.
- Cameron M (1948) Some species of Staphylinidae (Col.) from New Zealand. Annals and Magazine of Natural History (ser. 11) 14: 723–731.
- Caron E, Ribeiro-Costa CS (2008) First record of the tribe Diglottini from South America with description of *Diglotta brasiliensis* n. sp. (Coleoptera, Staphylinidae, Aleocharinae). Zootaxa 1776: 52–58.
- Casey TL (1885) New genera and species of Californian Coleoptera. Bulletin of the California Academy of Sciences 1: 285–336, pl. 1.
- Casey TL (1886) Descriptive notices of North American Coleoptera I. Bulletin of the California Academy of Sciences 2: 157–264, pl. 7.
- Casey TL (1887) On some new North American Pselaphidae. Bulletin of the California Academy of Sciences 2: 455–482, pl. 16.
- Casey TL (1889a) A preliminary monograph of the North American species of *Trogophloeus*. Annals of the New York Academy of Sciences 4: 322–383. doi:10.1111/j.1749-6632.1889.tb57043.x

- Casey TL (1889b) Coleopterological notices, I. Annals of the New York Academy of Sciences 5: 39–96. doi:10.1111/j.1749-6632.1890.tb57003.x
- Casey TL (1894) Coleopterological notices, V. Annals of the New York Academy of Sciences 7: 281–606, pl. 1.
- Casey TL (1904) On some new Coleoptera, including five new genera. Canadian Entomologist 36: 312–324. doi:10.4039/Ent36312-11
- Casey TL (1905) A revision of the American Paederini. Transactions of the Academy of Sciences of St. Louis 15: 17–248.
- Casey TL (1906) Observations on the staphylinid groups Aleocharinae and Xantholinini, chiefly of America. Transactions of the Academy of Science of St. Louis 16: 125–434.
- Casey TL (1910) New species of the staphylinid tribe Myrmedoniini. Memoirs on the Coleoptera (Lancaster, PA) 1: 1–183.
- Casey TL (1911) New American species of Aleocharinae and Myllaeninae. Memoirs on the Coleoptera (Lancaster, PA) 2: 1–245.
- Chamberlin JC, Ferris GF (1929) On *Liparocephalus* and allied genera (Coleoptera; Staphylinidae). Pan-Pacific Entomologist 5: 137–143, 153–162.
- Champion GC (1887) On the priority of various generic names in use in British Coleoptera. Entomologist's Monthly Magazine 23: 227–230.
- Champion GC (1899) Some remarks on the two species of *Diglossa* Haliday, occurring in Britain. Entomologist's Monthly Magazine 35: 264–265.
- Chandler DS (1992) Notes on *Briaraxis depressa* (Coleoptera: Pselaphidae). Entomological News 103: 15–18.
- Chandler DS (1997) A catalog of the Coleoptera of America north of Mexico. Family: Pselaphidae. USDA Handbook 529–31: ix + 118 pp.
- Chandler DS (2001) Biology, morphology, and systematics of the ant-like litter beetle genera of Australia (Coleoptera: Staphylinidae: Pselaphinae). Memoirs on Entomology International 15: i–viii, 1–560.
- Chandler DS (2002) 5. Subfamilia Pselaphinae Latreille, 1802. In: Navarrete–Heredia JL, Newton AF, Thayer MK, Ashe JS, Chandler DS, Guía ilustrada para los géneros de Staphylinidae (Coleoptera) de México [Illustrated guide to the genera of Staphylinidae (Coleoptera) of Mexico]. Comisión Nacional para el Conocimiento y Uso de la Biodiversidad (CONABIO). Universidad de Guadalajara, 49–102.
- Chelazzi G, Chelazzi L, Focardi S (1983) Dynamic zonation of staphylinoid beetles (Coleoptera: Staphylinidae) on a sandy beach in East Africa. In: McLachlan A, Erasmus T (Eds) Sandy beaches as ecosystems. W. Junk, Hague, 405–412.
- Cheng L (1976) Marine insects. North Holland, Amsterdam, xii + 581 pp.
- Chevrolat A (1843) Description d'une nouvelle espèce du genre *Myrmedonia*. Revue Zoologique (1843): 42–43.
- Cho YB (2008) Korean species of the genus *Bisnius* Stephens (Coleoptera: Staphylinidae). Entomological Research 38: 90–92. doi:10.1111/j.1748-5967.2008.00138.x
- Coiffait H (1956) Les "*Staphylinus*- et genres voisins de France et des régions voisines. Mémoires du Muséum National d'Histoire Naturelle (Paris) (n.s.) série A, Zoologie 8: 177–224.



- Coiffait H (1963) Classification des Philonthini européens. Description de formes nouvelles. *Revue Française d'Entomologie* 30: 5–29.
- Coiffait H (1974) Coléoptères Staphylinidae de la région paléarctique occidentale. II. Sous famille Staphilininae [sic] Tribus Philonthinae et Staphylininae. *Nouvelle Revue d'Entomologie* 4 (4) Supplément 3: 1–593.
- Coiffait H (1981a) Fam. Staphylinidae, Subfam. Xantholininae, Staphylininae, Paederinae, Oxytelinae, Aleocharinae. In: Wittmer W, Büttiker W (Eds) *Fauna of Saudi Arabia*. Ciba-Geigy, Basle, Switzerland, (Part 2) Vol. 3, 236–242.
- Coiffait H (1981b) Contribution à la connaissance des Staphylinidae (Coleoptera) des Îles Andaman. *Bollettino del Museo Civico di Storia Naturale di Verona* 7: 329–348.
- Craig PC (1970) The behavior and distribution of the intertidal sand beetle *Thinopinus pictus* (Coleoptera: Staphylinidae). *Ecology* 51: 1012–1017. doi:10.2307/1933627
- Curtis J (1830) *British entomology; being illustrations and descriptions of the genera of insects found in Great Britain and Ireland: containing coloured figures from nature of the most rare and beautiful species, and in many instances of the plants upon which they are found.* Curtis; London. edn.1, Vol. 7, pls. 290–337.
- Curtis J (1838) *British entomology; being illustrations and descriptions of the genera of insects found in Great Britain and Ireland: containing coloured figures from nature of the most rare and beautiful species, and in many instances of the plants upon which they are found.* Curtis; London. edn. 1, Vol. 15, pls. 674–721.
- Donisthorpe H St JK (1930) An annotated list of the additions to the British coleopterous fauna, made since the publication of the supplementary volume (VI) of Fowler's 'Coleoptera of the British Islands'. *Entomologist's Monthly Magazine* 66: 23–31, 94–101.
- Easton AM (1971) A new British *Atheta* (Col., Staphylinidae) in a new subgenus. *Entomologist's Monthly Magazine* 107: 24–26.
- Edward DA, Blyth JE, McKee R, Gilburn AS (2007) Change in the distribution of the strand line community: the seaweed fly (Diptera: Coelopidae). *Ecological Entomology* 32: 741–746. doi:10.1111/j.1365-2311.2007.00919.x
- Egglishaw J (1965) Observations on the fauna of wrack beds. *Transactions of the Society for British Entomology* 16: 189–216.
- Elliott P, King PE, Fordy MR (1983) Observations on staphylinid beetles living on rocky shores. *Journal of Natural History* 17: 575–581. doi:10.1080/00222938300770471
- Emberson RM (1998) The beetle (Coleoptera) fauna of the Chatham Islands. *New Zealand Entomologist* 21: 25–62.
- Enderlein G (1909) Die Insekten des antarktischen Gebietes. In: Drygalski E von (Ed) *Deutsche südpolare-Expedition 1901–1903*. Georg Reimer, Berlin, Bd. 10. *Zoologie* II: 361–532 + xl–lxiii pls.
- Eppelsheim E (1885) Beitrag zur Staphylinenfauna West-Afrika's. *Deutsche Entomologische Zeitschrift* 29: 97–147.
- Erichson WF (1837) *Die Käfer der Mark Brandenburg*. Morin, Berlin, Vol. 1, i–viii, 1–384.
- Erichson WF (1839a) *Die Käfer der Mark Brandenburg*. Morin, Berlin, Vol. 1, 385–740.
- Erichson WF (1839b) *Genera et species Staphylinorum, Insectorum Coleopterorum familiae*. Part 1, Morin, Berlin, 1–400.

- Erichson WF (1840) Genera et species Staphylinorum, Insectorum Coleopterorum familiae. Morin, Berlin, Part 2, 401–954.
- Esaki T (1956) Biological notes on *Brachypronomaea esakii* Sawada. Kontyû 24: 199–200.
- Evans PD, Ruscoe CNE, Treherne JE (1971) Observations of the biology and submergence behaviour of some littoral beetles. Journal of the Marine Biological Association of the United Kingdom 51: 375–386. doi:10.1017/S0025315400031854
- Evans WG (1980) Insecta, Chilopoda and Arachnida. In: Morris RH, Abbott DP and Hadlerlie EC (Eds) Intertidal invertebrates of California. Stanford University Press, Stanford, 641–658.
- Fagel G (1969) Contribution à la connaissance des Staphylinidae CVIII. Sur quelques espèces inédites provenant du bassin méditerranéen. Bulletin et Annales de la Société Royale d'Entomologie de Belgique 105: 103–125.
- Fagel G (1970) Contribution à la connaissance des Staphylinidae. CX. Sur quelques espèces du bassin de la Méditerranée orientale. Bulletin et Annales de la Société Royale d'Entomologie de Belgique 106: 145–164.
- Fairmaire L (1849) Essai sur les coléoptères de la Polynésie. Revue et Magasin de Zoologie (sér. 2) 1: 277–291.
- Fairmaire L (1852) Excursion entomologique dans la baie de la Somme. Annales de la Société Entomologique de France (sér. 2) 10: 663–691.
- Fairmaire L (1856) In: Fairmaire L, Laboulbène A, Faune entomologique française ou description des insectes qui se trouvent en France. Coléoptères. Deyrolle, Paris, vol. 1, 371–665.
- Fairmaire L (1861) Miscellanea entomologica. Annales de la Société Entomologique de France (sér. 4) 1: 577–596.
- Fairmaire L (1892) Coléoptères d'Obock. Troisième partie. Revue d'Entomologie (Caen) 11: 77–127.
- Fauvel A (1862a) [Une nouvelle espèce de coléoptères de la famille des staphyliniens propres à la France]. Bulletin de la Société Entomologique de France (sér. 4) 1: xxiv–xxxv.
- Fauvel A (1862b) Notice sur quelques aléochariens nouveaux ou peu connus et description de larves de *Phytosus* et *Leptusa*. Annales de la Société Entomologique de France (sér. 4) 2: 81–94, pl. 2. [note: this writer, Charles Alphonse Albert Fauvel, published under the name Albert Fauvel].
- Fauvel A (1862c) Diagnoses d'un nouveau genre et de deux espèces nouvelles de staphylinides de France. Annales de la Société Entomologique de France (sér. 4) 2: 292.
- Fauvel A (1868) Faune du Chili. Insectes coléoptères. Staphylinides. (suite et fin). Bulletin de la Société Linnéenne de Normandie (sér. 2): 1: 6–67, pl. 1.
- Fauvel A (1871) Staphylinides auct. Microptera Grav.—Brachélytres Latr. [formant la suite de sa Faune Gallo-Rhénane]. Bulletin de la Société Linnéenne de Normandie (sér. 2) 5 (1869–1870): 27–192.
- Fauvel A (1877) Les staphylinides de l'Australie et de la Polynésie. Annali del Museo Civico di Storia Naturale di Genova 10: 168–298.
- Fauvel A (1878a) Les staphylinides des Moluques et de la Nouvelle Guinée. Annali del Museo Civico di Storia Naturale di Genova 12: 171–315.

- Fauvel A (1878b) Les staphylinides de l'Australie et de la Polynésie (2<sup>e</sup> mémoire). Annali del Museo Civico di Storia Naturale di Genova 13: 465–598.
- Fauvel A (1879) Staphylinides nouveaux de la coll. Godeffroy. 83–84. In: Fairmaire L, Descriptions de coléoptères nouveaux ou peu connus du Musée Godeffroy. Journal du Musée Godeffroy 14: 80–114.
- Fauvel A (1889) Les Coléoptères de la Nouvelle-Calédonie et dépendances avec descriptions, notes et synonymies nouvelles (Suite). Revue d'Entomologie (Caen) 8: 242–271, 277–285.
- Fauvel A (1900) Sur les Oxyteliens de Nouvelle-Zélande. Revue d'Entomologie (Caen) 19: 181–189.
- Fauvel A (1901) Voyage de M. le Dr. Ed. Bugnion au Venezuela, en Colombie et aux Antilles. Staphylinides (Planches I et II). Revue d'Entomologie (Caen) 20: 69–91.
- Fauvel A (1902) Catalogue des staphylinides de la Barbarie, de la Basse-Égypte et des Îles Açores, Madères, Salvages et Canaries. Revue d'Entomologie (Caen) 21: 45–189.
- Fauvel A (1903) Mission de M. Maurice Maindron dans l'Inde meridionale. Revue d'Entomologie (Caen) 22: 149–163.
- Fauvel A (1904a) Staphylinides nouveaux du Sinaï et de la mer Rouge. Revue d'Entomologie (Caen) 23: 71–74.
- Fauvel A (1904b) Staphylinides exotiques nouveaux. 2<sup>e</sup> partie. Revue d'Entomologie (Caen) 23: 76–112.
- Fauvel A (1904c) Staphylinides nouveaux de Madagascar. Revue d'Entomologie (Caen) 23: 296–322.
- Fauvel A (1905) Staphylinides exotiques nouveaux. 3<sup>e</sup> partie. Revue d'Entomologie (Caen) 24: 113–147.
- Fenyès A (1909) New Aleocharinae (Staphylinidae, Col.) of the U.S. Entomological News 20: 418–425.
- Fenyès A (1921) Fenyès A (1921) New genera and species of Aleocharinae with a polytomic synopsis of the tribes. Bulletin of the Museum of Comparative Zoology 65 (2): 17–36.
- Frank JH (1982) The parasites of the Staphylinidae. A contribution towards an encyclopedia of the Staphylinidae. Florida Agricultural Experiment Stations Technical Bulletin 824: 1–118.
- Frank JH (1985a) New records of *Cafius caribeanus* and *C. subtilis* in Florida and Venezuela. Entomological News 96: 61–62.
- Frank JH (1985b) Associations of Scaritini and *Bledius* in the New World. Florida Entomologist 68: 480–482. doi:10.2307/3495137
- Frank JH, Carlysle TC, Rey JR (1986) Biogeography of the seashore Staphylinidae *Cafius bis-triatus* and *C. rufifrons* (Insecta: Coleoptera). Florida Scientist 49: 148–161.
- Frank JH, Thomas MC (1984a) Cocoon-spinning and the defensive function of the median gland in larvae of Aleocharinae (Coleoptera, Staphylinidae): a review. Quaestiones Entomologicae 20: 7–23.
- Frank JH, Thomas MC (1984b) *Heterota plumbea* and *Coenonica puncticollis* in Florida (Coleoptera: Staphylinidae). Florida Entomologist 67: 409–417. doi:10.2307/3494720
- Froggatt WW (1907) Australian insects. William Brooks, Sidney, 449 pp. + 37 pls.

- Ganglbauer L (1895) Die Käfer von Mitteleuropa. Die Käfer der österreichisch-ungarischen Monarchie, Deutschlands, der Schweiz, sowie des französischen und italienischen Alpengebietes. Zweiter Band. Familienreihe Staphylinoidea. 1. Theil. Staphylinidae, Pselaphidae. Carl Gerold's Sohn, Vienna, 880 pp.
- García-Castellanos D, Estrada F, Jiménez-Munt I, Gorini, C, Fernández M, Vergés J, Vicente R De (2009) Catastrophic flood of the Mediterranean after the Messinian salinity crisis. *Nature* 462: 778–782. doi:10.1038/nature08555
- Gemminger M, von Harold E (1868) Catalogus Coleopterorum hucusque descriptorum synonymicus et systematicus. Gummi, Munich, Vol. 2, 425–752.
- Gestro R (1889) Viaggio ad Assab, nel Mar Rosso, dei Signori G. Doria ed O. Beccari con il R. avviso „Esploratore“ dal 16 Novembre 1879 al 26 Febraio 1880. IV. Coleotteri. Annali del Museo Civico di Storia Naturale di Genova (ser. 2) 7: 5–72.
- Good JA (1998) The identification and habitat of micropterous *Diglossa mersa* (Haliday) and *D. sinuaticollis* (Mulsant & Rey) (Staphylinidae). *Coleopterist* 7: 73–76.
- Gravenhorst JLC (1802) Coleoptera Microptera Brunsvicensia nec non exoticorum quotquot exstant in collectionibus entomologorum Brunsvicensium in genera familias et species distribuit. Reichard, Brunswick, 206 pp.
- Gravenhorst JLC (1806) Monographia Coleopterorum Micropterorum. Dieterich, Gottingen, xvi + 236 pp. + 1 pl.
- Grebennikov VV, Newton AF (2009) Good-bye Scydmaenidae, or why the ant-like stone beetles should become megadiverse Staphylinidae, sensu latissimo (Coleoptera). *European Journal of Entomology* 106: 275–301.
- Griffiths CL, Griffiths RJ (1983) Biology and distribution of the littoral rove beetle *Psammathobledius punctatissimus* (LeConte) (Coleoptera: Staphylinidae). *Hydrobiologia* 101: 203–214. doi:10.1007/BF00009876
- Gusarov VI (2003a) Revision of some types of North American aleocharines (Coleoptera: Staphylinidae; Aleocharinae), with synonymic notes. *Zootaxa* 185: 1–134.
- Gusarov VI (2003b) A revision of Nearctic species of the genera *Adota* Casey, 1910 and *Psammotiba* Yosii & Sawada, 1976 (Coleoptera: Staphylinidae; Aleocharinae). *Zootaxa* 353: 1–35.
- Gusarov VI (2004) A revision of the Nearctic species of the genus *Halobrecta* Thomson, 1858 (Coleoptera: Staphylinidae; Aleocharinae) with notes on some Palearctic species of the genus. *Zootaxa* 746: 1–25.
- Gusarov VI, Makranczy G (2004) *Yosiityphlus* Sawada, 1971, a new synonym of *Thinobius* Kiesenwetter, 1844 (Coleoptera: Staphylinidae). *Zootaxa* 748: 1–4.
- Gyllenhal L (1810) Insecta Suecica descripta. Classis I. Coleoptera sive Eleuterata. Tom. I pars II. Leverentz, Skara, Sweden, xviii + 658 pp.
- Gyllenhal L (1827) Insecta Suecica descripta. Classis I. Coleoptera sive Eleuterata. Tom. I pars IV. Fleischer, Leipzig, x + 761 pp.
- Haghebaert G (1991) A review of *Diglossa* of the world (Coleoptera, Staphylinidae, Aleocharinae). *Bulletin et Annales de la Société Royale Belge d'Entomologie* 127: 223–234.

- Haghebaert G (1993) Description of two new *Phytosus* species, with notes on the known species (Coleoptera, Staphylinidae, Aleocharinae). Entomologische Blätter für Biologie und Systematik der Käfer 89: 160–166.
- Haghebaert G (1995) The first New Guinean records of the genus *Bryothinusa* Casey, 1904 with descriptions of three new species (Coleoptera, Staphylinidae, Aleocharinae). Bulletin et Annales de la Société Royale Belge d'Entomologie 131: 25–33.
- Haliday AH (1837) Notes about *Cilenum laterale* and a submarine species of Aleocharidae. Entomological Magazine 4: 251–253.
- Hammond PM (1975) The phylogeny of a remarkable new genus and species of gymnosine staphylinid (Coleoptera) from the Auckland Islands. Journal of Entomology (series B) 44: 153–173.
- Hammond PM (2000) Coastal Staphylinidae (rove beetles) in the British Isles, with special reference to saltmarshes. In: Sherwood BR, Gardiner BG, Harris T (Eds) British saltmarshes. Joint symposium on British Saltmarshes organized between the Linnean Society of London, the Royal Society for the Protection of Birds, and English Nature (London; 2000). Forrest Text, Cardigan, Wales, 247–302.
- Hardy J (1851) Staphylinidae Leach. In: Hardy J, Bold TJ, A catalogue of the insects of Northumberland and Durham. Coleoptera. Transactions of the Tyneside Naturalists' Field Club 2 (1): 21–97 [part]; also as 59–134 in Part II (1851) of a reprint edition. F and W Dodsworth, Newcastle upon Tyne [page 78, cited in the text, is page 116 of the reprint edition].
- Hasegawa M, Kanie N (1992) Coleoptera of sea-shore from Omotehama, Toyohashi, Japan. Toyohashi-shi Shizenshi Hakubutsukan Kenkyu Hokoku (Science Report of the Toyohashi Museum of Natural History) 2: 13–24.
- Hatch MH (1957) The beetles of the Pacific Northwest. Part II: Staphylinidae. University of Washington Press, Seattle, ix + 384 pp.
- Hayward P, Nelson-Smith A, Shields C (1996) Sea shore of Britain and Europe (Collins Pocket Guide). Harper Collins, London, 352 pp.
- Herman LH (1972) Revision of *Bledius* and related genera. Part I. The *aequatorialis*, *mandibularis*, and *semiferugineus* groups and two new genera (Coleoptera, Staphylinidae, Oxytelinae). Bulletin of the American Museum of Natural History 149: 111–254.
- Herman LH (1976) Revision of *Bledius* and related genera. Part II. The *armatus*, *basalis*, and *melanocephalus* groups (Coleoptera, Staphylinidae, Oxytelinae). Bulletin of the American Museum of Natural History 157: 71–172.
- Herman LH (1983) Revision of *Bledius*. Part III. The *annularis* and *emarginatus* groups (Coleoptera, Staphylinidae, Oxytelinae). Bulletin of the American Museum of Natural History 175: 1–146.
- Herman LH (1986) Revision of *Bledius*. Part IV. Classification of species groups, phylogeny, natural history, and catalogue (Coleoptera, Staphylinidae, Oxytelinae). Bulletin of the American Museum of Natural History 184: 1–368.
- Herman LH (2002) Case 3207. Staphylinidae Latreille, 1804 (Insecta, Coleoptera): proposed conservation of 65 specific names. Bulletin of Zoological Nomenclature 59: 99–113.

- Hoebeke ER (1985) A revision of the rove beetle tribe Falagriini of America north of Mexico (Coleoptera: Staphylinidae: Aleocharinae). *Journal of the New York Entomological Society* 93: 913–1018.
- Hogarth PJ (2007) *The biology of mangroves and seagrasses*, 2nd ed. Oxford Univ. Press, Oxford, x + 273 pp.
- Hollander J den (1983) Arthropod life in sandy shores. In: Wolff WJ (Ed) *Terrestrial and freshwater fauna of the Wedden Sea area*. Balkema, Rotterdam, 84–88.
- Hollander J den, Etten JCA van (1974) De oekologie van *Bledius arenarius* en *B. subniger* op het Oostvoornse strand (Coleoptera, Staphylinidae). *Entomologische Berichten* 34: 155–160.
- Holme F (1837) Notice of the coleopterous insects observed in the Scilly Islands in July and August, 1836. *Transactions of the Entomological Society of London* 2: 58–64.
- Horn GH (1871) Descriptions of new Coleoptera of the United States with notes on known species. *Transactions of the American Entomological Society* 3: 325–344.
- Horn GH (1884) Synopsis of the Philonthi of boreal America. *Transactions of the American Entomological Society* 11: 177–244.
- Hughes M, Weir A, Leschen R, Judd C, Gillen B (2004) New species and records of Laboulbeniales from the subantarctic islands of New Zealand. *Mycologia* 96: 1355–1369. doi:10.2307/3762152
- Irmeler U, Heller K (2002) Zonierung der Staphylinidae in einem Salzgrünland der schleswig-holsteinischen Nordseeküste. *Faunistisch-Oekologische Mitteilungen* 8: 219–229.
- Jałoszyński P (2003) *Chelonoidum araiorum* sp. nov., an unusual species of the Cephenniini (Coleoptera: Scydmaenidae) from the intertidal zone of the Pacific coast of central Honshu, Japan. *Bulletin of the National Science Museum, Tokyo. Ser. A.* 29 (4): 225–228.
- James GJ, Moore I, Legner EF (1971) The larval and pupal stages of four species of *Cafius* (Coleoptera: Staphylinidae) with notes on their biology and ecology. *Transactions of the San Diego Natural History Society* 16: 279–290.
- Jarrige J (1959) Un nouveau genre d'aleocharide marin de la zone intertropicale. *Naturaliste Malgache* 11: 63–66.
- Jarrige J (1964) Brachélytres recoltés en Nouvelle-Calédonie par M.Y. Plessis. *Cahiers du Pacifique* 6: 177–180.
- Jarrige J (1973) Un *Heterota* malgache nouveau (Col., Aleocharidae). *Bulletin de la Société Entomologique de France* 78: 256–259.
- Jeannel R (1940) III. Coléoptères In: *Croisière du Bougainville aux îles australes françaises*. Mémoires du Muséum National d'Histoire Naturelle, Paris (ser. 2) 19: 64–201.
- Jeannel R (1949) Les psélaphides de l'Afrique orientale (Coleoptera). *Mémoires du Muséum National d'Histoire Naturelle, Paris (n.s.)* 29: 1–226.
- Jeannel R (1952) Psélaphides de Saïgon. *Revue Française d'Entomologie* 19: 69–113.
- Jeannel R (1954) Les psélaphides de Madagascar. *Mémoires de l'Institut Scientifique de Madagascar (série E: Entomologie)* 4: 151–344.
- Jeon MJ, Ahn KJ (2002) First larval descriptions for *Cafius* Curtis (Coleoptera: Staphylinidae: Staphylininae) in Korea. *Journal of the Kansas Entomological Society* 78: 261–271. doi:10.2317/0410.21.1



- Jeon MJ, Ahn KJ (2007) Descriptions of late instars of three littoral *Cafius* species (Coleoptera: Staphylinidae) by association of life stage using DNA sequences. *Florida Entomologist* 90: 465–474. doi:10.1653/0015-4040(2007)90[465:DOLIOT]2.0.CO;2
- Jeon MJ, Ahn KJ (2008) Revision of the seashore genus *Salinamexus* (Coleoptera: Staphylinidae: Aleocharinae) with a description of *Salinamexus koreanus* sp. nov. from Korea. *Canadian Entomologist* 139: 189–194. doi:10.4039/n06-023
- Jeon MJ, Ahn KJ (2009) Description of late-instars of *Bryothinusa koreana* Ahn and Jeon (Coleoptera: Staphylinidae: Aleocharinae) by association of life stage based on DNA sequence data. *Florida Entomologist* 92: 367–373.
- Jones TW (1968) The zonal distribution of three species of Staphylinidae in the rocky intertidal zone in California. *Pan-Pacific Entomologist* 44: 203–210.
- Joy NH (1912) *Bledius arenarius*, var *fergussoni* var nov. *Entomologist's Monthly Magazine* 48: 44.
- Kemner NA (1925) Zur Kenntnis der Staphyliniden-Larven. I: Die Larven der Tribus Proteini-ni und Diglossini. *Entomologisk Tidskrift* 46: 61–77.
- Kiesenwetter EAH von (1844) Die Staphylinenfauna von Leipzig's Umgegen. *Stettiner Entomologische Zeitung* 5: 307–320, 340–356, 372–378.
- Kiesenwetter EAH von (1849) [Staphylinidae] species. In: Küster HC, Die Käfer Europa's. Nach der Natur beschrieben. Bauer und Raspe, Nuremberg, Heft 17, 17–20 [text occupies 8 pages which are not numbered]. [cf. Küster 1854].
- Kiesenwetter EAH von (1850) Funzig Diagnosen unbeschriebener oder wenig bekannter europäischer Käfer. *Stettiner Entomologische Zeitung* 11: 217–225.
- Kiesenwetter EAH von (1877) Staphylinidae. In: Kiesenwetter EAH von, Kirsch T, Die Käferfauna der Auckland-Inseln nach Herm. Krone's Sammlungen beschrieben. *Deutsche Entomologische Zeitschrift* 21: 160–164.
- Kincaid T (1961a) The ecology and morphology of *Thinobius frizzelli* Hatch, an intertidal beetle. *Calliostoma Co.*, Seattle, 12 pp. + 6 pls.
- Kincaid T (1961b) The staphylinid genera *Pontomalota* and *Thinusa*. *Calliostoma Co.*, Seattle, 10 pp. + 4 pls.
- King PE, Fordy M, Al-Khalifa MS (1979) Observations on the intertidal *Micralymma marinum* (Stroem) (Col., Staphylinidae). *Entomologist's Monthly Magazine* 115: 113–136.
- Kirschenblatt JD (1938). On some Far Eastern staphylinid beetles [in Russian]. *Trudy Gidrobiologicheskoi Ekspeditsii Zoologicheskogo Instituta Akademii Nauk v 1934 godu na Yaponskoe More* 1: 527–532. *Izdatelstvo Akademii Nauk USSR; Moscow*. [p. 533–536 of the same volume have a condensed version of the same article, in English]. We use the spelling of his family name and initials as given on p. 533. However, the author's first name, Яков, is normally transliterated into English as 'Yakov' with the first letter (Я) as Ya (it could be 'Ja' if transliterated into German). The author's family name Kirschenblatt is of German origin, transliterated into Russian as Киршенблат, and is best transliterated back from Russian in its original form (as given on p. 533), not as Kirshenblat. The page facing the title page provides an English translation of the title (Reports of the Japan Sea Hydrobiological Expedition of the Zoological Institute of the Academy of Sciences of the

- USSR in 1934. Part 1), editor (S. A. Sernov), and publisher (Academy of Sciences Press, Moscow). Apparently this was the only published volume].
- Klimaszewski J (1982a) Studies of Myllaenini (Coleoptera: Staphylinidae, Aleocharinae). 1. Systematics, phylogeny, and zoogeography of Nearctic *Myllaena* Erichson. Canadian Entomologist 114: 181–242.
- Klimaszewski J (1982b) A redefinition of Myllaenini Ganglbauer and redescription of *Camacopalpus* Motschulsky and *Polypea* Fauvel (Coleoptera: Staphylinidae). Canadian Entomologist 114: 411–429.
- Klimaszewski J (1984) A revision of the genus *Aleochara* Gravenhorst of America north of Mexico. Memoirs of the Entomological Society of Canada 129: 1–211.
- Klimaszewski J, Jansen RE (1994) Systematics, biology and distribution of *Aleochara* Gravenhorst from southern Africa. Part 3: Subgenus *Coprochara* Mulsant and Rey (Coleoptera: Staphylinidae). Annals of the Transvaal Museum 36: 147–170.
- Klimaszewski J, Majka CG (2007) Two new *Atheta* species (Coleoptera: Staphylinidae: Aleocharinae) from eastern Canada: taxonomy, bionomics, and distribution. Canadian Entomologist 139: 45–53.
- Klimaszewski J, Peck SB (1998) A review of the aleocharine rove beetles from the Galapagos Islands, Ecuador (Coleoptera: Staphylinidae, Aleocharinae). Revue Suisse de Zoologie 105: 221–260.
- Klimaszewski J, Watt JC (1997) Coleoptera: family-group review and keys to identification. Fauna of New Zealand 37: 1–199.
- Klimaszewski J, Majka CG, Langor D (2006) Review of the North American *Tarphiota* Casey, with a description of new seashore-inhabiting *Atheta* species exhibiting convergent characteristics (Coleoptera: Staphylinidae: Aleocharinae). Entomological Science 9: 67–78.
- Klimaszewski J, Maus C, Gardiner A (2002) The importance of tracking introduced species: New records of athetine rove beetles from South Atlantic Inaccessible Island (Coleoptera, Staphylinidae, Aleocharinae). Coleopterists Bulletin 56: 481–490.
- Klimaszewski J, Newton AF, Thayer MK (1996) A review of the New Zealand rove beetles (Coleoptera: Staphylinidae). New Zealand Journal of Zoology 23: 143–160.
- Klimaszewski J, Webster R, Assing V, Savard K (2008) *Diglotta mersa* and *Halobrecta flavipes* Thomson, two new species for the Canadian fauna (Coleoptera, Staphylinidae, Aleocharinae). Zookeys 2: 175–188.
- Koch C (1936) Wissenschaftliche Ergebnisse der entomologischen Expeditionen Seiner Durchlaucht des Fürsten Alessandro C. della Torre e Tasso nach Aegypten und auf die Halbinsel Sinai, XIII. Staphylinidae. Pubblicazioni del Museo Entomologico “Pietro Rossi” (Duino) 1: 115–232.
- Kraatz G (1856) Naturgeschichte der Insecten Deutschlands. Erste Abtheilung. Coleoptera. Zweiter Band. Nicolai, Berlin, 1–376.
- Kraatz G (1857) Naturgeschichte der Insecten Deutschlands. Erste Abtheilung. Coleoptera. Zweiter Band. Nicolai, Berlin, i–viii + 769–1080.
- Kraatz G (1859a) Die Staphylinen-Fauna von Ostindien, insbesondere der Insel Ceylan. Archiv für Naturgeschichte 25: 1–196.

- Kraatz G (1859b) *Phytosus balticus*. In: Schaum H, Beiträge zur europäischen Käferfauna zusammengestellt. Berliner Entomologische Zeitschrift 10: 52–53.
- Krogerus R (1925) Studien über Lebensweise und Entwicklung einiger *Bledius*-Arten. Acta Societatis pro Fauna et Flora Fennica 56 (3): 1–24.
- Küster HC (1854) Die Käfer Europa's. Nach der Natur beschrieben. Bauer und Raspe, Nuremberg, Heft 28, 100 p. [note: pages are not numbered consecutively. Instead, each species description is numbered independently, with that of *Myrmedonia tenuicornis* (Küster) having "XXVIII.3" at the foot of its first page (= vol. 28 species 3); that description occupies 2 otherwise unnumbered pages, followed by a description of *Myrmedonia limbata*, Paykull having "XXVIII.4" at the foot of its first page; cf. Kiesenwetter 1849].
- Larsson SG, Gigha G (1959) Coleoptera 1. Synopsis of the species. The Zoology of Iceland 3 (46a): 1–218.
- Last HR (1987) Staphylinidae from Papua New Guinea in the collection of the Bernice P. Bishop Museum Honolulu, Hawaii. Entomologische Abhandlungen (Dresden) 51: 25–56.
- Lawrence JF, Newton AF (2000) Families and subfamilies of Coleoptera (with selected genera, notes, references and data on family-group names). In: Pakaluk J, Slipinski SA (Eds) Biology, phylogeny and classification of Coleoptera. Papers celebrating the 80th birthday of Roy A. Crowson. Muzeum i Instytut Zoologii PAN, Warsaw, 779–1006.
- Lea AM (1910) On Australian and Tasmanian Coleoptera, with descriptions of new species. Part I. Proceedings of the Royal Society of Victoria 22 (1909): 113–152, pl. 30.
- Lea AM (1920) Description of a new staphylinid beetle from Macquarie Island. 30–31 [with further notes and illustrations by R.J. Tillyard, 18–20]. In: Tillyard RJ, The insects of Macquarie Island In Australasian Antarctic Expedition 1911–14 under the leadership of Sir Douglas Mawson. Scientific Reports. Series C-Zoology and Botany. Vol. V, Part 8, with appendices by Professor Brues CT and Lea AM. Govt. Printer, Sydney, 35 pp.
- Lea AM (1927) Descriptions of new Staphylinidae from Fiji. Records of the South Australian Museum 3: 273–278.
- Lea AM (1929) Notes on some miscellaneous Coleoptera, with descriptions of new species. Part VII. Transactions of the Royal Society of South Australia (1929): 203–244.
- Leach WE (1819) [New genera]. In: Samouelle G, The entomologist's useful compendium; or, an introduction to the knowledge of British insects, comprising the best means of identifying and preserving them, and a description of the apparatus generally used; together with the genera of Linné, and the modern method of arranging the classes Crustacea, Myriapoda, spiders, mites, and insects from their affinities, and structure, according to the views of Dr. Leach. Also an explanation of the terms used in entomology; a calendar of the times of appearance, and useful situations of near 3000 species of British insects; with instructions for collecting and fitting up objects for the microscope. Thomas Boys, London, 496 pp.
- LeConte JL (1849) On the Pselaphidae of the United States. Boston Journal of Natural History 6: 64–110.
- LeConte JL (1852) Descriptions of new species of Coleoptera from California. Annals of the Lyceum of Natural History of New York 5: 125–216.
- LeConte JL (1863) New species of North American Coleoptera. Part I. Smithsonian Miscellaneous Collections 6 (167): 1–186.

- LeConte JL (1877) On certain genera of Staphylinidae Oxytelini, Piestidae, and Micropeplidae, as represented in the fauna of the United States. Transactions of the American Entomological Society 6: 213–252.
- LeConte JL (1880) Short studies of North American Coleoptera. Transactions of the American Entomological Society 8: 163–218.
- Leech HB, Moore I (1971) Nearctic records of flights of *Cafius* and some related beetles at the seashore (Col., Staphylinidae and Hydrophilidae). Wasmann Journal of Biology 29: 65–70.
- Lengerken H von (1929) Die Salzkäfer der Nord- und Ostseeküste mit Berücksichtigung der angrenzenden Meere sowie des Mittelmeeres, des Schwarzen und des Kaspischen Meeres. Zeitschrift für Wissenschaftliche Zoologie 135: 1–162.
- Leschen RAB, Bullians MS, Michaux B, Ahn KJ (2002) Systematics of *Baeostethus chiltoni*, a subantarctic liparocephaline (Coleoptera: Staphylinidae: Aleocharinae): a Pangean relic or a more recent immigrant? Journal of the Royal Society of New Zealand 32: 189–201. doi: 10.1080/03014223.2002.9517691
- Lesne P, Mercier L (1922) Un staphylinide parasite des muscides fuscicoles *Aleochara (Polystoma) algarum* Fauvel. Caractères adaptatifs de la larve à la vie parasitaire. Annales de la Société Entomologique de France 91: 351–358.
- Little C, Williams GA, Trowbridge CD (2009) The biology of rocky shores. (2<sup>nd</sup> edn). Oxford Univ. Press, Oxford, 376 pp.
- Löbl I (1974) Beitrag zur Kenntnis der Pselaphidae (Coleoptera) der Koreanischen Volksdemokratischen Republik. Acta Zoologica Cracoviensia 19: 91–104.
- Löbl I (1977) Weitere Pselaphidae (Coleoptera) von der Koreanischen Volksdemokratischen Republik. Bulletin de l'Académie Polonaise des Sciences. Classe II. Série des Sciences Biologiques 25: 235–241.
- Lohse GA (1985) *Diglossa*—Studien. Entomologische Blätter für Biologie und Systematik der Käfer 81: 179–182.
- Lott DA (2008) Staphylinidae. In: Duff AG, Checklist of beetles of the British Isles. Wells, Somerset, UK: Duff, 2008 ed., 33–55.
- Majka CG, Klimaszewski J, Lauff RF (2008) The coastal rove beetles (Coleoptera, Staphylinidae) of Atlantic Canada: a survey and new records. ZooKeys 2: 115–150. doi:10.3897/zookeys.2.2
- Mäklin FG (1852) [New species and notes]. In: Mannerheim CG von, Zweiter Nachtrag zur Käfer-Fauna der nord-amerikanischen Länder des Russischen Reiches. Bulletin de la Société Impériale des Naturalistes de Moscou 25 (2): 283–387.
- Mäklin FG (1853) [Description of new taxa]. In: Mannerheim CG von, Dritter Nachtrag zur Käfer-Fauna der nord-amerikanischen Länder des Russischen Reiches. Bulletin de la Société Impériale des Naturalistes de Moscou 26 (3): 95–269.
- Mäklin FG (1878) Diagnoser öfver några nya siberiska insektarter. Öfversigt af Finnska Vetenskaps-Societetens Föreläsningar 19 (1876–1877): 15–32.
- Makranczy G (2006) Systematics and phylogenetic relationships of the genera in the *Carpe limus* group (Coleoptera: Staphylinidae: Oxytelinae). Annales Historico-Naturales Musei Nationalis Hungarici 98: 29–120.

- Malkin B (1958) Protective coloration in *Thinopinus pictus*. Coleopterists Bulletin 12: 20.
- Mannerheim CG von (1843) Beitrag zur Kaefer-Fauna der aleutischen Inseln, der Insel Sitka und Neu-Californiens. Bulletin de la Société Imperiale des Naturalistes de Moscou 16 (2): 175–314.
- Mannerheim CG von (1846) Nachtrag zur Kaefer-Fauna der aleutischen Inseln und der Insel Sitka. Bulletin de la Société Imperiale des Naturalistes de Moscou 19 (2): 501–516.
- Marris JWM (2000) The beetle (Coleoptera) fauna of the Antipodes Islands, with comments on the impact of mice; and an annotated checklist of the insect and arachnid fauna. Journal of the Royal Society of New Zealand 30 (2): 169–195. doi:10.1080/03014223.2000.9517616
- Maruyama M, Ahn KJ (2000a) The intertidal beetle *Paramblopusa borealis* (Casey) (Coleoptera, Staphylinidae, Aleocharinae) new to Japan. Japanese Journal of Systematic Entomology 6: 83.
- Maruyama M, Ahn KJ (2000b) Redescription of *Liparocephalus litoralis*, and key to the species of *Liparocephalus* (Coleoptera, Staphylinidae, Aleocharinae). Canadian Entomologist 132: 567–571. doi:10.4039/Ent132567-5
- Maruyama M, Hayashi M (2009) Description of the intertidal aleocharine *Halorhadinus sawadai* sp.n. from Japan, with notes on the genus *Halorhadinus* Sawada, 1971. Koleopterologische Rundschau 79: 71–82.
- Maruyama M, Klimaszewski J, Gusarov V (2008) *Osakatheta yasukoeae*, a new intertidal genus and species of athetine rove beetles (Coleoptera, Staphylinidae, Aleocharinae) from Japan. Zootaxa 1683: 39–50.
- Matsumura S (1911) Erster Beitrag zur Insekten-Fauna von Sachalin. Journal of the College of Agriculture, Tohoku Imperial University, Sapporo 4: 1–144.
- McCarthy D (2003) The trans-Pacific zipper effect: disjunct sister taxa and matching geological outlines that link the Pacific margins. Journal of Biogeography 30: 1545–1561. doi:10.1046/j.1365-2699.2003.00929.x
- McLachlan A, Brown AC (2006) The ecology of sandy shores, 2nd ed. Elsevier Science Publishers, Amsterdam, xii + 373 pp.
- Meyerdirk DE (1969) Adaptations of a rove beetle, *Diaulota fulviventris* Moore (Coleoptera: Staphylinidae) to the marine environment. MS Thesis, San Diego: San Diego State College.
- Mjöberg E (1906) Zur Kenntnis einiger unter Seetang lebenden Insekten. Zeitschrift für wissenschaftliche Insektenbiologie 2: 137–143.
- Moore I (1956a) A revision of the Pacific coast Phytosi with a review of the foreign genera (Coleoptera: Staphylinidae). Transactions of the San Diego Society of Natural History 12: 103–152.
- Moore I (1956b) Notes on some intertidal Coleoptera with descriptions of the early stages (Carabidae, Staphylinidae, Malachiidae). Transactions of the San Diego Society of Natural History 12: 207–230.
- Moore I (1964a) A new marine beetle from the Gulf of California. Transactions of the San Diego Society of Natural History 13: 173–176.

- Moore I (1964b) The larva of *Hadrotus crassus* (Mannerheim) (Coleoptera: Staphylinidae). Transactions of the San Diego Society of Natural History 13: 309–311.
- Moore I (1975) The larva of *Cafius sulcicollis* LeConte (Coleoptera: Staphylinidae). Pan-Pacific Entomologist 51: 140–142.
- Moore I (1976) *Giulianium campbelli*, a new genus and species of marine beetle from California (Coleoptera: Staphylinidae). Pan-Pacific Entomologist 52: 56–59.
- Moore I (1977) The larva of *Rothium sonorensis* Moore and Legner with a key to the known larvae of the genera of the marine Bolitocharini (Coleoptera: Staphylinidae). Psyche 84: 262–266.
- Moore I (1978a) Two new species of *Salinamexus* from western North America (Coleoptera: Staphylinidae). Entomological News 89: 113–115.
- Moore I (1978b) A second species of *Rothium*, an intertidal beetle from the Gulf of California. Pan-Pacific Entomologist 54: 155–156.
- Moore I, Legner EF (1971) *Bryothinusa chani*, a new species of marine beetle from Hong Kong (Coleoptera: Staphylinidae). Coleopterists Bulletin 25: 107–108.
- Moore I, Legner EF (1973) Succession of the coleopterous fauna in wrack. Wasmann Journal of Biology 31: 289–290.
- Moore I, Legner EF (1974a) Notes on *Bledius ornatus* LeConte, a seashore beetle, with description of the larva (Coleoptera: Staphylinidae). Wasmann Journal of Biology 32: 141–145.
- Moore I, Legner EF (1974b) The larva and pupa of *Carpelimus debilis* Casey (Coleoptera: Staphylinidae). Psyche 80: 289–294.
- Moore I, Legner EF (1975) A study of *Bryothinusa* comparing a tabular and a dichotomous key to the species. Bulletin of the Southern California Academy of Sciences 74: 109–112.
- Moore I, Legner EF (1976) Intertidal rove beetles (Coleoptera: Staphylinidae). In: Cheng L (Ed) Marine insects. North Holland, Amsterdam, 521–552.
- Moore I, Legner EF (1977) A report on some intertidal Staphylinidae from Sonora, Mexico with four new genera (Coleoptera). Pacific Insects 17: 459–471.
- Moore I, Legner EF, Chan T (1973) A review of the genus *Bryothinusa* with descriptions of three new species. Entomological News 84: 73–81.
- Moore I, Orth RE (1979a) *Diglossa legneri*, a new seashore beetle from California. Coleopterists Bulletin 33: 337–340.
- Moore I, Orth RE (1979b) Notes on *Bryothinusa* with description of the larva of *B. catalinae* Casey (Coleoptera: Staphylinidae). Psyche 85: 183–189.
- Morris RH, Abbott DP, Haderlie EC (Eds) (1980) Intertidal invertebrates of California. Stanford University Press, Stanford, ix + 290 pp.
- Motschulsky TV von (1854) Voyages. Lettres de M. de Motschulsky à M. Ménériés. Cairo sur le Mississippi [sic], le 26 novembre 1853. Etudes Entomologiques 3: 1–14.
- Motschulsky TV von (1858a) Enumération des nouvelles espèces de coléoptères rapportés de ses voyages. Bulletin de la Société Imperiale des Naturalistes de Moscou 31 (2): 634–670.
- Motschulsky TV von (1858b) Enumération des nouvelles espèces de coléoptères rapportés de ses voyages. Bulletin de la Société Imperiale des Naturalistes de Moscou 31 (3): 204–264.



- Motschulsky TV von (1860) Enumération des nouvelles espèces de coléoptères rapportés de ses voyages. 3-ième article. Bulletin de la Société Imperiale des Naturalistes de Moscou 33 (2): 539–588.
- Mulsant E, Rey C (1852) Description de quelques coléoptères nouveaux ou peu connus, de la tribu des brachélytres. Opusculs Entomologiques 1: 155–186, 189–190, pl. 1 [a reprint is paged 1–32].
- Mulsant E, Rey C (1853) Description de quelques coléoptères nouveaux ou peu connus, de la tribu des brachélytres. Opusculs Entomologiques 2: 35–85, pls. 1–2.
- Mulsant E, Rey C (1870) Description de quelques insectes nouveaux ou peu connus. Opusculs Entomologiques 14: 153–193.
- Mulsant E, Rey C (1873a) Histoire naturelle des coléoptères de France: brévipennes: aléochariens-suite. Deyrolle, Paris, 695 pp. + pl. 4.
- Mulsant E, Rey C (1873b) Description de divers coléoptères brévipennes nouveaux ou peu connus. Opusculs Entomologiques 15: 147–189.
- Mulsant E, Rey C (1874) Tribu des brévipennes. Famille des aléochariens. Septième branche: Myrmédoniaires. Annales de la Société d'Agriculture Histoire Naturelle et Arts Utiles de Lyon (sér. 4) 6 (1873): 33–738.
- Mulsant E, Rey C (1875) Histoire naturelle des coléoptères de France: brévipennes: aléochariens (suite)-Myrmédoniaires (2<sup>e</sup> partie). Deyrolle, Paris, 470 pp.
- Mulsant E, Rey C (1876) Tribu des brévipennes. Staphyliniens. Annales de la Société d'Agriculture Histoire Naturelle et Arts Utiles de Lyon (sér. 4) 8 (1875): 145–856, pls. i–vi.
- Mulsant E, Rey C (1878) Tribu des brévipennes. Cinquième famille, oxyporiens. Sixième famille, oxyteliens. Annales de la Société d'Agriculture Histoire Naturelle et Arts Utiles de Lyon (sér. 4) 10: 443–850, pls. i–vi.
- Mutchler AJ (1925) Coleoptera from the Williams Galapagos Expedition. Zoologica 5: 219–240.
- Nakane T (1963) New or little-known Coleoptera from Japan and its adjacent regions. XVII. Fragmenta Coleopterologica (1963) (5): 21–22.
- Newton AF, Chandler DS (1989) World catalog of the genera of Pselaphidae (Coleoptera). Fieldiana Zoology (n.s.) 53: i–iv, 1–93.
- Newton AF, Thayer MK (2007) Tree of Life Web Project. 1995. Staphylinidae Latreille 1802. Rove beetles. Version 01 January 1995 (temporary). <http://tolweb.org/Staphylinidae/9621/1995.01.01> in The Tree of Life Web Project, <http://tolweb.org/> [accessed 17.III.2010].
- Newton AF, Thayer MK, Ashe JS, Chandler DS (2001) Staphylinidae. In: Arnett RH, Jr, Thomas MC (Eds) American Beetles. CRC Press, Boca Raton, FL, Volume 1: 272–418.
- Nomura S, Nguyen TT, Nguyen TTA, Ahn KJ (2006) The first record of *Prosthecarthron sauteri* Raffray (Staphylinidae, Pselaphinae) from Vietnam. Elytra, Tokyo 34: 95–97.
- Notman H (1920) Staphylinidae from Florida in the collection of the American Museum, with descriptions of new genera and species. Bulletin of the American Museum of Natural History 42: 693–732.
- OED (1971) The compact edition of the Oxford English Dictionary. Complete text reproduced micrographically. Oxford University Press, Oxford.

- O'Keefe ST (2005) Scydmaenidae Leach, 1815. In: Beutel RG, Leschen RAB (Eds) Handbook of Zoology. Volume IV. Arthropoda: Insecta. Coleoptera Volume 1: Morphology and Systematics. Walter de Gruyter, Berlin, 280–288.
- Oldroyd H (1965) The natural history of flies. W. W. Norton and Co., New York, xiv + 324 pp.
- Orth RE, Moore I (1980) A revision of the species of *Cafius* Curtis from the west coast of North America with notes of the east coast species (Coleoptera: Staphylinidae). Transactions of the San Diego Society of Natural History 19: 181–211.
- Pace R (1984) Una nuova specie del genere *Bryothinusa* Casey delle Isole Samoa. Nouvelle Revue d'Entomologie (n.s.) 1: 67–70.
- Pace R (1985) Aleocharinae delle Comore raccolte da Yves Gomyi (Coleoptera, Staphylinidae). Revue Suisse de Zoologie 92: 621–629.
- Pace R (1986) Revisione di quattro specie dell'tribù Diglottini (Coleoptera, Staphylinidae) descritte da Fauvel. Annali del Museo Civico di Storia Naturale di Genova 80: 273–280.
- Pace R (1990) Aleocharinae delle Filippine. Mémoires du Muséum National d'Histoire Naturelle (Paris) série A, Zoologie 147: 57–113.
- Pace R (1992) Aleocharinae di Gibuti (Coleoptera, Staphylinidae). Bollettino del Museo di Storia Naturale di Verona 16: 180–186.
- Pace R (1993) Nuove Aleocharinae orientali (Coleoptera, Staphylinidae). Bollettino del Museo Civico di Storia Naturale di Verona 17: 127–180.
- Pace R (1994) Aleocharinae della sottoregione Africana Orientale al Museo di Ginevra (Coleoptera: Staphylinidae) Parte 1. Revue Suisse de Zoologie 101: 117–193.
- Pace R (1999a) Aleocharinae del Cile (Coleoptera, Staphylinidae). Bollettino del Museo Civico di Storia Naturale di Verona 23: 119–210.
- Pace R (1999b) Aleocharinae di Hong Kong (Coleoptera, Staphylinidae). Revue Suisse de Zoologie 106: 663–689.
- Pace R (2000a) Aleocharinae di Papua-Nuova Guinea (Coleoptera, Staphylinidae). Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Entomologie 70: 109–163.
- Pace R (2000b) Aleocharinae del Cile, nuove o poco note (Coleoptera, Staphylinidae). Bollettino del Museo Regionale di Scienze Naturali, Torino 17: 355–510.
- Pace R (2003) New or little known Aleocharinae from the Australian Region (Coleoptera, Staphylinidae). In: Daccordi M, Giachino PM (Eds) Results of the Zoological Missions to Australia of the Regional Museum of Natural Sciences of Turin, Italy. Museo Regionale di Scienze Naturali, Turin, I: 111–186.
- Pace R (2008) Aleocharinae (Insecta, Coleoptera, Staphylinidae) de Madagascar. Zoosystema 30: 555–577.
- Park JS, Jeon MJ, Ahn KJ (2008) Description of *Heterota sunjaei* sp. nov. from Korean sea-shores and an annotated catalog of the littoral genus *Heterota* (Coleoptera: Staphylinidae: Aleocharinae). Canadian Entomologist 140: 111–118. doi:10.4039/n07-015
- Paulian R (1938a) Les larves des espèces françaises du genre *Bledius* Mann. Bulletin de la Laboratoire Maritime de Dinard 19: 25–32.
- Paulian R (1938b) Etude biologique d'*Aleochara algarum* Fauvel. Bulletin de la Société Entomologique de France 63: 343–352.

- Paulian R (1941) Les premiers états des Staphylinidea (Coleoptera). Étude de morphologie comparée. Mémoires du Muséum National d'Histoire Naturelle (Nouvelle série) 15: 1–361 + pls. I–III.
- Paulian R (1942) Observations sur *Bledius spectabilis* Kraatz. Bulletin de la Laboratoire Maritime de Dinard 24: 62–72.
- Paulian R (1959) Observations sur la faune intercotidale de Madagascar. Naturaliste Malgache 11: 53–61.
- Peck SB (2005) A checklist of the beetles of Cuba with data on distributions and bionomics (Insecta: Coleoptera). Arthropods of Florida and Neighboring Land Areas 18: i–vi, 1–241.
- Peschke K, Fuldner D (1977) Übersicht und neue Untersuchungen zur Lebensweise der parasitoiden Aleocharinae (Coleoptera: Staphylinidae). Zoologische Jahrbücher für Systematik 104: 242–262.
- Peyron E (1857) Description de quelques coléoptères nouveaux et observations diverses. Annales de la Société Entomologique de France (sér. 3) 5: 715–723.
- Pope RD (1977) Kloet and Hincks. A check list of British insects. Second edition (completely revised). Part 3: Coleoptera and Strepsiptera. Handbooks for the Identification of British Insects 11 (3): i–xiv, 1–105.
- Raffray A (1890) Étude sur les psélaphides (Suite). V. Tableaux synoptiques. — Notes et synonymie. Revue d'Entomologie (Caen) 9: 81–172.
- Raffray A (1903) Voyage de M. L. Biró dans les possessions allemandes de la Nouvelle-Guinée. Annales Historico-Naturales Musei Nationalis Hungarici 1: 29–101, pls. 1–2.
- Raffray A (1915) H. Sauter's Formosa-Ausbeute. Pselaphidae (Col.) II. Supplementa Entomologica 3 (1914): 1–5.
- Rambousek FJ (1910) O bulharských Staphylinidech [Sur les staphylinides de Bulgarie]. Věstník Královské České Společnosti Náuk 4 (1909): 1–23. [note: the author provides a French translation interspersed with the Czech text].
- Redtenbacher L (1867) Reise der Österreichischen Fregatte Novara um die Erde in den Jahren 1857, 1858, 1859 unter des Befehl des Commodore B. von Wüllerstorff-Urbair. Zoologischer Theil. Zweiter Band. Coleopteren. Karl Gerold's Sohn, Vienna, iv + 249 pp. + 5 pls.
- Reitter E (1882) Versuch einer systematischen Eintheilung der Clavigeriden und Pselaphiden. Verhandlungen des Naturforschenden Vereines in Brünn 20: 177–211.
- Reitter E (1884) Beitrag zur Pselaphiden- und Scydmaeniden-Fauna von Java und Borneo. II. Stück. Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien 33 (1883): 387–428, pl. xx.
- Richards LJ (1982) Prey selection by an intertidal beetle: field test of an optimal diet model. Oecologia 55: 325–332. doi:10.1007/BF00376920
- Richards LJ (1983) Feeding and activity patterns of an intertidal beetle. Journal of Experimental Marine Biology and Ecology 73: 213–224. doi:10.1016/0022-0981(83)90048-5
- Richards LJ (1984) Field studies of foraging behaviour of an intertidal beetle. Ecological Entomology 9: 189–194. doi:10.1111/j.1365-2311.1984.tb00714.x
- Rottenberg A (1870) Beiträge zur Coleopteren-Fauna von Sicilien. Berliner Entomologische Zeitschrift 14: 11–40.

- Rougemont GD de (2001) The staphylinid beetles of Hong Kong. *Memoirs of the Hong Kong Natural History Society* 24: 1–146.
- Rye EC (1864) New British species, corrections, &c, noticed since the publication of the *Entomologist's Annual*, 1863. *Entomologist's Annual* (1864): 30–84.
- Sakaguti K (1944) A new intertidal rove-beetle from the Pacific coast of Japan. *Transactions of the Kansai Entomological Society* 14: 20–21, pls. i–ii.
- Saulcy F de (1864) Descriptions des espèces nouvelles de coléoptères recueillies en Syrie, en Egypte et en Palestine. *Annales de la Société Entomologique de France (sér. 4)* 4: 421–440, 629–660.
- Saunders LG (1928) Some marine insects of the Pacific coast of Canada. *Annals of the Entomological Society of America* 21: 521–545.
- Sawada K (1955) Marine insects of the Tokara Islands, VIII. Family Staphylinidae (Coleoptera). *Publications of the Seto Marine Biological Laboratory* 5: 81–87, pls. 14–15.
- Sawada K (1956) A new intertidal species of Staphylinidae from Ishigakajima, Ryukyu Islands (Coleoptera). *Kontyû* 24: 197–199.
- Sawada K (1971a) Aleocharinae (Staphylinidae, Coleoptera) from the campus of the Seto Marine Biological Laboratory. *Publications of the Seto Marine Biological Laboratory* 18: 291–315.
- Sawada K (1971b) Aleocharinae (Staphylinidae, Coleoptera) from the intertidal zone of Japan. *Publications of the Seto Marine Biological Laboratory* 19: 81–110.
- Sawada K (1971c) On the new genus and species of Leptotyphlinae of Japan (Coleoptera, Staphylinidae). *Revue d'Écologie et de Biologie du Sol* 8: 327–330.
- Sawada K (1985) *Atheta* and its allies of southeast Asia (Coleoptera: Staphylinidae) IV. Ten oriental species described by M. Bernhauer and M. Cameron. *Contributions of the Biological Laboratory of Kyoto University* 27: 91–110.
- Sawada K (1987) *Atheta* and its allies of southeast Asia (Coleoptera: Staphylinidae) V. Singaporean species described in Cameron, 1920. *Contributions of the Biological Laboratory of Kyoto University* 27: 137–150.
- Sawada K (1989a) On a new genus and new species of Aleocharinae (Coleoptera: Staphylinidae) from mangrove forests in Singapore. *Raffles Bulletin of Zoology* 37: 83–86.
- Sawada K (1989b) New species of Aleocharinae from Japan, I (Staphylinidae; Coleoptera). *Contributions of the Biological Laboratory of Kyoto University* 27: 273–307.
- Sawada K (1991) On new genera and species of intertidal Aleocharinae (Coleoptera: Staphylinidae) and Goniacerinae (Pselaphinae) from Singapore and Japan. *Raffles Bulletin of Zoology* 39: 141–152.
- Sawada K (1992) New genus and species of intertidal Brachyglutini (Coleoptera: Pselaphidae) from Japan. *Raffles Museum of Zoology* 40: 55–60.
- Say T (1834) Descriptions of new North American insects and observations on some already described. *Transactions of the American Philosophical Society* 4: 409–470. doi:10.2307/1004840
- Schaufuss LW (1880) Beschreibung sechzig neuer Pselaphiden [as appendix paged 5–25 in a presentation volume with the following dedication] *Der Société entomologique de Belgique zu Brüssel zur Feier Ihres fünfundzwanzigsten Stiftungstages die herzlichsten Festg-*

- rüsse aus dem Museum Ludwig Salvator in Oberblasewitz-Dresden. F. Thomass; Dresden. [note: a reprint of this article, with identical title, was later in 1880 published in *Nunquam Otiosus* (Dresden) 3: 481–511].
- Scheerpeltz O (1933) Staphylinidae VII. In: Schenkling S (Ed) *Coleopterorum Catalogus*. W. Junk, Berlin, pars 129: 989–1500.
- Scheerpeltz O (1934) Staphylinidae VIII. In: Schenkling S (Ed) *Coleopterorum Catalogus*. W. Junk, Berlin, pars 130: 1501–1881.
- Scheerpeltz O (1957a) Eine neue Art der Gattung *Cameronium* (Col. Staphylinidae). *Entomologische Blätter* 53: 147–150.
- Scheerpeltz O (1957b) Staphylinidae (Coleoptera) von Sumba und Flores. 4. Beitrag zur Kenntnis der Staphyliniden der orientalischen Region. *Verhandlungen der Naturforschenden Gesellschaft in Basel* 68: 217–357.
- Scheerpeltz O (1965) Eine neue Art der Gattung *Myrmecopora* Saulcy von den pazifischen Küste Nordamerikas (Col., Staph). *Deutsche Entomologische Zeitschrift* (N.F.) 12: 49–51.
- Scheerpeltz O (1969) Eine neue Art der Gattung *Pareiobledius* Bernhauer, nebst einer Übersicht über die bis heute bekannt gewordenen Arten dieser Gattung (Col. Staphylin.). *Koleopterologische Rundschau* 46/47: 127–130.
- Scheerpeltz O (1972) Studien an den paläarktischen arten der gattung *Myrmecopora* Saulcy (Col. Staphylinidae). *Koleopterologische Rundschau* 50: 93–109.
- Scher HD, Martin EE (2006) Timing and climatic consequences of the opening of Drake Passage. *Science* 312: 228–430. doi:10.1126/science.1120044
- Schiødte JMC (1845) Om slaegten *Micralymma*. *Naturhistorisk Tidsskrift* (ser. 2) 1: 370–379, pl. iv. [note: Jørgen Matthias Christian Schiødte was a Danish biologist; the spelling of his name as Schiödte is inappropriate because Danish does not have a letter ö].
- Schiødte JMC (1864) De metamorphosi eleutheratorum observationes. *Naturhistorisk Tidsskrift* (ser. 3) 3: 131–224, pls. i–xii.
- Schneider O (1898) Die Tierwelt der Nordsee-Insel Borkum unter Berücksichtigung der von den übrigen östfriesischen Inseln bekannten Arten. *Abhandlungen herausgegeben von Naturwissenschaftlicher Verein zu Bremen* 16: 1–174.
- Scriba W (1866) Ueber Staphylinen. *Berliner Entomologische Zeitschrift* 10: 289–290.
- Scotese CR (2010). PALEOMAP Project. <http://www.scotese.com/earth.htm> [accessed 20.XI.2010].
- Scott H (1916) Another staphylinid parasitic on a dipteran. *Entomologist's Monthly Magazine* 52: 206.
- Scott H (1920) Notes on (I) the parasitic staphylinid *Aleochara algarum* Fauvel, and its hosts, the phycodromid flies; (II) a case of supposed parasitism in the genus *Homalota*. *Entomologist's Monthly Magazine* 56: 148–157.
- Sharp D (1869) A revision of the British species of *Homalota*. *Transactions of the Entomological Society of London* (1869): 91–272.
- Sharp D (1870) Characters of a new genus and descriptions of new species of Aleocharidae from Britain. *Entomologist's Monthly Magazine* 6: 279–282.
- Sharp D (1871a) On a variety of *Philonthus xantholoma*. *Entomologist's Monthly Magazine* 7: 181.

- Sharp D (1871b) Description of a new genus and species of Staphylinidae (from South Australia). *Entomologist's Monthly Magazine* 7: 217–218.
- Sharp D (1874) The Staphylinidae of Japan. *Transactions of the Entomological Society of London* (1874): 1–103.
- Sharp D (1886) On New Zealand Coleoptera, with descriptions of new genera and species. *Scientific Transactions of the Royal Dublin Society* (ser. 2) 3: 351–454, pl. xii.
- Sharp D (1888) The Staphylinidae of Japan. *Annals and Magazine of Natural History* (ser. 6) 2: 277–295, 369–387, 451–464.
- Sharp D (1889) The Staphylinidae of Japan. *Annals and Magazine of Natural History* (ser. 6) 3: 28–44, 108–121, 249–267.
- Sharp D (1900) Some undescribed species of *Trogophloeus*, with a new genus. *Entomologist's Monthly Magazine* 36: 230–234.
- Sherwood BR, Gardiner BG, Harris T (2000) British saltmarshes. Joint symposium on British Saltmarshes organized between the Linnean Society of London, the Royal Society for the Protection of Birds, and English Nature (London; 2000). Forrester Text, Cardigan, Wales, xvi + 417 pp.
- Smetana A (1971) Revision of the tribe Quediini of America north of Mexico (Coleoptera: Staphylinidae). *Memoirs of the Entomological Society of Canada* 79: i–vi, 1–303.
- Smetana A (2009) BOOK REVIEW. Majka CG, Klimaszewski J (2008) Biodiversity, bio-systematics, and ecology of Canadian Coleoptera. *Coleopterists Bulletin* 63: 380–382. doi:10.1649/1194.1
- Solier AJJ (1849) Estafilinoideos. In: Gay C, *Historia física y política de Chile según documentos adquiridos en esta república durante doce años de residencia en ella y publicada bajo los auspicios del supremo gobierno*. Zoología. Gay, Paris, 4: 302–356.
- Steel WO (1949) A new genus and species of Pacific (Chatham Is.) Staphylinidae. *Entomologist's Monthly Magazine* 85: 309–310.
- Steel WO (1958) Notes on the Omaliinae (Col., Staphylinidae). (9). The genus *Micralymma* Westwood. *Entomologist's Monthly Magazine* 94: 138–142.
- Steel WO (1962) Notes on the Omaliinae (Col., Staphylinidae). (11). The genus *Micromalium* Melichar, with further notes on *Micralymma* Westwood. *Entomologist's Monthly Magazine* 97(1961): 237–239.
- Steel WO (1964) Insects of Campbell Island. Coleoptera: Staphylinidae. *Pacific Insects Monograph* 7: 340–375.
- Steel WO (1970) The larvae of the genera of Omaliinae (Coleoptera: Staphylinidae) with particular reference to the British fauna. *Transactions of the Royal Entomological Society of London* 122: 1–47. doi:10.1111/j.1365-2311.1970.tb00524.x
- Stephens JF (1829) The nomenclature of British insects; being a compendious list of such species as are contained in the systematic catalogue of British insects, and forming a guide to their classification. Baldwin and Cradock, London, 68 pp.
- Stephens JF (1832) Illustrations of British entomology: or, a synopsis of indigenous insects: containing their generic and specific distinctions; with an account of their metamorphoses, times of appearance, localities, food and economy, as far as practicable. Mandibulata. Baldwin and Cradock, London, Vol. 5, 240 pp. + pls. 24–26.



- Stephens JF (1833) Illustrations of British entomology: or, a synopsis of indigenous insects: containing their generic and specific distinctions; with an account of their metamorphoses, times of appearance, localities, food and economy, as far as practicable. Mandibulata. Baldwin and Cradock, London, Vol. 5: 241–304 + pl. 27.
- Strand E (1935) Revision von Gattungsnamen palaearktischer Coleoptera. Folia Zoologica et Hydrobiologica 7: 282–299.
- Strøm H (1783) Norske insecters beskrivelse med anmaerkninger. Nye Samling af det Kongelige Danske Videnskabers Selskabs Skrifter 2: 49–93. [note: Hans Strøm was Norwegian, and in Norwegian there is no letter ö, so printing of the name as Ström is inappropriate; see Apelseth, A. 1997. Bibliografi over Hans Strøm <http://www2.hivolda.no/ahf/hansstrom/bibliografi.htm>].
- Tanokuchi Y (1989) Some Pselaphidae inhabiting the mangrove forests of Singapore and Thailand, with description of a new genus and eight new species. Raffles Bulletin of Zoology 37: 87–115.
- Thayer MK (1985) *Micralymma marinum* (Stroem) in North America: biological notes and new distributional records (Coleoptera: Staphylinidae). Psyche 92: 49–55.
- Thayer MK (2005) Staphylinidae Latreille, 1802. In: Beutel RG, Leschen RAB (Eds) Handbook of Zoology. Volume IV. Arthropoda: Insecta. Coleoptera Volume 1: Morphology and Systematics. Walter de Gruyter, Berlin, 296–344.
- Thomson CG (1852) Öfversigt af de i Sveriges funna arter af släktet *Homalota* (Mannerheim). Öfversigt af Kongliga Svenska Vetenskapsakademiens Förhandlingar 9: 131–149.
- Thomson CG (1857) Arter af släktet *Omalium*, funna i Sverige. Öfversigt af Kongliga Vetenskapsakademiens Förhandlingar 13 (1856): 223–238.
- Thomson CG (1858) Försök till uppställning af sveriges Staphyliner. Öfversigt af Kongliga Svenska Vetenskapsakademiens Förhandlingar 15: 27–40.
- Thomson CG (1859) Skandinaviens Coleoptera, synoptisk bearbetade. Tom. I. Berlinoska, Lund, 290 pp.
- Thomson CG (1861) Skandinaviens Coleoptera, synoptisk bearbetade. Tom. III. Berlinoska, Lund, 278 pp.
- Topp W, Ring RA (1988a) Adaptations of Coleoptera to the marine environment. I. Observations on rove beetles (Staphylinidae) from sandy beaches. Canadian Journal of Zoology 66: 2464–2468. doi:10.1139/z88-365
- Topp W, Ring RA (1988b) Adaptations of Coleoptera to the marine environment. II. Observations on rove beetles (Staphylinidae) from rocky shores. Canadian Journal of Zoology 66: 2469–2474. doi:10.1139/z88-366
- Tottenham CE (1949) New generic names in Staphylinidae (Coleoptera). Proceedings of the Royal Entomological Society of London (B) 18: 78. doi:10.1111/j.1365-3113.1949.tb01425.x
- Tottenham CE (1955) Studies in the genus *Philonthus* Stephens (Coleoptera: Staphylinidae) Parts II, II, and IV. Transactions of the Royal Entomological Society of London 106: 153–195. doi:10.1111/j.1365-2311.1955.tb00783.x
- Tronquet M (2003) *Arena tabida* (Kiesenwetter, 1850) et *Pseudopasilia testacea* (Brisout, 1863) ou la confusion des genres (Coleoptera, Staphylinidae). Revue de l'Association Rousillonnaise d'Entomologie 12: 19–26.

- Vauloger de Beaupré M (1897) Description de deux coléoptères nouveaux du nord de l'Afrique. Bulletin de la Société Entomologique de France (1897): 237–239.
- Wagner H (1933) In: Neresheimer J, Wagner H, Beiträge zur Coleopterenfauna der Mark Brandenburg. Entomologische Blätter für Biologie und Systematik der Käfer 31: 106–120.
- Walker F (1858) Characters of some apparently undescribed Ceylon insects. Annals and Magazine of Natural History (ser. 3) 2: 202–209.
- Waterhouse CO (1875) On the Coleoptera of Kerguelen's Island. Entomologist's Monthly Magazine 12: 54–57.
- Waterhouse CO (1879) Zoology of Kerguelen Island: Coleoptera. Philosophical Transactions of the Royal Society of London 168: 230–234, pl. xiv.
- Waterhouse FH (1879) Descriptions of new Coleoptera of geographical interest, collected by Charles Darwin, Esq. Journal of the Linnean Society of London, Zoology 14: 530–534. doi:10.1098/rstl.1879.0021
- Waterhouse GR (1858) Description of four new species of Staphylinidae. Zoologist 16: 6073–6075.
- Waterhouse GR (1859) [Description of four new species of Staphylinidae]. Proceedings of the Entomological Society of London (1859): 14–17. [note: although the meeting was held in February 1858, publication seems not to have been until 1859, by which time the descriptions had been published in The Zoologist, cited above].
- Waterhouse GR (1863) [Some British species of *Homalota*]. Proceedings of the Entomological Society of London (1863): 136–140.
- Weise J (1877) Japanische Staphylinidae [sic] und Pselaphidae In: Beiträge zur Käferfauna von Japan, meist auf R. Hiller's Sammlungen basiert. Deutsche Entomologische Zeitschrift 21: 88–100.
- Wendeler H (1930) Neue exotische Staphyliniden (Coleoptera). Neue Beiträge zur Systematischen Insektenkunde 4: 248–252.
- Westwood JO (1838) Notes upon subaquatic insects, with the description of a new genus of British Staphylinidae. Magazine of Zoology and Botany 2: 124–132, pl. 4.
- White A (1846) Insects of New Zealand. Pt. 11. In: Richardson J, Gray JE (Eds) The zoology of the voyage of H.M.S. Erebus and Terror, under the command of Captain Sir James Clark Ross, R.N., F.R.S., during the years 1839–1843. Longman, Brown, Green and Longman, London, 1–24.
- Wikipedia (2010) Sea of Japan naming dispute. Seen November 2010. Search for Wiki+“Sea of Japan Naming dispute”
- Wollaston TV (1860) On additions to the Madeiran Coleoptera. Annals and Magazine of Natural History (ser. 3) 6: 48–54.
- Wollaston TV (1864) Catalogue of the coleopterous insects of the Canaries in the collection of the British Museum. British Museum (Natural History), London, xii + 648 pp.
- Wollaston TV (1865) Coleoptera Atlantidum, being an enumeration of the coleopterous insects of the Madeiras, Salvages, and Canaries. Van Voorst, London, xlvii + 526 pp. + 1 map + appendix (140 pp.).

- Wong MH, Chan TD (1977) The ecology of the marine rove beetle, *Bryothinusa* spp. (Coleoptera: Staphylinidae) in Hong Kong. *Hydrobiologia* 53: 253–256. doi:10.1007/BF00818547
- Wyatt TD (1982) The salt marsh beetle. *Antenna* 6: 303–304.
- Wyatt TD (1986) How a subsocial intertidal beetle, *Bledius spectabilis*, prevents flooding and anoxia in its burrow. *Behavioral Ecology and Sociobiology* 19: 323–331. doi:10.1007/BF00295705
- Wyatt TD, Foster WA (1988) Distribution and abundance of the intertidal saltmarsh beetle *Bledius spectabilis*. *Ecological Entomology* 13: 453–464. doi:10.1111/j.1365-2311.1988.tb00378.x
- Yosii R, Sawada K (1976) Studies on the genus *Atheta* Thomson and its allies (Coleoptera, Staphylinidae). II. Diagnostic characters of genera and subgenera with description of representative species. *Contributions from the Biological Laboratory, Kyoto University* 25: 11–140.
- Zanetti A (1987) Coleoptera. Staphylinidae. Omaliinae. In: *Fauna d'Italia* 25. Calderini, Bologna, xii + 472 pp.
- Zerche L (1998) *Amblopusa magna* sp. n.—eine neue Art der amphipazifischen Gattung *Amblopusa* Casey, 1893 aus dem Fernen Osten Russlands und ihre Stellung im phylogenetischen System (Coleoptera: Staphylinidae, Aleocharinae, Liparocephalini). *Beiträge zur Entomologie* 48: 103–113.