

A review of *Aleurodaphis* (Hemiptera, Aphididae, Hormaphidinae) with the description of one new species and keys to species

Li-Yun Jiang[†], Ge-Xia Qiao[‡]

Key Laboratory of Zoological Systematics and Evolution and National Zoological Museum of China, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, P. R. China

[†] [urn:lsid:zoobank.org:author:3386E41A-0B18-4558-97D0-F96E2489F5D5](https://doi.org/urn:lsid:zoobank.org:author:3386E41A-0B18-4558-97D0-F96E2489F5D5)

[‡] [urn:lsid:zoobank.org:author:6075A192-E433-4782-9F4D-013126A42DC1](https://doi.org/urn:lsid:zoobank.org:author:6075A192-E433-4782-9F4D-013126A42DC1)

Corresponding author: Ge-Xia Qiao (qiaogx@ioz.ac.cn)

Academic editor: Mike Wilson | Received 17 June 2011 | Accepted 16 August 2011 | Published 7 October 2011

[urn:lsid:zoobank.org:pub:A8E2B269-844B-4CE7-ACE2-AE72D79F48F5](https://doi.org/urn:lsid:zoobank.org:pub:A8E2B269-844B-4CE7-ACE2-AE72D79F48F5)

Citation: Jiang L-Y, Qiao G-X(2011) A review of *Aleurodaphis* (Hemiptera, Aphididae, Hormaphidinae) with the description of one new species and keys to species. ZooKeys 135: 41–56. doi: 10.3897/zookeys.135.1721

Abstract

The genus *Aleurodaphis* van der Goot is reviewed. One new species *Aleurodaphis sinojackiae* Qiao & Jiang, sp. n. on *Sinojackia xylocarpa* from Jiangsu and Zhejiang, China is described. *Aleurodaphis sinisalicis* Zhang, 1982 is synonymised with *A. blumeae* van der Goot, 1917. Keys to species, morphological description and features of the new species, host plants, and distribution are provided. The specimens including types are deposited in British Natural History Museum, London (BMNH), Kōgakkan University, Japan and the National Zoological Museum of China, Institute of Zoology, Chinese Academy of Sciences, Beijing, China (NZMC).

Keywords

Hormaphidinae, *Aleurodaphis*, new species, synonym

Introduction

Aleurodaphis is erected in 1917 by van der Goot. He described the species *Aleurodaphis blumeae* as the type of the genus, from *Blumea*. Its remarkable characters are the followings, body of apterae is aleurodiform, frontal horn is absent and wax glands are arranged along crenulated margin of body. Takahashi studied the specimens from East

Asia, and described two species *A. mikaniae* Takahashi, 1925 and *A. asteris* Takahashi & Sorin, 1958. More than 30 years later, one new species was found from India, *A. antennata* Chakrabarti & Maity, (1980) 1982 and one new species was reported in China, *A. sinisalicis* Zhang, 1982. Sorin and Miyazaki (2004) reviewed the genus from Japan with descriptions of three new species, *A. impatientis*, *A. ligulariae* and *A. stewartiae*. After identifying the specimens from China and checking the specimens of the genus in British Natural History Museum, one new species, *A. sinojackiae* Qiao & Jiang, sp. n. is found, and *A. sinisalicis* Zhang, 1982 is synonymised with *A. blumae* van der Goot, 1917 here. Therefore, the genus has eight known species in the world (Remaudière and Remaudière 1997; Sorin and Miyazaki 2004), including the new species described here.

Material and methods

Aphid terminology in this paper follows Sorin and Miyazaki (2004) and Ghosh (1988). The unit of measurements in this paper is millimeters (mm).

In Tables 1-2, the following abbreviations have been used: Ant. I – IV = antennal segments I – IV; Ant. V b = base of antennal segment V; pt = processus terminalis; URS = ultimate rostral segment; Hind T & F = hind trochanter & femur; 2HTs = second hind tarsal segment.

Specimen depositories. The holotype and some paratypes of the new species are deposited in British Natural History Museum, London (BMNH), while the other paratypes in the National Zoological Museum of China, Institute of Zoology, Chinese Academy of Sciences, Beijing, China (NZMC) and Kôgakkan University, Japan. All the other specimens studied are deposited in BMNH and NZMC.

Systematics

Aleurodaphis van der Goot, 1917

<http://species-id.net/wiki/Aleurodaphis>

Aleurodaphis van der Goot, 1917: 239.

Aleurodaphis van der Goot: Baker, 1929: 86; Takahashi, 1931: 92; Takahashi and Sorin, 1958: 31; Raychaudhuri, Ghosh, Pal and Ghosh, 1980: 36; Ghosh, 1988: 249; Noordam, 1991: 47; Tao, 1990: 58; Blackman and Eastop, 1994: 551; Remaudière and Remaudière, 1997: 181; Tao, 1999: 17.

Type species. *Aleurodaphis blumae* van der Goot, 1917.

Diagnosis. Body oval and flat. In apterous females: body aleyrodiform, absence of frontal horns, and wax glands arranged along the crenulated margin of body. Head and prothorax, meso- and metathorax, abdominal tergites I–VII fused, respectively; only abdominal tergite VIII free; antennae 4 or 5-segmented, primary rhinaria

small and ciliated; eyes with 3 facets. Dorsal setae fine and sparse. Rostrum reaching mid-coxae, at most hind coxae. Ultimate rostral segment obviously longer than second hind tarsal segment. Legs short; first tarsal chaetotaxy: 2–4, 2–4, 2–4; dorsal-apical setae on second hind tarsal segments with funnel-shaped apex. Siphunculi ring-shaped. Cauda knobbed and anal plate bilobed. In alate viviparous females: antennae 5-segmented, with secondary rhinaria near ring-shaped, without cilia; eyes normal; first tarsal chaetotaxy: 4, 4, 4, sometimes 3 or 2; fore wings with media once branched, pterostigma extended and two cubitus fused or separated at base; hind wings with two obliques.

Host plants. The range of host plants in *Aleurodaphis* is quite wide, including Compositae (*Aster*, *Blumea*, *Carpesium*, *Chrysanthemum*, *Kalimeris*, *Ligularia*, *Parasenecio*, *Senecio*), Balsaminaceae (*Impatiens*), Gramineae (*Bambusa*), Moraceae (*Ficus*), Plantaginaceae (*Plantago*), Scrophulariaceae (*Mazus*), Styrcaceae (*Sinojackia*), Theaceae (*Stewartia*), Verbenaceae (*Callicarpa*) and Violaceae.

Biology. Five species, *A. asteris*, *A. blumeae*, *A. impatientis*, *A. ligulariae* and *A. mikaniae*, mainly feeding on Compositae species, have monoecious and anholocyclic life cycle. *A. sinojackiae* Qiao & Jiang, sp. n. and *A. stewartiae* can form galls on the leaves of the primary host plants, but their secondary hosts are unknown. The details of *A. antennata* were unreported (Ghosh 1988; Blackman and Eastop 1994, 2006; Sorin and Miyazaki 2004).

Distribution. China, Japan, India and Indonesia.

Keys to species of *Aleurodaphis*

Apterous viviparous females

- 1 Body without marginal wax glands; on *Stewartia*, in curled leaves..... *A. stewartiae*
- Body with marginal wax glands 2
- 2 Marginal wax glands arranged in each segment, not connecting with each other (Fig. 20); on *Sinojackia*, in curled leaves *A. sinojackiae* Qiao & Jiang, sp. n.
- Marginal wax glands arranged consecutively along the crenulated margin of body (Figs 1, 2), not in curled leaves..... 3
- 3 Ultimate rostral segment slender and long, 4.60–5.67 times as long as its basal width (Fig. 2) *A. blumeae*
- Ultimate rostral segment stout and short, less than 3.30 times as long as its basal width..... 4
- 4 Each first tarsal segment with 2 setae; marginal wax glands along the crenulated margin of body with 120 wax facets at most..... *A. asteris*
- Each first tarsal segment with more than 2 setae; marginal wax glands along the crenulated margin of body with 150 wax facets at least 5
- 5 Ultimate rostral segment 1.16–1.40 times as long as second hind tarsal segment 6

- Ultimate rostral segment 1.41–2.02 times as long as second hind tarsal segment 7
- 6 Ultimate rostral segment 1.16–1.30 times as long as second hind tarsal segment; dorsal of body without obvious mastoid process; on *Bambusa*.....
..... *A. antennata*
- Ultimate rostral segment 1.40 times as long as second hind tarsal segment; dorsal of body with obvious mastoid process; on Compositae and Balsaminaceae (Impatiens) *A. mikaniae*
- 7 First tarsal chaetotaxy: 3 or 4, 3 or 4, 3 or 4; triommatidia elongate, with the outer-most facet placed widely apart from the other two; on *Impatiens*
..... *A. impatientis*
- First tarsal chaetotaxy: 2 or 3, 2 or 3, 2; triommatidia thickset, with the outer-most facet placed close to the other two; on *Ligularia* *A. ligulariae*

Alate viviparous females

- 1 Antennae 4-segmented, with more than 30 setae on each of segments III and IV *A. ligulariae*
- Antennae 5-segmented, with 2–5 setae on each of segments III and IV 2
- 2 Ultimate rostral segment less than 3.00 times as long as its basal width..... 3
- Ultimate rostral segment more than 3.00 times as long as its basal width.... 4
- 3 Antennal segment III with 10–14 secondary rhinaria; first tarsal chaetotaxy: 4, 4, 4, sometimes 3, 3, 3..... *A. sinojackiae* Qiao & Jiang, sp. n.
- Antennal segment III with 24–27 secondary rhinaria; first tarsal chaetotaxy: 3, 3, 3, sometimes 2, 2, 2..... *A. mikaniae*
- 4 Ultimate rostral segment 4.60–7.50 times as long as its basal width, and 1.96–2.32 times as long as second hind tarsal segment *A. blumeae*
- Ultimate rostral segment 3.38–4.00 times as long as its basal width, and 1.19–1.80 times as long as second hind tarsal segment 5
- 5 First tarsal chaetotaxy: 3, 3, 3; antennal segment III with 25–29 secondary rhinaria..... *A. stewartiae*
- First tarsal chaetotaxy: 4, 4, 4 or 3–4, 3–4, 2–4; antennal segment III with 9–12 secondary rhinaria..... *A. impatientis*

Aleurodaphis antennata Chakrabarti & Maity, (1980) 1982

http://species-id.net/wiki/Aleurodaphis_antennata

Aleurodaphis antennata Chakrabarti & Maity, (1980) 1982: 56.

Aleurodaphis antennata Chakrabarti & Maity: Ghosh, 1988: 252; Blackman and Eastop, 1994: 551; Remaudière and Remaudière, 1997: 179.

Host plants. *Bambusa* sp.

Distribution. India (Ghosh 1988) .

***Aleurodaphis asteris* Takahashi & Sorin, 1958**

http://species-id.net/wiki/Aleurodaphis_asteris

Aleurodaphis asteris Takahashi & Sorin, 1958: 31.

Aleurodaphis asteris Takahashi & Sorin: Zhang, Zhong and Zhang, 1992: 142; Remaudière and Remaudière, 1997: 179; Tao, 1999: 17; Sorin and Miyazaki, 2004: 166.

Material examined. **CHINA** (NZMC): 2 apterous viviparous females, 15 April 1991, Jiangle, Fujiang, No. 10054, on Violaceae, coll. W. Y. Zhang; 8 apterous viviparous females, 13 August 2003, Motuo, Tibet, No. 15371, host plants unknown, coll. G. X. Qiao and X. L. Huang; **JAPAN** (BMNH): 24 apterous viviparous females, 5 August 1966, Osaka, Chihaya, on *Aster* sp., coll. M. Sorin.; 9 apterous viviparous females, 29 May 1964, Osaka, Kongo Mt., on *Aster* sp., coll. v. d. Bosch; 4 apterous viviparous females, 7 June 1966, Kyushu, Hikosan, on *Kalimeris* sp., coll. H. Takada; 8 apterous viviparous females, 6 August 1980, Kyoto, Kibune Mt., on *Aster yomena*, coll. R. L. Blackman; **KOREA** (BMNH): 2 apterous viviparous females, 15 September 1963, Ul-nungdo, on *Aster incisus*, coll. W. H. Paik; 1 apterous viviparous female, 14 September 1963, Pusan, on *Chrysanthemum zawidskii*, coll. W. H. Paik.

Host plants. *Carpesium abrotanoides*, *Aster yomena*, *A. incisus*, *Chrysanthemum zawidskii*, *Kalimeris* sp. and Violaceae.

Biology. The species feed on the stems, leafstalks, flower stalks and leaves of the host plants.

Distribution. China, Japan and Korea.

***Aleurodaphis blumeae* van der Goot, 1917**

http://species-id.net/wiki/Aleurodaphis_blumeae

Figs 1–2

Aleurodaphis blumeae van der Goot, 1917: 240.


Aleurodaphis nobukii Shinji, 1923: 301.

Astegopteryx japonica Takahashi, 1923: 150.

Aleurodaphis sinisalicis Zhang, 1982: 20. syn. n.

Aleurodaphis blumeae van der Goot: Takahashi, 1921: 92; Takahashi, 1923: 150; Takahashi, 1924: 98; Raychaudhuri, Ghosh, Pal and Ghosh, 1980: 362; Ghosh, 1988: 256; Noordam, 1991: 47; Tao, 1990: 59; 1999: 17; Remaudière and Remaudière, 1997: 179; Sorin and Miyazaki, 2004: 166.

Comments. The type specimens of *Aleurodaphis sinisalicis* Zhang, 1982 were checked, including 48 apterous viviparous females, 25 July 1963, Sichuan (Guanxian County), No. Y0399, on *Salix* sp., coll. G. X. Zhang and T. S. Zhong. The result confirmed the queries of Blackman and Eastop (1994) and Remaudière and Remaudière (1997) that *A. sinisalicis* (Fig. 2) was a synonym of *A. blumeae* (Fig. 1).



Figures 1–2. Apterous viviparous female. **1** dorsal view of body, *Aleurodaphis blumeae* van der Goot **2** dorsal view of body, syntype of *Aleurodaphis sinisalicis* Zhang. Scale bars = 0.10 mm.

The original descriptions of *Aleurodaphis sinisalicis* Zhang, 1982 were accurate, but the morphological characters of *A. blumeae* in his diagnosis were wrong. Perhaps, this is the main reason why Zhang (1992) described it as a new species. In the original descriptions of *Aleurodaphis sinisalicis* Zhang, 1982, the diagnosis was: the ratio of body length to antennae length was 4.70 (*A. blumeae*: 2.70), the base of cauda restricted (*A. blumeae*: not restricted), and the anal plate bilobed (*A. blumeae*: not bilobed). Actually, the morphological characters of *A. blumeae* in this diagnosis were inaccurate. In *A. blumeae*, the ratio of body length to antennae length was 4.80 instead of 2.70, the base of cauda restricted instead of not restricted, and the anal plate bilobed instead of not bilobed.

The host plant of *A. sinisalicis*, *Salix* sp., is perhaps mis-recorded.

Material examined. CHINA (NZMC): 6 apterous viviparous females, 17 August 2004, Guizhou (Daozhen County), No. 15597, host plants unknown, coll. J. Y. Yang; 6 apterous viviparous females, 17 July 2001, Shaanxi (Nanzheng County), No. Y8606, host plants unknown, coll. S. H. Wang; 7 apterous viviparous females, 8 September 1995, Jiangxi (Jinggangshan City), No. 10852, on Compositae, coll. G. X. Zhang; 9 apterous viviparous females and 7 alate viviparous females, 25 April 1984, Shaanxi (Yangling County), No. 64, on *Carpesium cernuum*, coll. X. F. Dai; 5 alate viviparous females, May 1984, Shaanxi (Yangling County), No. Y6227, host plants unknown, coll. X. F. Dai; 3 apterous viviparous females, 26 June 1983, Zhejiang (Lin'an City),

No. Y2692, on *Carpesium abrotanoides*, collector unknown; 5 apterous viviparous females, 8 April 1998, Guangxi (Napo County), No. 11772, on *Callicarpa bodinieri*, coll. G. X. Qiao; 14 apterous viviparous females, 21 April 1998, Guangxi (Fangchenggang City), No. 11840, on *Senecio scandens*, coll. G. X. Qiao; 16 apterous viviparous females, 22 March 1998, Guangxi (Pingxiang City), No. 11580, on *Plantago asiatica*, coll. G. X. Qiao; 7 apterous viviparous females, Hunan, No. 8887, on Compositae, the collector unknown; 3 apterous viviparous females and 2 nymphs, February 1925, Taiwan (Taihoku), on *Ficus* sp., coll. R. Takahashi (BMNH); 107 apterous viviparous females, 1 alate viviparous female and 24 nymphs, 21 May 1985, Zhejiang (Hangzhou City), on *Carpesium abrotanoides*, coll. V. F. Eastop (BMNH); **JAPAN** (BMNH): 2 apterous viviparous females and 4 nymphs, 29 August 1913, Kumamoto, on *Blumea* sp., coll. Theobald; 10 alate viviparous females, 22 September 1957, Osaka, on *Carpesium abrotanoides* var. *tumbergianum*, coll. M. Sorin; 2 apterous viviparous females, 2 alate viviparous females and 4 nymphs, 30 July 1957, Tokyo, Takao Mt., on *Blumea* sp., coll. R. Takahashi; 7 apterous viviparous females and 12 nymphs, 16 August 1991, Chiba, Sayama, on *Carpesium* sp., coll. D. L. Stern; **KOREA** (BMNH): 2 apterous viviparous females, 15 September 1963, Ulnungdo, on *Mazus miguelii*, coll. W. H. Paik; 1 apterous viviparous female, July 1969, Lri, host plants unknown, coll. W. H. Paik; **INDONESIA** (BMNH): 6 apterous viviparous females and 2 alate viviparous females, 13 July 1916, Garoet, on Compositae, coll. D. van der Goot; **MALAYSIA** (BMNH): 3 apterous viviparous females and 2 nymphs, 23 September 1944, Cameron Highlands, on *Blumea* sp., coll. R. Takahashi; **PHILIPPINES** (BMNH): 1 alate viviparous female, September 1962, Davao Exp. Station, trap in Abacca grove, coll. M. R. Gavarra; 1 alate viviparous female, July 1963, Davao Exp. Station, host plants unknown, coll. M. R. Gavarra; 1 alate viviparous female, January 1964, Davao Exp. Station, host plants unknown, coll. M. R. Gavarra; 1 alate viviparous female, March 1964, Davao Exp. Station, host plants unknown, coll. M. R. Gavarra; 3 apterous viviparous females and 2 nymphs, 13 September 1964, Makiling, on *Blumea* sp., coll. V. S. Calilung.

Host plants. *Carpesium cernuum*, *C. abrotanoides*, *C. abrotanoides* var. *tumbergianum*, *Senecio scandens*, *Blumea chinensis*, *Callicarpa bodinieri*, *Mazus miguelii*, *Ficus* sp. and *Plantago asiatica*. The common hosts are various Compositae.

Biology. This species feeds on the lower surface of leaves, along the main veins. It can infest *Blumea* on stems and undersides of young leaves, causing slight leaf-curl (Calilung 1967).

Distribution. China, Japan, Korea, Indonesia, Malaysia and Philippine.

Aleurodaphis impatientis Sorin & Miyazaki, 2004

http://species-id.net/wiki/Aleurodaphis_impatientis

Aleurodaphis impatientis Sorin & Miyazaki, 2004: 167.

Host plants. *Impatiens textori* and *I. noli-tangere*.

Biology. The species is viviparous throughout the year on *Impatiens* spp. Alate viviparous females appear in the latter part of September. Adult apterous viviparous females pass the winter on the stalks near and just below the ground level. The hibernated adult females move to the seedlings and start to feed on the leaves and stalks in mid-April (Sorin and Miyazaki 2004).

Distribution. Japan (Sorin and Miyazaki 2004).

***Aleurodaphis ligulariae* Sorin & Miyazaki, 2004**

http://species-id.net/wiki/Aleurodaphis_ligulariae

Aleurodaphis ligulariae Sorin & Miyazaki, 2004: 170.

Host plant. *Ligularia fischeri*.

Biology. The aphid lives on the lower side of the leaves and the apical part of the stem, as well as on the flower stalk at the tips of the host plant shoots. The alate viviparous females appear in the latter part of September. Adult apterous viviparous females pass the winter on the basal part of the stem just below the ground level and on fallen leaves in the ground litter (Sorin and Miyazaki 2004).

Distribution. Japan (Sorin and Miyazaki 2004).

***Aleurodaphis mikaniae* Takahashi, 1925**

http://species-id.net/wiki/Aleurodaphis_mikaniae

Aleurodaphis mikaniae Takahashi, 1925: 51.

Aleurodaphis mikaniae Takahashi: Remaudière and Remaudière, 1997: 179; Tao, 1999: 17.

Material examined. CHINA (NZMC): 6 apterous viviparous females and 3 alate viviparous females, 24 August 2004, Guizhou (Daozhen County), No. 15638, host plants unknown, coll. J. Y. Yang; 4 apterous viviparous females, 15 August 2003, Sichuan (Baoxing County), No. 15017, on *Parasenecio* sp., coll. K. Guo; 7 apterous viviparous females, 27 June 1999, Shaanxi (Foping County), No. 12336, on Compositae, coll. T. L. He; 20 apterous viviparous females, 12 October 1988, Hunan (Zhangjiajie City), No. 8962, on *Impatiens* sp., coll. T. S. Zhong and G. X. Zhang; 4 apterous viviparous females, 26 September 1974, Guizhou (Guiyang City), No. Y2123, on *Senecio scandens*, coll. Y. Y. Rao; 10 apterous viviparous females, 31 March 1982, Yunnan (Kunming City), No. 7373, on *Senecio scandens*, coll. G. X. Zhang; 6 apterous viviparous females, 12 October 1996, Shaanxi (Zhouzhi County), No. 11096, host plants unknown, coll. G. X. Qiao; 4 apterous viviparous females, 18 August 2003, Sichuan (Baoxing County), host plants unknown, coll. K. Guo; 10 apterous viviparous females, 12 July 2002, Shaanxi (Meixian County), No. 13559, host plants unknown, coll. E. B. Ma; 14 apterous viviparous females, July 1936, Taiwan (Shinkwan),

host plants unknown, coll. R. Takahashi; **JAPAN** (BMNH): 41 apterous viviparous females, 6 August 1980, Kyoto, Kibune Mt., on *Impatiens* sp., coll. R. L. Blackman.

Host plants. *Parasenecio* sp., *Impatiens* sp., and *Senecio scandens*.

Distribution. China and Japan.

***Aleurodaphis sinojackiae* Qiao & Jiang, sp. n.**

urn:lsid:zoobank.org:act:4FE949A7-BF92-425C-BEDC-176EB4CA495A


http://species-id.net/wiki/Aleurodaphis_sinojackiae

Figs 3–36

Locus typicus. China (Jiangsu and Zhejiang).

Etymology. The new species is named after its host plant, *Sinojackia xylocarpa*.

Description of mounted specimens. *Apterous viviparous females* (Table 1; Figs 3–13, 20–28). Body oval (Fig. 20). Measurements: body 1.86–2.09 long, 1.01–1.24 wide. Cephalic setae, marginal setae on abdominal tergite I and dorsal setae on abdominal tergite VIII 0.04–0.06, 0.04, 0.06–0.07 long. Antennae 0.61–0.68 long, segment III 0.19–0.23 long. Setae on segment III 0.03 long. Ultimate rostral segment



Figures 3–19. *Aleurodaphis sinojackiae* Qiao & Jiang, sp. n. **3–13** Apterous viviparous female. **3** dorsal view of head **4** antennae **5** ultimate rostral segment **6** dorsal view of thorax **7** hind tarsal segments **8** abdominal tergite VIII **9** dorsal setae and sculptures on abdominal tergite VI **10** siphunculus **11** cauda **12** anal plate **13** genital plate. **14–19** Alate viviparous female. **14** antennae **15** ultimate rostral segment **16** basal half of fore wing **17** siphunculus **18** cauda **19** anal plate. Scale bars = 0.10 mm.

Table 1. Measurements of apterous viviparous females of *Aleurodaphis sinojackiae* Qiao & Jiang, sp. n. (mm)

No.	Body length	Body width	Ant. I	Ant. II	Ant. III	Ant. IV	Ant. V b	Ant. V pt	URS	Hind T & F	2HTs	Cauda length
1	1.92	1.15	0.07	0.08	0.21	0.12	0.13	0.04	0.13	0.36	0.10	0.10
2	1.98	1.13	0.07	0.07	0.20	0.10	0.12	0.03	0.15	0.36	0.10	0.09
3*	2.05	1.21	0.07	0.08	0.23	0.12	0.13	0.03	0.14	0.38	0.10	0.09
4	1.89	1.07	0.07	0.08	0.19	0.12	0.12	0.03	0.12	0.33	0.10	0.09
5	1.86	1.09	0.08	0.07	0.21	0.11	0.12	0.04	0.14	0.37	0.11	0.08
6	2.09	1.01	0.08	0.08	0.23	0.12	0.14	0.04	0.13	0.39	0.11	0.09
7	2.05	1.24	0.07	0.08	0.22	0.12	0.13	0.03	0.15	0.37	0.10	0.09
8	2.03	1.05	0.07	0.08	0.22	0.11	0.13	0.04	0.14	0.37	0.10	0.09
Average	1.98	1.12	0.07	0.08	0.21	0.11	0.13	0.03	0.14	0.37	0.10	0.09


Remark. * Holotype; for abbreviations see Materials and Methods.

Table 2. Measurements of alate viviparous females of *Aleurodaphis sinojackiae* Qiao & Jiang, sp. n. (mm)

No.	Body length	Body width	Ant. I	Ant. II	Ant. III	Ant. IV	Ant. V b	Ant. V pt	URS	Hind femur	2HTs	Cauda length
1	1.84	0.81	0.08	0.07	0.28	0.14	0.14	0.04	0.14	0.40	0.09	0.08
2	2.00	0.82	0.07	0.07	0.25	0.14	0.13	0.04	0.13	0.39	0.11	0.08
4	1.49	0.66	0.06	0.06	0.18	0.11	0.09	0.02	0.10	0.29	0.08	0.06
5	1.45	0.63	0.05	0.05	0.19	0.09	0.08	0.02	0.10	0.30	0.08	0.06
6	1.32	0.64	0.05	0.05	0.21	0.11	0.11	0.03	0.11	0.31	0.08	0.05
7	1.42	0.56	0.06	0.05	0.18	0.11	0.09	0.02	0.11	0.31	0.07	0.06
8	1.30	0.58	0.06	0.05	0.19	0.10	0.09	0.03	0.11	0.26	0.08	0.06
Average	1.45	0.64	0.06	0.05	0.20	0.11	0.10	0.03	0.11	0.30	0.08	0.06

0.12–0.15 long. Hind trochanter and femur 0.33–0.39 long, hind tibia 0.38–0.44 long, second hind tarsal segment 0.10–0.11 long. Setae on hind tibia 0.04–0.06 long. Apical diameter of siphunculi 0.04–0.05. Cauda 0.08–0.10 long.

Head and pronotum (Figs 3, 21), mesonotum and metanotum (Fig. 6), abdominal tergites I–VII fused (Fig. 20), respectively; tergite VIII free (Fig. 8). Antennae, rostrum and legs brown; cauda, anal plate and genital plate dark brown. Dorsum of body rough, covered with dense sculptures on dorsum of head and thoracic notums, and with sparse sculptures on abdominal tergites (Fig. 9). Dorsum of body with round marginal wax glands, composited with big facets (Figs 8, 25, 26). Pro- and metanotum each with 13 wax glands, mesonotum with 8 wax glands, abdominal tergites I–VII each with 3–6 pairs of wax glands; tergite VIII with 10–13 wax glands. Dorsal setae of body fine and short (Fig. 9). Head with 2 pairs of cephalic and spinal setae, 3 pairs of marginal setae; pronotum with 2 pairs of spinal, 1 pair of pleural and 2 pairs of marginal setae; mesonotum with 2 pairs of spinal, 3 pairs of pleural and 2 pairs of marginal setae; metanotum with 1 pair of spinal, 3 pairs of pleural and 2 pairs of marginal setae; abdominal tergite I with 1 pair of spinal, pleural and marginal setae; tergites II–VII each with 1 pair of spinal and marginal setae; tergite VIII with 1 pair of spinal and 5 marginal setae.



Figures 20–28. *Aleurodaphis sinojackiae* Qiao & Jiang, sp. n. Apterous viviparous female. **20** dorsal view of body **21** dorsal view of head and pronotum **22** antenna **23** ultimate rostral segment **24** fore tarsal segments **25** siphunculus **26** abdominal tergite VIII and cauda **27** anal plate **28** genital plate. Scale bars = 0.10 mm.



Figures 29–34. *Aleurodaphis sinojackiae* Qiao & Jiang, sp. n. Alate viviparous female. **29** dorsal view of body **30** antennae **31** ultimate rostral segment **32** siphunculus **33** cauda **34** anal plate. Scale bars = 0.10 mm.

Cephalic setae, marginal setae on abdominal tergite I, setae on abdominal tergite VIII 1.20–1.60, 0.45–1.20 and 1.09–2.00 times as long as widest diameter of antennal segment III, respectively. Spiracles oval, closed, on brown oval spiracular plates.

Head: Front flat and straight. Eyes with 3 facets. Antennae 5-segmented (Figs 4, 22), with spinulose imbrications on segments III–V, 0.33–0.38 times as long as body. Length in proportion of segments I–V: 31–38 : 34–42 : 100 : 50–62 : 56–65+13–21, respectively. Processus terminalis 0.25–0.33 times as long as base of the segment V. Segments I–V each with 2–4, 2 or 3, 0 or 1, 1 or 2, 2+0 setae, respectively. Processus terminalis with 5 or 6 apical setae. Setae on segment III 0.58 times as long as widest diameter of the segment. Primary rhinaria small and round. Rostrum short, reaching mid-coxae. Ultimate rostral segment acute wedge-shaped (Figs 5, 23), 2.67–3.33 times as long as its basal width, 1.14–1.43 times as long as second hind tarsal segment; with 2 pairs of primary setae and 1 or 2 pairs of secondary setae.

Thorax: Mesosternal furca with two separated arms, each arm 1.21–1.41 times as long as widest diameter of antennal segment III. Legs normal. Trochanter and femora fused, hind trochanter and femur 1.63–1.85 times as long as antennal segment III, hind tibia 0.20–0.22 times as long as body; setae on hind tibia 0.88–0.94 times as long as its mid-diameter. First tarsal chaetotaxy: 4, 4, 4, sometimes 3, 3, 4 or 4, 4, 3. Second hind tarsal segment with 2 setae between claws and each seta with funnel-shaped apex (Figs 7, 24).

Abdomen: Siphunculi pore-like (Figs 10, 25), on abdominal tergite VI, apical diameter 1.00–1.40 times as long as widest diameter of antennal segment III. Cauda, anal plate and genital plate with spinulose imbrications. Cauda knobbed (Figs 11, 26), constricted in middle, 0.55–0.68 times as long as its basal width, with 9 or 10 apical setae. Anal plate bilobed (Figs 12, 27), each with 6–8 setae. Genital plate broad band-shaped (Figs 13, 28), with 3 or 4 anterior setae and 14–23 middle and posterior marginal setae. Two gonapophyses each with 5 short setae.

Alate viviparous females (Table 2; Figs 4–19, 29–34). Body oval (Fig. 29). Measurements: body 1.30–2.00 long, 0.56–0.82 wide. Cephalic setae, marginal setae on abdominal tergite I and dorsal setae on abdominal tergite VIII 0.016–0.021, 0.015–0.017, 0.020–0.026 long, respectively. Antennae 0.49–0.74 long, segment III 0.18–0.28 long. Ultimate rostral segment 0.10–0.14 long. Hind femur 0.26–0.40 long, hind tibia 0.36 long, second hind tarsal segment 0.07–0.11 long. Setae on hind tibia 0.030 long. Fore wing 1.64–1.74 long. Apical diameter of siphunculi 0.04–0.05. Cauda 0.05–0.08 long.

Dorsum of body dark brown, antenna, apex of rostrum, legs, cauda, anal plate and genital plate brown. Dorsal setae of body fine, short and pointed, slightly longer than ventral setae. Head with 2 pairs of cephalic setae, 2 pairs of setae between antennae and 2 pairs of setae between eyes; abdominal tergites I–VII each with 1 pair of spinal and marginal setae; tergite VIII with 1 pair of spinal setae. Cephalic setae, marginal setae on abdominal tergite I, setae on abdominal tergite VIII 0.51–0.67, 0.50–0.54 and 0.64–0.83 times as long as widest diameter of antennal segment III, respectively.

Head: Front rounded. Antennae 5-segmented (Figs 14, 30), with sparse imbrications on segments I–II and dense spinulose imbrications on segments III–V. Whole length of antennae 0.37–0.43 times as long as body, length in proportion of segments I–V: 21–32 : 24–33 : 100 : 49–60 : 41–53+10–15, respectively. Processus terminalis 0.29–0.37 times as long as base of the segment V. Segments I–V each with 3–5, 2 or 3, 0 or 1, 1, 1 or 2+0 setae, respectively. Processus terminalis with 5 apical setae. Primary rhinaria irregular ring-shaped. Segments III, IV and base of Segment V each with 10–14, 3–6 and 2–4 secondary rhinaria, respectively. Rostrum short, reaching mid-coxae. Ultimate rostral segment (Figs 15, 31) 2.50–2.86 times as long as its basal width, 1.20–1.54 times as long as second hind tarsal segment; with 2 or 3 pairs of primary setae and 1 or 2 pairs of secondary setae.

Thorax: Legs normal. Hind femur 1.50–1.62 times as long as antennal segment III, hind tibia 0.25–0.28 times as long as body; setae on hind tibia 0.91–1.20 times as long as its mid-diameter. First tarsal chaetotaxy: 4, 4, 3. Fore wing (Figs 16, 29) 1.17–1.34 times as long as body, 2.00–2.42 times as long as width of the wing. Media

once branched. Pterostigma long and curved to the apex of the wing. Hind wings with one thick longitudinal vein and two oblique veins.

Abdomen: Siphunculi pore-like (Figs 17, 32), apical diameter 1.33–1.51 times as long as widest diameter of antennal segment III. Cauda knobbed (Figs 18, 33), constricted in middle, 0.68–0.86 times as long as its basal width, with 6–8 apical setae. Anal plate bilobed (Figs 19, 34), each with 6 or 7 setae. Genital plate broad band-shaped, with 3 anterior setae and 12–15 posterior marginal setae. Two gonapophyses each with 5 or 6 setae.

Embryos. Body oval, with wax glands arranged along crenulated margin in both apterae and alatae. Cephalic setae short and pointed. Antennae 4-segmented. Rostrum and legs well developed. Legs covered with dense setae. Siphunculi visible.

Type material examined. Holotype, 1 apterous viviparous female, **CHINA:** Zhejiang (Hangzhou City), 21 May 1985, on *Sinojackia xylocarpa*, coll. V. F. Eastop (BMNH). Paratypes, 28 apterous viviparous females, 2 alate viviparous females and 8 nymphs, with the same collection data as holotype (BMNH); 9 apterous viviparous females and 11 alate viviparous females, **CHINA:** Jiangsu (Zhongshan Botanic Garden, Nanjing City, Alt. 100m), No. Y7116, 10 June 1987, on *Sinojackia xylocarpa*, coll. T. S. Zhong (NZMC); 1 apterous viviparous female and 1 alate viviparous female, **CHINA:** Jiangsu (Zhongshan Botanic Garden, Nanjing City, Alt. 100m), No. Y7116, 10 June 1987, on *Sinojackia xylocarpa*, coll. T. S. Zhong (Kôgakkan University, Japan).

Host plants. *Sinojackia xylocarpa*.

Biology. The species induced the leaves of host plants to curl and form boat-shaped leaf galls.

Diagnosis. The new species differs from the other known species as follows: in apterous viviparous female: wax glands arranged in each segment, not connecting with each other (the other species: arranged continuously along the edge of body as a crenulation, or without wax glands); in alate viviparous female compared to the most similar species *A. mikaniae*: antennal segment III with 10–14 secondary rhinaria (*A. mikaniae*: 24–27); first tarsal chaetotaxy: 4, 4, 4, sometimes 3, 3, 3 (*A. mikaniae*: 3, 3, 3, sometimes 2, 2, 2).

Remark. As the detailed biological information is very important to research the taxonomic position of the genus and species identification, the life cycle of the new species will receive further study in future.

Aleurodaphis stewartiae Sorin & Miyazaki, 2004

http://species-id.net/wiki/Aleurodaphis_stewartiae

Aleurodaphis stewartiae Sorin & Miyazaki, 2004: 174.

Host plants. Primary host: *Stewartia monadelphica*. Secondary hosts unknown.

Biology. The aphid induces a leaf gall, which is formed by rolling the marginal part of the leaf upwards. The gall is about 47.5 long and 7.2 wide, with a surface rough to the touch. The alate viviparous females emerge in early August, and then disappear

from the host tree, probably emigrating to some unknown secondary host (Sorin and Miyazaki 2004).

Distribution. Japan (Sorin and Miyazaki 2004).

Acknowledgments

The authors cordially thank Prof. M Sorin in Kôgakkan University, Japan for his checking specimens of the new species and providing the related Japanese references for our research, and are deeply indebted to all the specimen collectors in this study; RL Blackman, VD Bosch, VS Calilung, XF Dai, VF Eastop, MR Gavarra, van der Goot, K Guo, TL He, XL Huang, EB Ma, WH Paik, YY Rao, M Sorin, DL Stern, H Takada, R Takahashi, Theobald, SH Wang, JY Yang, GX Zhang, WY Zhang and TS Zhong. We also thank Miss CP Liu for making slides of the new species. The work was supported by the National Science Funds for Distinguished Young Scientists (No. 31025024), National Natural Sciences Foundation of China (Nos. 30830017, 30970391), National Science Fund for Fostering Talents in Basic Research (No. J0930004), a grant (No. O529YX5105) from the Key Laboratory of the Zoological Systematics and Evolution of the Chinese Academy of Sciences, and the Ministry of Science and Technology of the People's Republic of China (MOST GRANT No. 2006FY110500).

References

- Baker AC (1929) Generic classification of the Hemipterous family Aphididae. *Bulletin, United States, Department of Agriculture* 826: 1–109.
- Blackman RL, Eastop VF (1994) *Aphids on the World's Trees. An Identification and Information Guide*. CAB International in Association with the Natural History Museum, Wallingford, UK, 987 pp.
- Blackman RL, Eastop VF (2006) *Aphids on the World's Herbaceous Plants and Shrubs*. The Natural History Museum, John Wiley & Sons Ltd, Chichester, UK, 1439 pp.
- Calilung VJ (1967) A faunistic study of Philippine Aphids. *Philippine Agricultural Scientist* 51: 93–170.
- Chakrabarti S, Maity SP (1982) New genus, new species and new records of Cerataphidine aphids Homoptera: Aphididae) from northwest Himalaya. *Proceedings of the Zoological Society (Calcutta)* 33(1–2): 55–63.
- Eastop VF, Hille Ris Lambers D (1976) *Survey of the World's Aphids*. Dr. W. Junk b.v., Publishers, The Hague, 573 pp.
- van der Goot P (1917) Zur Kenntnis der Blattläuse Javas. *Contrib. Faune Indes Neerl* 1(3): 1–301.
- Ghosh AK (1988) *The Fauna of India and the Adjacent Countries (Homoptera: Aphidoidea) Part 4 Subfamilies: Phloeomyzinae, Anoeciinae and Hormaphidinae*. *Zoological Survey of India, Calcutta*, 429 pp.

- Noordam D (1991) Hormaphidinae from Java (Homoptera: Aphididae). Zoologische Verhandlungen (Leiden) 270: 1–525.
- Raychaudhuri DN, Ghosh MR, Pal PK, Ghosh AK (1980) Taxonomy of aphids of north-east India and Bhutan. Taxonomic accounts of the subfamilies. Subfamily Hormaphidinae. 358–394. In Raychaudhuri DN. Aphids of north-east India and Bhutan. The Zoological Society, i–viii, Calcutta, 521pp.
- Remaudière G, Remaudière M (1997) Catalogue of the World's Aphididae. Institut National de la Recherche Agronomique, Paris, 473 pp.
- Shinji GO (1923) One new species of aphids from Saitama and Maroika. Doubts. Zasshi. 35(417): 301–309.
- Sorin M, Miyazaki M (2004) *Aleurodaphis* of Japan (Hemiptera, Aphididae). Japanese Journal of Systematic Entomology 10(2), 165–177.
- Takahashi R (1921) Aphididae of Formosa. Part I. Agriculture Experiment Station Government of Formosa, Report 20: 1–97.
- Takahashi R (1923) Aphididae of Formosa. Part 2. Department of Agriculture Government Research Institute Formosa, Report 4: 1–173.
- Takahashi R (1924) Aphididae of Formosa. Part 3. Department of Agriculture Government Research Institute Formosa, Report 10: 1–121.
- Takahashi R (1925) Aphididae of Formosa. Part 4. Department of Agriculture Government Research Institute Formosa, Report 16: 1–65.
- Takahashi R (1931) Aphididae of Formosa. Part 6. Department of Agriculture Government Research Institute Formosa, Report 53: 1–127.
- Takahashi R, Sorin M (1958) Notes on the aphid of *Aleurodaphis* in Japan (Homoptera). Akitu, Kyoto 7: 31–32.
- Tao CC (1990) Aphid-fauna of Taiwan Province, China. Taiwan Provincial Museum Press, Taipei, 327pp.
- Tao CC (1999) List of Aphidoidea (Homoptera) of China. Taiwan Agricultural Research Institute Special Publication 77: 1–144.
- Zhang GX, Zhong TS (1982) New species and new subspecies of Chinese Aphidoidea (Homoptera). Sinozoologia 2: 19–28.
- Zhang GX, Zhong TS, Zhang WY (1992) Aphidoidea. 137–175. In: Peng JW, Liu YQ. Eds. 1992. Iconography of forest insects in Hunan China. Academia Sinica and Hunan Forestry Institute, Hunan, China. Hunan Science and Technology Press, Changsha, 473pp.