

The order Zoantharia Rafinesque, 1815 (Cnidaria, Anthozoa: Hexacorallia): supraspecific classification and nomenclature

Martyn E. Y. Low^{1,4}, Frederic Sinniger², James Davis Reimer³

1 Lee Kong Chian Museum of Natural History, National University of Singapore, 2 Conservatory Drive, Singapore 117377, Republic of Singapore **2** Tropical Biosphere Research Center, University of the Ryukyus, 3422 Sesoko, Motobu, Okinawa 905-0227, Japan **3** Molecular Invertebrate Systematics and Ecology (MISE) Laboratory, Department of Marine and Environmental Sciences, Graduate School of Engineering and Science, University of the Ryukyus, 1 Senbaru, Nishihara, Okinawa 903-0213, Japan; and Tropical Biosphere Research Center, University of the Ryukyus, 1 Senbaru, Nishihara, Okinawa 903-0213, Japan **4** former address: Department of Marine and Environmental Sciences, Graduate School of Engineering and Science, University of the Ryukyus, 1 Senbaru, Nishihara, Okinawa 903-0213, Japan

Corresponding authors: James Davis Reimer (jreimer@sci.u-ryukyu.ac.jp); Martyn E. Y. Low (martyn@natural-history.org)

Academic editor: L. van Ofwegen | Received 31 August 2016 | Accepted 28 November 2016 | Published 14 December 2016

<http://zoobank.org/903D6413-C802-4864-A662-D71C50740E2D>

Citation: Low MEY, Sinniger F, Reimer JD (2016) The order Zoantharia Rafinesque, 1815 (Cnidaria, Anthozoa: Hexacorallia): supraspecific classification and nomenclature. ZooKeys 641: 1–80. <https://doi.org/10.3897/zookeys.641.10346>

Abstract

Many supraspecific zoantharian names have long and complicated histories. The present list is provided to advise researchers on the current state of supraspecific nomenclature of the zoantharians, particularly given the recent attention paid to the taxonomy, phylogeny, and biodiversity of this order. At the same time, several taxonomic issues brought to light by recent research are resolved. Details on the taxonomic and nomenclatural history of most groups are provided, along with appendices of invalid supraspecific names.

Keywords

zoantharians, family-group, genus-group, taxonomy, historical literature

Contents

Introduction.....	3
Systematics.....	4
Phylum Cnidaria Hatschek, 1888	4
Class Anthozoa Ehrenberg, 1831.....	5
Subclass Hexacorallia Haeckel, 1896.....	5
Order Zoantharia (Zoanthidea) Rafinesque, 1815.....	5
Suborder Brachycnemina Haddon and Shackleton, 1891.....	5
Neozoanthidae Herberts, 1972.....	6
<i>Neozoanthus</i> Herberts, 1972.....	6
Sphenopidae Hertwig, 1882.....	6
<i>Sphenopus</i> Steenstrup, 1856.....	7
<i>Palythoa</i> Lamouroux, 1816.....	7
Zoanthidae Rafinesque, 1815.....	8
<i>Zoanthus</i> Lamarck, 1801.....	8
<i>Acrozoanthus</i> Saville-Kent, 1893.....	9
<i>Isaurus</i> Gray, 1828	10
Suborder Macrocnemina Haddon and Shackleton, 1891	11
Epizoanthidae Delage and Hérouard, 1901	11
<i>Epizoanthus</i> Gray, 1867.....	11
<i>Palaeozoanthus</i> Carlgren, 1924	12
<i>Thoracactis</i> Gravier, 1918.....	12
Hydrozoanthidae Sinniger, Reimer and Pawlowski, 2010.....	13
<i>Hydrozoanthus</i> Sinniger, Reimer and Pawlowski, 2010	13
<i>Terrazoanthus</i> Reimer and Fujii, 2010	13
Microzoanthidae Fujii and Reimer, 2011	14
<i>Microzoanthus</i> Fujii and Reimer, 2011	14
Nanozoanthidae Fujii and Reimer, 2013	15
<i>Nanozoanthus</i> Fujii and Reimer, 2013.....	15
Parazoanthidae Delage and Hérouard, 1901.....	15
<i>Parazoanthus</i> Haddon and Shackleton, 1891.....	16
<i>Antipathozoanthus</i> Sinniger, Reimer and Pawlowski, 2010.....	17
<i>Bergia</i> Duchassaing de Fombressin and Michelotti, 1860.....	18
<i>Bullagummizoanthus</i> Sinniger, Ocaña and Baco, 2013.....	18
<i>Corallizoanthus</i> Reimer, in Reimer, Nonaka, Sinniger and Iwase, 2008.....	18
<i>Hurlizoanthus</i> Sinniger, Ocaña and Baco, 2013.....	19
<i>Isozoanthus</i> Carlgren, in Chun, 1903.....	19
<i>Kauluzoanthus</i> Sinniger, Ocaña and Baco, 2013.....	20
<i>Kulamanamana</i> Sinniger, Ocaña and Baco, 2013.....	20
<i>Mesozoanthus</i> Sinniger and Häussermann, 2009.....	21
<i>Savalia</i> Nardo, 1844	21
<i>Umimayanthus</i> Montenegro, Sinniger and Reimer, 2015.....	22

<i>Zibrowius</i> Sinniger, Ocaña and Baco, 2013	22
Suborder <i>incerta sedis</i>	22
Abysoanthidae Reimer and Fujiwara, in Reimer, Sinniger, Fujiwara, Hirano and Maruyama, 2007	23
<i>Abysoanthus</i> Reimer and Fujiwara, in Reimer, Sinniger, Fujiwara, Hirano and Maruyama, 2007	23
Family <i>incerta sedis</i>	23
<i>Stephanidium</i> Hertwig, 1888.....	23
Discussion.....	24
Acknowledgements.....	24
References	25
Appendix 1: Nomenclator of family-group names	49
Appendix 2: Nomenclator of genus-group names.....	52
Appendix 3: Nomenclatural remarks	64

Introduction

The Zoantharia are an order of Hexacorallia most closely related to the sea anemones (Actiniaria) (Kayal et al. 2013). They are found in most marine ecosystems. Despite their cosmopolitan distribution and high frequency in many ecosystems, compared to both Actiniaria and Scleractinia, research on this order has been relatively scant. This is due to both their high levels of intraspecific morphological variation (e.g. Burnett et al. 1997, Reimer et al. 2004, Ong et al. 2013) and the presence of incrustations in their body walls (for all zoantharian families except the Zoanthidae), which makes internal histological examination problematic (Reimer et al. 2010c). Taxonomic identification problems have been compounded by a general lack of utility in some traditionally used diagnostic characters (Sinniger et al. 2010a, Swain 2010). Thus, despite being common in many ecosystems (Karlson 1980), until very recently zoantharian taxonomy has been confused (Burnett et al. 1997).

However, recent molecular phylogenetic examination of zoantharians, combined with a reassessment of their taxonomy and diagnostic characters, has led to the group becoming somewhat ordered (Reimer et al. 2007a, 2008a, Reimer and Fujii 2010, Sinniger et al. 2010a, 2013, Fujii and Reimer 2011, 2013, Swain and Swain 2014; Swain et al. 2015), setting an example for the reorganization of other anthozoan groups. This follows in the footsteps of the database Biogeoinformatics of Hexacorallia (Fautin and Buddemeier 2008), which was established as a repository for taxonomic and nomenclatural information on the Hexacorallia (see also Fautin 2016).

The order Zoantharia (= Zoanthidea or Zoanthiniaria) is currently thought to be comprised of nine families—Abysoanthidae Reimer and Fujiwara, in Reimer, Sinniger, Fujiwara, Hirano and Maruyama, 2007, Epizoanthidae Delage and Hérourard, 1910, Hydrozoanthidae Sinniger, Reimer and Pawlowski, 2010, Microzoanthidae Fujii and Reimer, 2011, Nanozoanthidae Fujii and Reimer, 2013, Neozoanthidae

Herberts, 1972, Parazoanthidae Delage and Hérourard, 1901, Sphenopidae Hertwig, 1882, and Zoanthidae Rafinesque, 1815 (Daly et al. 2007: 143, 144; Hoeksema and Reimer 2013).

There is also a bioinformatics rationale for the compilation of such a taxonomic listing. As electronic name-lists of available and valid names become increasingly accessible, the need to link such lists with nomenclators of names will become an imperative. To this end, listings such as the present one can provide clear linkages between valid names and the huge mass of historical names (most of which are synonyms and some being potentially valid), which will effectively reduce the historical ‘synonymy load’ (Low and Reimer 2012a: 83) and provide a clear list of available supraspecific names of Zoantharia for future researchers.

Additional notes for introductory text

All valid taxa are treated in the systematics section, with the taxon (genus and species) listed. Synonyms are also listed. Justification for these synonymies is given in the sections that follow (Appendices 1–3). We use the general term “zoantharian” to refer to species within this order, unless when quoting original text in which another term (usually “zoanthid”) was used.

Statistics

A total of 16 family-group names are treated. Nine are considered to be valid, six are synonyms, and one is not referable to the order Zoantharia. A total of 102 names at the genus rank are treated. Of these, 28 names are considered to be valid, 38 to be synonyms, 19 are incorrect spellings, 18 are not referable to the order Zoantharia, and two—*Stephanidium* Hertwig, 1888, and *Triga* Gray, 1867—are of uncertain placement and validity (see Table 1).

Systematics

Phylum Cnidaria Hatschek, 1888

Cnidaria Hatschek, 1888 (in 1888–1891): 40.

Remarks. See discussion on the nomenclature of the phylum in Fautin and Daly (2009: 315).

Class Anthozoa Ehrenberg, 1831

Anthozoa Ehrenberg, 1831: 44.

Subclass Hexacorallia Haeckel, 1896

Zoantharia de Blainville, 1830: 274.

Hexacorallia Haeckel, 1896: 217.

Order Zoantharia Rafinesque, 1815

Zoanthia Rafinesque, 1815: 155.

Zoantharia Gray, 1832: 94.

Zoanthiniaria van Beneden, 1897: 150.

Zoanthidea Bourne, 1900: 58.

Diagnosis. Anthozoans with body walls usually incrustated with sand and/or other detritus (except for the family Zoanthidae), tentacles always arranged in two rows or cycles. Majority of species are colonial.

Remarks. Herein, we choose to use the name *Zoantharia* Rafinesque, 1815. Although *Zoantharia* Rafinesque, 1815, has identical spelling with the supraordinal name *Zoantharia* de Blainville, 1830, the latter name has fallen from common use—*Hexacorallia* Haeckel, 1896, being favoured. Furthermore, using the name *Zoanthidea* Bourne, 1900, can potentially cause confusion when used in the non-technical form “zoanthids” to denote members of the order, as it would be identical with the term “zoanthids” to denote members of the family *Zoanthidae* Rafinesque, 1815.

Suborder Brachycnemina Haddon and Shackleton, 1891

Brachycneminae Haddon and Shackleton, 1891a: 626.

Diagnosis. Zoantharians with the fifth mesenteries from the dorsal directive being incomplete.

Remarks.—Recent molecular phylogenetic research indicates the two suborders of *Zoantharia* may not be monophyletics (e.g. Fujii and Reimer 2013). Three families are currently assigned to *Brachycnemina* Haddon and Shackleton, 1891.

Neozoanthidae Herberts, 1972

Neozoanthidae Herberts, 1972: 137.

Type genus. *Neozoanthus* Herberts, 1972.

Gender. Masculine.

Diagnosis. Zooxanthellate, brachycnemic zoantharians with only partial incrustation in ectoderm, rarely extending to mesoglea, no incrustation around the top, oral ends of polyps (Herberts 1972).

Remarks. A monogeneric and monospecific family-group. This taxon was originally defined as “Zoanthaires à arrangement mésentérique de type brachycnémique et à sphincter endodermique” (Herberts 1972: 137). However, recent work has further divided zoantharian sphincter muscles into ten different basic types; that of Neozoanthidae is discontinuous endodermal (Swain et al. 2015). Thus, the above diagnosis has been slightly modified from that of Herberts’ (1972).

Recent molecular phylogenetic work calls into the question the validity of this family, as molecular data derived from *Neozoanthus* specimens cluster within the Zoanthidae radiation (Reimer et al. 2011a). Incrustation does not extend to mesoglea and is only found in the ectoderm. The families Neozoanthidae and Zoanthidae are not synonymised herein, as the Neozoanthidae also has some phylogenetic relation with the Hydrozoanthidae and the relationships between these families will require additional research (Reimer et al. 2011a).

Neozoanthus Herberts, 1972

Neozoanthus Herberts, 1972: 137.

Type species. *Neozoanthus tulearensis* Herberts, 1972, by monotypy.

Gender. Masculine.

Diagnosis. As for family above.

Remarks. Molecular phylogenetic results (Reimer et al. 2011a) indicate that this genus appears to be very closely related to the genus *Isaurus* Gray, 1828 (a genus assigned to the family Zoanthidae). Three species included in this group are from the Indo-Pacific.

Sphenopidae Hertwig, 1882

Palythoidae (as “Palythoae”) Duchassaing de Fonbressin and Michelotti, 1860: 37.

Sphenopidae Hertwig, 1882: 120.

Type genus. *Sphenopus* Steenstrup, 1856.

Gender. Masculine.

Diagnosis. Brachycnemic zoantharians with sand/detritus incrustation in the ectoderm and mesoglea.

Remarks. The family-group name Sphenopidae Hertwig, 1882, is currently threatened by the senior subjective synonym Palythoidae Duchassaing de Fonbressin and Michelotti, 1860. To maintain widespread and current usage of the former name, and in accordance with Article 23.9.2 (ICZN 1999: 28, 29), an application is being prepared to request the International Commission on Zoological Nomenclature to suppress the senior subjective synonym Palythoidae Duchassaing de Fonbressin and Michelotti, 1860, in favour of Sphenopidae Hertwig, 1882. See additional discussion in Appendix 3.

The family-group Sphenopidae was established by Hertwig (1882), with the inclusion of only the type genus, *Sphenopus* Steenstrup, 1856. Two genera are currently assigned to Sphenopidae Hertwig, 1882.

***Sphenopus* Steenstrup, 1856**

Sphenopus Steenstrup, 1856: 37.

Type species. *Sabella marsupialis* Gmelin, 1791, by monotypy.

Gender. Masculine.

Diagnosis. Unitary (=solitary, non-colonial) brachycnemic zoantharians with sand/detritus incrustation in the ectoderm and mesoglea.

Remarks. Distinct, large unitary polyps found embedded in sandy habitats with the oral disc clear of, and not attached to substrate (Soong et al. 1999, Reimer et al. 2012b, 2015b). Recent molecular work has also indicated that this this genus-group may form a clade with some *Palythoa* spp. (Reimer et al. 2012b, Fujii and Reimer 2016).

***Palythoa* Lamouroux, 1816**

Palythoa Lamouroux, 1816: 359.

Corticifera Le Sueur, 1817: 178.

Polythoa Schweigger, 1819: 100 [incorrect spelling].

Polythea Gistel, 1848: 181 [incorrect spelling].

Gemmaria Duchassaing de Fonbressin and Michelotti, 1860: 55.

Polythoa (*Corticithoa*) Andres, 1883a: 521, 535–538.

Polythoa (*Gemmithoa*) Andres, 1883a: 521, 532, 533.

Parapalythoa Verrill, 1900: 560.

Protopolythoa Verrill, 1900: 562.

Haplotella Stechow, 1919: 853.

Type species. *Palythoe* [*sic*] *stellata* Lamouroux, 1816 [= *Alcyonium mammillosum* Ellis, in Ellis and Solander, 1786], by subsequent designation by Haddon and Shackleton (1891b: 691) (see also discussion in Appendix 3).

Gender. Feminine.

Diagnosis. Colonial brachycnemic zoantharians with sand/detritus incrustation in the ectoderm and mesoglea. Currently, all species except for two described are zooxanthellate (Irei et al. 2015).

Remarks. Specimens examined with a sphincter muscle of linear mesogleal type (Swain et al. 2015).

Protopalychtha was originally separated from *Palythoa* based on primarily polyp shape. Pax (1910) recommended merging *Palythoa* and *Protopalychtha*, and he defined three groups; “immersae”, “intermediae” and “liberae”, which have no taxonomic rank (see Ryland and Lancaster 2003). In 1923, Carlgren adopted this inclusive nomenclature. Similarly, Burnett et al. (1997) discussed the need to merge these genera but retained *Protopalychtha*. Ryland and Lancaster (2003) also discussed this issue and argued to keep *Protopalychtha* citing additional characters (zooxanthellate eggs in *Protopalychtha*; afterwards potentially observed in *Palythoa* spp. (Shiroma et al. 2010)). Herein, based on the several recent molecular studies (Sinniger et al. 2005, Reimer et al. 2006, 2007b, 2012a), we formally synonymise *Palythoa* and *Protopalychtha*, with *Palythoa* having priority.

Although at least 272 species-group names have been established in the genera *Palythoa* and *Protopalychtha* in the literature (see Fautin and Buddemeier 2008), with more research, these species-group names will likely be found to be synonyms—the result of high level of intraspecific morphological variation (see Burnett et al. 1994, Reimer et al. 2006, Hibino et al. 2014).

Zoanthidae Rafinesque, 1815

Zoanthidae (as “Zoanthia”) Rafinesque, 1815: 155.

Zoanthidae Gray, 1832: 95.

Type genus. *Zoanthus* Lamarck, 1801.

Diagnosis. Brachycnemic zoantharians with no or little sand/detritus encrustation. Continuous or divided marginal muscle, and zoanthinae larvae (Ryland and Lancaster 2003).

Remarks. All members of this family are zooxanthellate (with endosymbiotic *Symbiodinium* spp.), and are found in sub-tropical and tropical waters. Authorship of the family-group name Zoanthidae should be attributed to Rafinesque (1815), and not Gray (1832, 1840) (see Low and Reimer 2012a). Three genera are currently assigned to Zoanthidae Rafinesque, 1815.

Zoanthus Lamarck, 1801

Zoanthus Cuvier, 1800: tables 9–10 [*nomen nudum*].

Zoanthus (as “*Zoantha*”) Lamarck, 1801: 363.

Mammillifera Le Sueur, 1817: 177.

Actimastus Rafinesque, 1818: 271 [unnecessary replacement name]

Mamillifera Quoy and Gaimard, 1834: 169 [incorrect spelling]

Anthozoon Gistel, 1848: 181 [unnecessary replacement name]

Mammilifera Gistel, 1848: 181 [incorrect spelling]

Polythoa (*Mammithoa*) Andres, 1883a: 521.

Polythoa (*Mammothoa*) Andres, 1883a: 533 [incorrect spelling].

Zoanthus (*Rhyzanthus*) Andres, 1883a: 538.

Type species. *Actinia sociata* Ellis, 1768, by monotypy.

Gender. Masculine.

Diagnosis. Zooxanthellate, absence of mineral incrustations in the column/coenenchyme (excluding superficial surface attachments) brachycnemic zoantharians with smooth, usually erect polyps—except in *Zoanthus praelongus* Carlgren, 1954 (see discussion below)—with no endodermal invaginations. Mesogleal sphincter muscle with clear distal and proximal sections, mesogleal canal system but no encircling sinus (Duerden 1898, Swain and Swain 2014).

Remarks. Species of the genus *Zoanthus* and *Acrozoanthus* have a double sphincter muscle, which is unique among zoantharians. Referred to in Swain et al. (2015) as discontinuous mesogleal type.

Despite over 150 species having been described in or assigned to this genus (see Fautin and Buddemeier 2008), many of the species-group names are likely to be synonyms—due to the high level of intraspecific morphological variation (see Burnett et al. 1995, 1997, Reimer et al. 2004). One species, *Zoanthus praelongus* Carlgren, 1954, has recumbent (non-erect) polyps (as in the genus *Isaurus*) but is clearly referable to *Zoanthus* (see Carlgren 1954, Muirhead and Ryland 1985, Reimer et al. 2008b).

Acrozoanthus Saville-Kent, 1893

Acrozoanthus Saville-Kent, 1893: 153, 154.

Type species. *Acrozoanthus australiae* Saville-Kent, 1893, by monotypy.

Gender. Masculine.

Diagnosis. Zooxanthellate, absence of mineral incrustations in the column/coenenchyme (excluding superficial surface attachments) brachycnemic zoantharians inhabiting the outside of eunicid worms, with a ‘budding’ method of asexual reproduction (Reimer et al. 2011b).

Remarks. Specimens examined with a discontinuous mesogleal sphincter muscle (Swain et al. 2015).

The genus *Acrozoanthus* has a long and complicated taxonomic history. Reimer et al. (2011b) discussed that “despite its very similar appearance, *Acrozoanthus australiae* was placed into a genus separate from *Zoanthus* due to the presence of an axial skeleton

(Saville-Kent 1893). Later it was shown that this skeleton was in fact a result of habitat preference as *Acrozoanthus* inhabits the outside of eunicid worm tubes, and the genus was subsequently merged back again into the genus *Zoanthus* (Haddon 1895). Subsequent to its original description, this species was not mentioned in literature again until its rediscovery by Ryland (1997), based on examination of a single specimen. Further work by Ryland and co-workers described the nematocysts of *Acrozoanthus australiae* (Ryland et al. 2004) and also an unusual ‘budding’ method of asexual reproduction (Ryland 1997), which was theorized to potentially confirm the placement of this species in its own genus”.

Molecular phylogenetic data indicate that *Acrozoanthus* is within the *Zoanthus* (see Reimer et al. 2011b) monophyly. However, due to a need for more detailed investigations, we refrain from synonymising *Acrozoanthus* with *Zoanthus*.

***Isaurus* Gray, 1828**

Isaura Lamouroux, in Audouin, Bourdon, de Candolle, d’Aubebard de Férussac, Deshayes, Deslongchamps, É. Geoffroy Saint-Hilaire, I. Geoffroy Saint-Hilaire, Guérin, Guillemain, de Jussieu, Kunth, Delafosse, Lamouroux, Latreille, Prévost, Richard and Bory de Saint-Vincent, 1826: 23 [*nomen oblitum*].

Isaurus Gray, 1828: 8 [*nomen protectum*]

Isaura Agassiz, 1845: 14 [unjustified emendation].

Antinedia Duchassaing de Fombressin and Michelotti, 1866: 136.

Pales Gray, 1867: 8.

Panceria Andres, 1877: 221–226.

Polythoa (*Monothoa*) Andres, 1883a: 521.

Zoanthus (*Monanthus*) Andres, 1883a: 538, 549, 541, 543.

Isaua Volpi and Benvenuti, 2003: 72 [incorrect spelling].

Type species. *Isaurus tuberculatus* Gray, 1828, by monotypy.

Gender. Masculine.

Diagnosis. Zooxanthellate, absence of mineral incrustations in the column/coenenchyme (excluding superficial surface attachments) brachycnemic zoantharians with recumbent (non-erect) polyps. Often have tubercles (raised bumps = endodermal invagination) on the outer surface of polyps, except in *Isaurus maculatus* Muirhead and Ryland, 1985, which has a smooth polyp surface.

Remarks. Specimens examined with an orthogonally-reticulate mesogleal sphincter muscle (Swain et al. 2015).

Ryland and Lancaster (2003) recognised only three valid species in this genus, although 22 species have been described in or assigned to this genus.

Suborder Macrocnemina Haddon and Shackleton, 1891

Macrocneminae Haddon and Shackleton, 1891a: 626.

Diagnosis. Zoantharians with the fifth mesenteries from the dorsal directive being complete.

Remarks. As stated above, the two suborders of Zoantharia appear not to be monophyletic (Sinniger et al. 2005). Three families are currently assigned to Macrocnemina Haddon and Shackleton, 1891.

Epizoanthidae Delage and Hérouard, 1901

Mardoellidae Danielssen, 1890: 116, 117 [*nomen oblitum*].

Epizoanthinae Delage and Hérouard, 1901: 664 [*nomen protectum*].

Type genus. *Epizoanthus* Gray, 1867.

Diagnosis. Macrocnemic zoantharians with a simple mesogleal muscle (Sinniger and Häussermann 2009: 26).

Remarks. The family-group Epizoanthidae (as “Epizoanthinae”) was established by Delage and Hérouard (1901: 664, 665) as monotypic, with the inclusion of only the type genus. Three genera are currently assigned to Epizoanthidae.

Recent molecular and morphological studies (e.g. Swain et al. 2015) suggest that this diagnosis should be revised following a complete revision of the suborder. Currently, molecular signatures such as those suggested in Sinniger et al. (2008) and in Sinniger et al. (2013) appear efficient to distinguish macrocnemic genera.

***Epizoanthus* Gray, 1867**

Sidisia Gray, 1858: 489 [suppressed in Opinion 1689, ICZN 1992].

Epizoanthus Gray, 1867: 237 [conserved in Opinion 1689, ICZN 1992].

Carolia Gray, 1867: 239 [invalid name, junior homonym].

Polythoa (*Endeithoa*) Andres, 1883a: 521, 531.

Verrillia Andres, 1883a: 520, 545.

Zoanthus (*Corticanthus*) Andres, 1883a: 538, 541.

Mardoell Danielssen, 1890: 117–126.

Marodellia Blanchard, 1893: 130 [unjustified emendation]

Mardoella Bell, 1906: 762 [incorrect spelling].

Lirrevia Delphy, 1939: 270.

Type species. *Dysidea papillosa* Johnston, 1842, by monotypy (see also Opinion 1689, ICZN 1992).

Gender. Masculine.

Diagnosis. As for family but readily distinguishable from *Palaeozoanthus* by the presence of non-fertile micromesenteries (Sinniger and Häussermann, 2009: 26).

Remarks. Most species with reticulate mesogleal muscle, *E. illoricatus* Tischbierék, 1930 with simplified mesogleal sphincter muscle (Swain et al. 2015).

As discussed in Sinniger and Häussermann (2009: 26), this genus is characterised by the “[p]olyps usually strongly encrusted with sand particles. Species found on rocky substrata or gastropod shells often inhabited by pagurids; some cases of free-living species reported (*E. lindhali*, *E. vagus*). In colonial species, polyps linked by stolons or, in pagurid-associated species, by a continuous coenenchyme. No symbioses with *Symbiodinium zooxanthellae*”.

***Palaeozoanthus* Carlgren, 1924**

Palaeozoanthus Carlgren, 1924: 470–473.

Type species. *Paleozoanthus reticulatus* Carlgren, 1924, by original description and monotypy.

Gender. Masculine.

Diagnosis. Macrocnemic zoantharians with a simple mesogleal muscle, and fertile micromesenteries.

Remarks. This genus is monospecific, and is comprised of the type species *Paleozoanthus reticulatus*, which has not been encountered since it was first described. Due to similarities in sphincter muscles (Swain et al. 2015) further studies are needed to determine if this genus corresponds to *Terrazoanthus*.

***Thoracactis* Gravier, 1918**

Thoracactis Gravier, 1918: 12.

Thoracactus Walsh, 1967: 49 [unjustified emendation and junior objective synonym].

Toracactis Herberts, 1972: 80 [incorrect spelling].

Type species. *Thoracactis topsenti* Gravier, 1918, by monotypy.

Gender. Masculine.

Diagnosis. Rudimentary sphincter muscles, azooxanthellate, with no mesogleal channels or lacunae, found on hexactinellid sponges.

Remarks. This is a monospecific genus comprised of only the type species *Thoracactis topsenti*. In describing *Thoracactis topsenti*, Gravier (1918) incorrectly identified it as an actinian (anemone) based on the lack of zooxanthellae, channels, gaps, or cell islets. Although currently placed in the family Epizoanthidae, based on the current

understanding of the type species, this genus is referable to the family Parazoanthidae, although an examination of the type material will be necessary to confirm this (see Reimer et al. 2010a: 158).

Hydrozoanthidae Sinniger, Reimer and Pawlowski, 2010

Hydrozoanthidae Sinniger, Reimer and Pawlowski, 2010: 60.

Type genus. *Hydrozoanthus* Sinniger, Reimer and Pawlowski, 2010.

Diagnosis. “This family is erected to group former Parazoanthidae species sharing specific insertions and deletions in mt-16S rDNA, especially in the V5 region (as defined in Sinniger *et al.*, 2005) of this gene” and “Phylogenetically, species in this family are more closely related to brachycnemic zoanthids (especially from the genus *Palythoa*) than to other parazoanthids.” (Sinniger et al. 2010a: 61).

Remarks. “This family groups several tropical and sub-tropical macrocnemic zoanthids; including species associated with hydrozoans and also several other non-hydrozoan associated species.” (Sinniger et al. 2010a: 61). This family is currently comprised of two genera— *Hydrozoanthus* Sinniger, Reimer and Pawlowski, 2010, and *Terrazoanthus* Reimer and Fujii, 2010.

***Hydrozoanthus* Sinniger, Reimer and Pawlowski, 2010**

Hydrozoanthus Sinniger, Reimer and Pawlowski, 2010: 60.

Type species. *Parazoanthus tunicans* Duerden, 1900, by original designation.

Gender. Masculine.

Diagnosis. A hydrozoanthid associated with hydrozoans.

Remarks. Examined species with branchiform endodermal sphincter muscle (Swain et al. 2015).

***Terrazoanthus* Reimer and Fujii, 2010**

Terrazoanthus Reimer and Fujii, 2010: 20.

Type species. *Terrazoanthus onoi* Reimer and Fujii, 2010, by original designation.

Gender. Masculine.

Diagnosis. This genus is characterised by being a member of the Hydrozoanthidae that is found on rocky substrates (as opposed to being obligate symbionts with hydrozoans). Some species in this genus are also brightly coloured (see Reimer and Fujii 2010: 20).

Remarks. Sphincter muscle transitional, with distal half mesogleal and proximal half endodermal, with encrustations to endodermal surface of mesoglea (Swain and Swain 2014) (=meso-endo transitional [Swain et al. 2015]), although *T. minutus* (Duerden, 1898) has a simplified mesogleal sphincter muscle (Swain et al. 2015).

Described species currently referable to the genus *Terrazoanthus* are mainly from the East Pacific, with *T. minutus* from the Caribbean, and it is likely that several more undescribed species exist in the Atlantic (see Reimer et al. 2010a, 2012a) and in the Central Indo-Pacific region (Reimer et al. 2014b).

The diagnosis of *Terrazoanthus* is in need of revision with the placement of *T. patagonichus* (Carlgren, 1898) into this genus by Swain et al. (2015) as this species is associated with hydroids (McMurrich 1904).

Microzoanthidae Fujii and Reimer, 2011

Microzoanthidae Fujii and Reimer, 2011: 420, 421.

Type genus. *Microzoanthus* Fujii and Reimer, 2011.

Diagnosis. As discussed in Fujii and Reimer (2011: 421), this family is characterised by “[c]olonies attached to bottom side (downward facing side) of dead coral rubble, asperous stones, inside narrow cracks, or occasionally on dead coral rubble on muddy seafloor. Azooxanthellate, macrocnemic. Polyps connected by narrow stolon or solitary. Sand particles encrusted in column. Irregularly sized sand particles encrusted into ectoderm. Tentacles two to three times as long as expanded oral disc diameter. Edge of oral disc shaped in regular, repeating zig-zagged pattern”.

Remarks. This is a monotypic family and comprises only the genus *Microzoanthus* Fujii and Reimer, 2011, with two species.

***Microzoanthus* Fujii and Reimer, 2011**

Microzoanthus Fujii and Reimer, 2011: 420, 421.

Type species. *Microzoanthus occultus* Fujii and Reimer, 2011, by original designation.

Gender. Masculine.

Diagnosis. As for family (Fujii and Reimer 2011). Encircling sinus present just beneath ectodermal surface (Swain and Swain 2014).

Remarks. Examined specimens with spindly-cteniform endodermal sphincter muscle (Swain et al. 2015).

Currently two species, reported only from the Pacific Ocean (Fujii and Reimer 2013) and the Red Sea (Reimer et al. 2014c).

Nanozoanthidae Fujii and Reimer, 2013

Nanozoanthidae Fujii and Reimer, 2013: 512.

Type genus. *Nanozoanthus* Fujii and Reimer, 2013.

Diagnosis. “Well developed polyps connected by narrow stolon. Mineral particles encrusted in column from aboral end to the edge of the oral disc. Irregularly sized sand particles encrusted into ectoderm and slightly into mesoglea. Zig-zagged, white-colored pattern following outside edge of oral disc. Macrocnemic mesenterial arrangement. Sphincter muscle mesogleal. No lacunae or ring sinus. Zooxanthellate. Mitochondrial cytochrome oxidase subunit I and 16S ribosomal DNA sequences significantly differ from all other known zoanthid genera (Fig. 1, 2).” (Fujii and Reimer 2013).

Remarks. A monogeneric family. Molecular data position this family in an intermediate position between the Brachycnemina and Macrocnemina, although currently it is placed within Macrocnemina (Fujii and Reimer 2013).

***Nanozoanthus* Fujii and Reimer, 2013**

Nanozoanthus Fujii and Reimer, 2013: 512–515.

Type species. *Nanozoanthus harenaceus* Fujii and Reimer, 2013, by original designation.

Gender. Masculine.

Diagnosis. As for family above.

Remarks. This is a monospecific genus currently, with specimens reported from southern Japan, Western Australia, and the Red Sea (Reimer et al. 2015a). However, molecular data indicate that the European species *Isozoanthus sulcatus* (Gosse, 1859) likely also belongs to this genus (Fujii and Reimer 2013).

Parazoanthidae Delage and Hérouard, 1901

Savaliidae (as “Savalini”) Nardo, 1844: 433.

Bergidae Verrill, 1865a: 147 [*nomen oblitum*].

Gerardidae Verrill, 1865a: 148.

Savagliidae Brook, 1889: 51, 74, 79.

Parazoanthidae Delage and Hérouard, 1901: 665 [*nomen protectum*].

Savaliidae Poche, 1914: 104.

Heterozoanthidae Pax and Müller, 1956: 3.

Type genus. *Parazoanthus* Haddon and Shackleton, 1891.

Diagnosis. As discussed in Sinniger and Häussermann (2009: 28), this family “[...] traditionally groups macrocnemic zoanthids possessing an endodermal sphincter. Mem-

ber species are frequently associated with other organisms, which are used as substrata”. Excludes species that form monophyly with Brachycnemina (Sinniger et al. 2010a).

Remarks. Precedence of Bergiidae Verrill, 1865, and *Bergia* Duchassaing de Fombressin and Michelotti, 1860, and respectively Parazoanthidae Delage and Hérourard, 1901, and *Parazoanthus* Haddon and Shackleton, 1891a, was reversed in Low and Reimer (2011a). In accordance with Article 23.9.2 (ICZN 1999: 28, 29), an application is being prepared to request that the International Commission on Zoological Nomenclature suppress the senior subjective synonyms Savaliidae Nardo, 1844, Gerardiidae Verrill, 1865, and Savagliidae Brook, 1889, in favour of Parazoanthidae Delage and Hérourard, 1901, to maintain current and widespread usage. See additional discussion in Appendix 3.

In addition to the type genus, *Parazoanthus* Haddon and Shackleton, 1891, twelve other valid genera are currently assigned to the family Parazoanthidae: *Antipathozoanthus* Sinniger, Reimer and Pawlowski, 2010, *Bergia* Duchassaing de Fombressin and Michelotti, 1860, *Bullagummizoanthus* Sinniger, Ocaña and Baco, 2013, *Corallizoanthus* Reimer, in Reimer, Nonaka, Sinniger and Iwase, 2008, *Hurlizoanthus* Sinniger, Ocaña and Baco, 2013, *Isozoanthus* Carlgren, in Chun, 1903, *Kauluzoanthus* Sinniger, Ocaña and Baco, 2013, *Kulamanamana* Sinniger, Ocaña and Baco, 2013, *Mesozoanthus* Sinniger and Häussermann, 2009, *Savalia* Nardo, 1844, *Umimayanthus* Montenegro, Sinniger and Reimer, 2015, and *Zibrowius* Sinniger, Ocaña and Baco, 2013.

***Parazoanthus* Haddon and Shackleton, 1891**

Heterozoanthus Verrill, 1870: 371 [*nomen oblitum*].

Parazoanthus Haddon and Shackleton, 1891a: 653, 654 [*nomen protectum*].

Type species. *Palythoa axinella* Schmidt, 1862, by original designation.

Gender. Masculine.

Diagnosis. Originally described as well-developed canal system in the mesoglea of the column, forming a ring sinus. Zoantharians often associated with sponges but not Hydrozoa, lacking skeletal secretion (Sinniger et al. 2010a). Examined species with endodermal sphincter muscle (=branchiform endodermal muscle [Swain et al. 2015]), encrustations reaching to endodermal surface of mesoglea (Swain and Swain 2014).

Remarks. “The original morphological description of *Parazoanthus* mentions several characteristics such as diffuse endodermal sphincter, encircling sinus, endodermal canals, lacunae and cell-islets in the mesoglea, continuous ectoderm and body-wall incrustated with mineral particles, often with numerous sponge spicules present in the incrustations. As shown in Sinniger et al. (2005) and here, these morphological characteristics alone do not ascertain the monophyly of *Parazoanthus*. Morphological characteristics in zoanthids can often become artifactual due to both complications

encountered in making thin cuttings of heavily sediment incrusting polyps, and in interpreting the results of such sections. In the past, the large majority of epizoic macrozoanths were described as belonging to *Parazoanthus* despite clearly different ecologies in many cases.

Thus, the results of this study strongly suggest that only zoanthid species able to associate with sponges should remain in *Parazoanthus*, as the type species of this genus, *P. axinellae* from the Mediterranean Sea, is regularly associated with demosponges.” (Sinniger et al. 2010a: 69).

There is a need for a new diagnosis of this genus-grouping. With the recent erection of *Umimayanthus* Montenegro, Sinniger and Reimer, 2015 and the resurrection of *Bergia* Duchassaing de Fombressin and Michelotti, 1860 in Montenegro et al. (2015a), the genus *Parazoanthus* now consists only of the former phylogenetic grouping of *Parazoanthus* ‘clade C’ sensu Sinniger et al. (2010a) and is monophyletic (Montenegro et al. 2015a: 71). *Parazoanthus* can be distinguished from *Bergia* and *Umimayanthus* by 16S-rDNA sequences, lacking the unique 60 bp deletion of *Bergia* and the unique insertion and deletion of *Umimayanthus*. Thus, the molecular characters described in Sinniger et al. (2008, 2013) appear to be efficient in identifying to genus level and could be used as diagnostic characters.

***Antipathozoanthus* Sinniger, Reimer and Pawlowski, 2010**

Antipathozoanthus Sinniger, Reimer and Pawlowski, 2010: 61.

Type species. *Gerardia macaronesicus* Ocaña and Brito, 2003, by original designation.

Gender. Masculine.

Diagnosis. Sinniger et al. (2010a: 63) originally diagnosed this genus as a group that “grows exclusively on antipatharians” and lacking skeletal secretion.

Remarks. No mesogleal canals or sinus, encrustation to outer mesoglea (Swain and Swain 2014), examined species with either branchiform endodermal or endo-meso transitional sphincter muscle (Swain et al. 2015).

Sinniger et al. (2010a: 63) discussed that “[t]he type species *A. macaronesicus* was originally included in the description of *Savalia* (*Gerardia*) *macaronesica* (Ocaña and Brito 2003), and later the description was amended and the authors suggested the possible placement of this species in a separate genus (Ocaña et al. 2007). The species name was accorded to the genus gender. Skeletal secretion (similar to *Savalia* spp.) was advanced by Ocaña and Brito (2003) as occurring in *Antipathozoanthus macaronesicus*, and this remains to be studied in detail in order to assess whether this is an isolated characteristic or if it is taxonomically informative at genus level.”

It also appears at least one member of this genus can be found on gorgonian octocorals (Bo et al. 2012).

***Bergia* Duchassaing de Fonbressin and Michelotti, 1860**

Bergia Duchassaing de Fonbressin and Michelotti, 1860: 54.

Type species. *Bergia catenularis* Duchassaing de Fonbressin and Michelotti, 1860, by subsequent designation by Duerden (1903: 496).

Gender. Feminine.

Diagnosis. "... can be distinguished from all other zoantharians including *Parazoanthus* spp., *Umimayanthus* spp. and *Epizoanthus* spp. by a unique deletion of 60 bp (from position 133 to 192 in our alignment) and several consecutive base substitutions in the 16S-rDNA region. These characters clearly separate this genus from all other genera inside the family Parazoanthidae, as well as from the genus *Epizoanthus*" (Montenegro et al. 2015a: 68).

Remarks. Long considered to be a junior subjective synonym of *Parazoanthus* Haddon and Shackelton, 1891, recent molecular and morphological work by Montenegro et al. (2015a) have shown that the type species, *Bergia catenularis* Duchassaing de Fonbressin and Michelotti, 1860, represents a generic-level monophyly and resurrected the genus-group name *Bergia* Duchassaing de Fonbressin and Michelotti, 1860, for this grouping.

This genus-grouping currently contains three species all found in the Atlantic Ocean, although there is evidence of undescribed species in the Pacific Ocean (Montenegro et al. 2015a: 68).

Examined species in this genus-grouping have either branchiform endodermal or simplified mesogleal sphincter muscles (Swain et al. 2015).

***Bullagummizoanthus* Sinniger, Ocaña and Baco, 2013**

Bullagummizoanthus Sinniger, Ocaña and Baco, 2013: [9].

Type species. *Bullagummizoanthus emilyacadiaarum* Sinniger, Ocaña and Baco, 2013, by original designation.

Gender. Masculine.

Diagnosis. Characteristic insertion/deletion pattern in the 16S V5 region sensu Sinniger et al. (2005) (Sinniger et al. 2013).

Remarks. Monospecific deep-sea genus, appears to be specifically epibiotic on paragorgiid octocorals, and described from and only reported from the Hawaiian Archipelago, although likely present throughout the Pacific (Sinniger et al. 2013).

***Corallizoanthus* Reimer, in Reimer, Nonaka, Sinniger and Iwase, 2008**

Corallizoanthus Reimer, in Reimer, Nonaka, Sinniger and Iwase, 2008: 940.

Type species. *Corallizoanthus tsukaharai* Reimer, in Reimer, Nonaka, Sinniger and Iwase, 2008, by monotypy.

Gender. Masculine.

Diagnosis. Characterised by its association with living precious corals (Alcyonacea: Coralliidae), and unlike the gorgonian-associated genus *Savalia*, does not secrete its own scleroproteinous axis. Additionally, polyps are primarily but not always unitary (solitary; non-colonial) (see Reimer et al. 2008a: 940). Encrustations to center of mesoglea, sphincter muscle is cyclically transitional (Swain and Swain 2014, Swain et al. 2015).

Remarks. A monospecific genus thus far only reported from the Ryukyu Islands, Japan.

***Hurlizoanthus* Sinniger, Ocaña and Baco, 2013**

Hurlizoanthus Sinniger, Ocaña and Baco, 2013: [7].

Type species. *Hurlizoanthus parrishi* Sinniger, Ocaña and Baco, 2013, by original designation.

Gender. Masculine.

Diagnosis. Macrocnemic genus associated with primnoids. Characteristic insertion/deletion pattern in the 16S V5 region sensu Sinniger et al. (2005) (Sinniger et al. 2013).

Remarks. Currently this deep-sea genus includes only one species, known from a few locations in the Hawaiian Archipelago (Sinniger et al. 2013).

***Isozoanthus* Carlgren, in Chun, 1903**

Polythoa (*Taeniothoa*) Andres, 1883a: 521, 532 [*nomen oblitum*].

Isozoanthus Carlgren, in Chun, 1903: 520 [*nomen protectum*].

Type species. *Isozoanthus giganteus* Carlgren, in Chun, 1903, by monotypy (Articles 68.3, 68.3.1, ICZN, 1999: 71).

Gender. Masculine.

Diagnosis. Macrocnemic zoanthids with a marginal endodermal sphincter muscle (=cteniform endodermal [Swain et al. 2015]) and inconspicuous mesogleal ring-sinus.

Remarks. The genus *Isozoanthus* was first made available in Carlgren (in Chun 1903: 520). The type species by monotypy is *Isozoanthus giganteus* (first very briefly diagnosed and figured by Carlgren (in Chun, 1903: 520, unnumbered fig.). Carlgren's (in Nordgaard 1905: 159) use of "*Isozoanthus* (*Epizoanthus*) *arborescens*" and subsequent designation (i.e. Carlgren 1913: 39) of *Epizoanthus arborescens* (Danielsen, 1890) as type species are thus invalid (Article 67.2, ICZN, 1999: 66, 67) (see also Williams 2000: 195).

The status of this genus is currently very confused. With the utility of the characters of the sphincter muscle in zoantharians in question (see Swain 2010, Sinniger et al. 2010a), it is clear that more research is needed to clarify the diagnosis and taxonomic position of *Isozoanthus*. The taxonomy is further complicated by the recently described species (the hydroid-associated *Hydrozoanthus antumbrosus* (Swain, 2009) originally described within *Isozoanthus*), and the octocoral-associated *Isozoanthus primonodius* Carreiro-Silva, Braga-Henriques, Sampaio, de Matos, Porteiro and Ocaña, 2010) clearly belong to other genera based on ecology and morphology. Furthermore, only limited molecular data is available for the type species *Isozoanthus giganteus*. Data in Swain (2010) indicate that *Isozoanthus giganteus* is highly divergent from both the well-researched species of *Isozoanthus*—*Isozoanthus sulcatus* (Gosse, 1859)—and all known zoantharians. Recent work by Fujii and Reimer (2013) shows that *I. sulcatus* is likely within the Nanazoanthidae Fujii and Reimer, 2013. However, as additional information on *Isozoanthus giganteus* is lacking, the diagnosis is retained with the caveat that any assignment of species to this genus should include: 1) phylogenetic confirmation of a close relationship with *Isozoanthus giganteus*; and 2) the elimination of any possibility that the species in question does not belong to another parazoanthid genera based on morphology, ecological associations, and/or habitat.

***Kauluzoanthus* Sinniger, Ocaña and Baco, 2013**

Kauluzoanthus Sinniger, Ocaña and Baco, 2013: [8].

Type species. *Kauluzoanthus kerbyi* Sinniger, Ocaña and Baco, 2013, by original designation.

Gender. Masculine.

Diagnosis. Polyps do not contract when fixed. Characteristic insertion/deletion pattern in the 16S V5 region sensu Sinniger et al. (2005) (Sinniger et al. 2013).

Remarks. Currently this genus comprises only one species, which was reported as parasitic on *Kulamanana haumea*.

***Kulamanamana* Sinniger, Ocaña and Baco, 2013**

Kulamanamana Sinniger, Ocaña and Baco, 2013: 4.

Type species. *Kulamanamana haumaeae* Sinniger, Ocaña and Baco, 2013, by original designation.

Gender. Feminine.

Diagnosis. Macrocnemic genus associated with octocorals and secreting a golden to dark brown scleroproteic skeleton. Ectoderm absent of mineral particles, with well-

developed coenenchyme completely covering the host. Characteristic insertion/deletion pattern in the 16S V5 region sensu Sinniger et al. (2005) (Sinniger et al. 2013).

Remarks. The type species has been reported to live primarily on isidid corals (bamboo corals) (Sinniger et al. 2013: 6). Reported from the Hawaiian Archipelago, Line and Jarvis Islands, Palmyra Atoll, Kingman Reef, all in the Pacific (Sinniger et al. 2013).

***Mesozoanthus* Sinniger and Häussermann, 2009**

Mesozoanthus Sinniger and Häussermann, 2009: 31, 32.

Type species. *Mesozoanthus fossii* Sinniger and Häussermann, 2009, by original designation and monotypy.

Gender. Masculine.

Diagnosis. “Macrocnemic with *Parazoanthus*-like growth-form. Well-developed polyps with long and pointed tentacles; polyps form clusters linked by a basal coenenchyme. DNA sequences significantly differ from those in other genera...” and “In contrast to *Parazoanthus*, members of *Mesozoanthus* usually occur in small patches and are not known to colonise demosponges. No symbioses with *Symbiodinium* zooxanthellae.” (Sinniger and Häussermann 2009: 32).

Remarks. Only two species of this genus are known, from temperate waters along the west coast of the Americas.

***Savalia* Nardo, 1844**

Savalia Nardo, 1844: 433, 434.

Savaglia Nardo: 1877: 674–676 [unjustified emendation].

Gerardia Lacaze-Duthiers, 1864a: 87.

Type species. *Gorgonia savaglia* Bertolini, 1819, by monotypy (Articles 68.3, 68.3.1, ICZN 1999: 71).

Gender. Masculine.

Diagnosis. No mesogleal canal system in the column. Secreting a black or dark brown horny skeleton, azooxanthellate.

Remarks. Examined species in this genus-grouping have branchiform endodermal or cyclical transitional sphincter muscle (Swain et al. 2015). Distinction from the other skeleton-secreting zoantharians such as *Kulamanamana* or potentially *Antipathozoanthus* can be made through habitat (shallow vs. deep-sea) and/or molecular signatures.

There has been historical and recent controversy over the correct name for this genus-group, and this is discussed in detail in Appendix 3.

***Umimayanthus* Montenegro, Sinniger and Reimer, 2015**

Umimayanthus Montenegro, Sinniger and Reimer, 2015: 76.

Type species. *Umimayanthus chanpuru* Montenegro, Sinniger and Reimer, 2015, by original designation.

Gender. Masculine.

Diagnosis. "...can be distinguished from all zoantharians including *Parazoanthus* spp. by a highly conservative and unique insertion of 9 bp in length (from position 556 to 564 in alignment) and one deletion of 14 bp long (from position 574 to 587) in the mt 16S-rDNA region" (Montenegro et al. 2015b: 76).

Remarks. Specimens examined from this genus-grouping have a branchiform endodermal sphincter muscle (Swain et al. 2015).

"...exclusively associated with sponges, usually encrusting and cushion sponges, occasionally with massive sponges. Polyps generally scattered over the sponge surface, but can form defined stoloniferous chains in lines, or form groups of two to three connected polyps. Polyps may be solitary or connected to each other by a stolon through a thin but clearly visible coenenchyme either over or under the sponge surface. Polyps with sand particles and detritus incrustated in column. Tentacles equal or longer than the expanded oral disc diameter." (Montenegro et al. 2015b: 76).

This genus-group currently includes four described species; three in the Indo-Pacific and one in the Atlantic (Montenegro et al. 2015b).

***Zibrowius* Sinniger, Ocaña and Baco, 2013**

Zibrowius Sinniger, Ocaña and Baco, 2013: [7].

Type species. *Zibrowius ammophilus* Sinniger, Ocaña and Baco, 2013, by original designation.

Gender. Masculine.

Diagnosis. "Sand incrustated, arborescent fan shaped colonies, golden skeleton, well developed coenenchyme completely covering the host, can be confused with *Kulamanamana*, but easily distinguished by the presence of sand incrustation in the ectoderm, characteristic insertion/deletion pattern in the 16S V5 region" sensu Sinniger et al. [2005] (Sinniger et al. 2013: 7)

Remarks. Until now, only reported from the Cross Seamount in the central Pacific Ocean.

Suborder *incerta sedis*

Remarks. Two families are currently not assigned to any suborder.

Abyssoanthidae Reimer and Fujiwara, in Reimer, Sinniger, Fujiwara, Hirano and Maruyama, 2007

Abyssoanthidae Reimer and Fujiwara, in Reimer, Sinniger, Fujiwara, Hirano and Maruyama, 2007: 258.

Type genus. *Abyssoanthus* Reimer and Fujiwara, in Reimer, Sinniger, Fujiwara, Hirano and Maruyama, 2007.

Diagnosis. “Sand/detritus/sediment-encrusted Zoantharia with unitary (noncolonial) free-living polyps, attached to hard substrates at abyssal (non-continental shelf deep-sea) depths surrounding methane cold seeps or other chemosynthetic ecosystems.” Reimer and Sinniger (2010: 454).

Remarks. A monogeneric family. Molecular data places this family in an unresolved position distant from the Brachycnemina and Macrocnemina.

***Abyssoanthus* Reimer and Fujiwara, in Reimer, Sinniger, Fujiwara, Hirano and Maruyama, 2007**

Abyssoanthus Reimer and Fujiwara, in Reimer, Sinniger, Fujiwara, Hirano and Maruyama, 2007: 258.

Type species. *Abyssoanthus nankaiensis* Reimer and Fujiwara, in Reimer, Sinniger, Fujiwara, Hirano and Maruyama, 2007, by original designation and monotypy.

Gender. Masculine.

Diagnosis. As for family above.

Remarks. Two species known, both from waters around Japan. There may be additional species from the Mediterranean, in non-chemosynthetic environments (Sinniger et al. 2010b).

Family *incerta sedis****Stephanidium* Hertwig, 1888**

Stephanidium Hertwig, 1888: 52.

Type species. *Stephanidium schulzii* Hertwig, 1888, by monotypy.

Gender. Neuter.

Diagnosis. Very small (diameter 1.5–2.2 mm, height 1 mm), unitary polyps, with microcnemes and macrocnemes, although their arrangement could not be clearly seen. Mesenteriel insertions make body wall to have a furrowed appearance, with spherical evaginations on the body wall above the area where the sphincter muscle is present. Twenty-six mesenteries.

Remarks. From the original description, possibly a species of zoantharian, but type material needs to be located and examined. We make no decision as to the validity of this genus and species in the event that the identification of this genus and species requires a reversal of precedence Article 23.9 (ICZN 1999: 27, 28) with a later (but more widely-used name). Also, the genus-group name *Stephanidium* Hertwig, 1888, is preoccupied by *Stephanidium* Ehrenberg, 1839 (*incerta sedis*).

Discussion

As can be seen from examining the nomenclatural and taxonomic history of the various supraspecific names in this paper, many taxa of the order Zoantharia have a confused history. However, over the past two decades, phylogenetic and detailed morphological examinations of zoantharians have resulted in a new understanding of the evolutionary relationships within the order (Sinniger et al. 2005, Reimer et al. 2007a, 2008a, Reimer and Fujii 2010, Sinniger et al. 2010a, 2013, Fujii and Reimer 2011, 2013; Swain et al. 2015). Combined with a recent effort to organize the nomenclature of zoantharians (Low and Reimer 2011a, b, 2012a, b), it can be said that much of the Zoantharia nomenclature is now stable, and generally reflects our current understanding of their evolutionary history.

However, as seen in this manuscript, there are still some nomenclatural issues that remain to be resolved. For example, the validity of the genera *Sphenopus* and *Acrozoanthus* still have to be thoroughly examined. Furthermore, it is clear from recent molecular phylogenetic results (Fujii and Reimer 2013) that the taxonomy and nomenclature of the suborders are in urgent need of a revision. Finally, at the ordinal level, it appears that Zoantharia is much closer to the Actiniaria (sea anemones) than has been previously thought (Fujii and Reimer 2013). As these two orders have been speculated to be basal cnidarians (Kayal et al. 2013), after clarification of their evolutionary history, subsequent nomenclatural amendment may be needed.

Additional work examining the utility of ecological and morphological traits of Zoantharia as diagnostic characters is needed to allow the linkage of current phylogenetic results with past literature (Swain et al. 2016). Despite, or perhaps due to its confused and challenging morphological taxonomy, the order Zoantharia is now among the most advanced hexacorallian order in terms of the use of molecular tools to clarify phylogenetic relationships between taxa at various taxonomic levels. As such, and despite potential and known differences in molecular evolution of other anthozoans (Stampar et al. 2014), Zoantharia is a potentially good model for clarifying the taxonomy of other hexacorallian orders and anthozoan groups.

Acknowledgements

Carey Ashworth (University of the Ryukyus, Japan) is thanked for help with translating references; Anne-Marie Damiano and Catherine de la Bigne (Musée Océanographique,

Monaco) for providing literature. Siong-Kiat Tan (National University of Singapore) provided comments on nomenclatural issues. Sung-yin Yang (University of the Ryukyus, Japan) helped getting references. Saulo Bambi and Cecilia Volpi of the Museo di Storia Naturale Università di Firenze (sez. Zoologica “La Specola”), Italy, are thanked for their assistance in locating and photographing the syntype of *Gemmaria rusei* Duchassaing de Fombressin and Michelotti, 1860. Ray B. Williams (Tring, United Kingdom) provided information on the works of Angelo Andres and Philip Henry Gosse. Neal L. Evenhuis (Bishop Museum, Honolulu, Hawaii) provided copies of literature and provided comments on many of the dates of publications of the works cited. Daphne G. Fautin (University of Kansas) generously shared her bibliography (now published as Fautin 2016) of the Cnidaria (from the database “Hexacorallians of the World”) with us and her discussions with the first author through the years have greatly improved the content of this work. Peter K. L. Ng (National University of Singapore) has also provided guidance and advice on nomenclatural matters through the years. Helmut Zibrowius provided us with a tremendous amount of old literature on Zoantharia. Comments from Marymegan Daly (Ohio State University) and an anonymous reviewer on an earlier manuscript (that now forms part of Appendix 3) greatly improved the information presented therein. This work was sponsored in part by the International Research Hub Project for Climate Change and Coral Reef/Island Dynamics at the University of the Ryukyus, as well as by a JSPS Kiban B grant entitled “Global evolution of Brachycnema and their *Symbiodinium*” and by a Japan Society for the Promotion of Science (JSPS) ‘Zuno-Junkan’ grant entitled “Studies on origin and maintenance of marine biodiversity and systematic conservation planning” to the senior author.

References

The question of nomenclatural priority (while correctly a minor and largely theoretical exercise) underpins the system of nomenclature currently in use. In this order-wide review of the supraspecies nomenclature and taxonomy, we have had to determine the relative priority of many synonyms, and have come across accurate date of publication information for much of the literature. Many of these references have a nomenclatural impact beyond the order Zoantharia, and these data are included for the benefit of the taxonomic community at large.

- Abildgaard PC (1791) Nyere Efterretning om det Skaldyr fra Middelhavet, som Forskål har beskrevet under Navn af *Anomia tridentata*. Skrifter af Naturhistorie-Selskabet 1 (2): 171–175, pl. 10.
- Acosta A, Casas M, Vargas CA, Camacho JE (2005) Lista de Zoantharia (Cnidaria: Anthozoa) del Caribe y de Colombia. Biota Colombiana, 6 (2): 147–162.
- Agassiz L (1844) Nomina systematica generum polyporum (Anthozoorum et bryozoorum cum polythalmiis) tam viventium quam fossilium. Secundum ordinem alphabeticum disposita, adjunctis auctoribus, libris in quibus reperiuntur, anno editionis, etymologia et familiis ad

- qua pertinent. Jent et Gassmann, Soloduri [= Soloturn], vi + 28 + [4] pp. [31 Dec 1844 (Bowley and Smith, 1968: 35)]
- Agassiz L (1846) Nomenclatoris zoologici index universalis, continens nomina systematica classium, ordinum, familiarum et generum animalium omnium, tam viventium quam fossilium, secundum ordinem alphabeticum unicum disposita, adjectis homonymiis plantarum, nec non variis adnotationibus et emendationibus. Jent et Gassmann, Soloduri [= Solothurn], viii + 393 pp. [29 Dec 1846 (Evenhuis, 1997: 50)]
- Alencar DB, Melo AA, Silva GC, Lima RL, Pires-Cavalcante KMS, Carneiro RF, Rabelo AS, Sousa OV, Vieira RHSE, Viana FA, Sampaio AH, Saker-Sampaio S (2015) Antioxidant, hemolytic, antimicrobial, and cytotoxic activities of the tropical Atlantic marine zoanthid *Palythoa caribaeorum*. Anais da Academia Brasileira de Ciências 87 (2): 1113–1123. <https://doi.org/10.1590/0001-3765201520140370>
- Altuna Á, Sinniger F, Aldrey JM (2010) Occurrence of *Savalia savaglia* (Anthozoa: Zoantharia) in the Ría de Arousa (Galicia, north-western Spain, north-eastern Atlantic). Marine Biodiversity Records 3: e110. <https://doi.org/10.1017/s1755267210000965>
- Andres A (1877) On a new genus and species of zoanthina malacodermata (*Panceria spongiosa*, sp. n.). Quarterly Journal of Microscopical Science (n.s.) 17 (67): 221–226, pl. 16. [Jul 1877 (contents, p. v)]
- Andres A (1881) Prodromus neapolitanae actiniarum faunae addito generalis actiniarum bibliographiae catalogo. Mitteilungen aus der Zoologischen Station zu Neapel zugleich ein Repertorium für Mittelmeerkunde 2 (3): 305–371.
- Andres A (1883a) Le Attinie. Atti della Reale Accademia dei Lincei, ser. 3, Memorie della Classe di Scienze Fisiche, Matematiche e Naturali 14: 211–673, pls. 1–13. [> 1 Apr 1883 (Williams, 2012: 95); reissued as Andres (1883b)]
- Andres A (1883b) Le Attinie. Reale Accademia dei Lincei, Roma, x + 460 pp., pls. 1–13. [reissue of Andres (1883a) (Williams 2012: 96)]
- Andres A (1884) Le Attinie. Fauna und Flora des Golfes von Neapel, Wilhelm Engelmann, Leipzig, 9, i–x, 1–460, pls. 1–13. [> 30 Mar 1884 (?) (Williams 2012: 102); re-edition of Andres (1883a, b)]
- Anonymous (1786) The Natural History of many curious and uncommon Zoophytes, collected from various Parts of the Globe, by the late John Ellis, Esq. F.R.S. Soc. Reg. Upsal. Soc. Systematically arranged and described by the late Daniel Solander, M.D. F.R.S. &c. With Sixty-two Plates. 4to. 1l. 16s. in Boards. White. The Critical Review: Or, Annals of Literature 62 [Jul 1786]: 1–6.
- Anonymous (1842) Johnston's (G.) History of British Sponges and Lithophytes, 8vo, cloth, 1l 10s. In: New publications, from February 6 to March 6th, 1842. Bent's Monthly Literary Advertiser, Register of Books, Engravings, &c., 1842 (452) [11 Apr 1842]: 63.
- Anonymous (1850) Animaux radiaires des Antilles; par P. Duchassaing, D. M. In-8° de 2 feuilles, plus 2 pl. Imp. De Plon, à Paris. Bibliographie de la France, ou Journal général de l'Imprimerie et de la Librairie, et des Cartes géographiques, Gravures, Lithographies, Oeuvres de Musique 39 (32) [10 Aug 1850]: 401.
- Anonymous (1857) Suites a Buffon. Histoire naturelle des Coralliaires ou Polypes proprement dits (coraux, gorgones, éponges, etc., etc.), par MM. Milne-Edwards et J. Haime. Tomes

- 1^{er} et II, avec 2 livraisons de planches. Prix: figures noires, 19 fr., et fig. color., 25 fr. Feuilleton commercial du Journal Général de l'Imprimerie et de la Librairie (2) 1 (49) [5 Dec 1857]: 633–648.
- Anonymous (1893) The Great Barrier Reef of Australia, its Products and Potentialities, by W. Saville-Kent, F.L.S., F.Z.S., F.I.Inst., &c., illustrated. The Publishers' Circular 5 (1392) [4 Mar 1893]: 249.
- Anonymous (1897) The Naturalist in Australia. By W. Saville-Kent, F.L.S., F.Z.S., &c., Past President Royal Society of Queensland; Author of 'The Great Barrier Reef of Australia'. Illustrated by 50 full-page Collotypes, 9 Coloured Plates by Keulemans and other Artists, and over one hundred Illustrations in the Text. xv + 302 pp. The Saturday Review of Politics, Literature, Science and Art 83 [5 Jun 1897]: 644.
- Audouin JV (1826) Explication sommaire des planches de polypes de l'Égypte et de la Syrie, publiées par Jules-César Savigny, membre de l'Institut; offrant un exposé des caractères naturels des genres, avec la distinction des espèces. pp. 225–244. In: Description de l'Égypte ou Recueil des observations et des recherches qui ont été faites en Égypte pendant l'expédition de l'armée française, publié par les ordres de Sa Majesté l'empereur Napoléon le Grand. Histoire naturelle. Tome premier. Quatrième partie [Text]. Imprimerie Impériale, Paris, 336 pp. [authorship and date of publication (ICZN, 1987: 220; Sherborn, 1897: 287); reissued as Audouin (1828)]
- Audouin JV (1828) Explication sommaire des planches de polypes de l'Égypte et de la Syrie, publiées par Jules-César Savigny, membre de l'Institut; offrant un exposé des caractères naturels des genres, avec la distinction des espèces. pp. 40–78. In: Description de l'Égypte ou Recueil des observations et des recherches qui ont été faites en Égypte pendant l'expédition de l'armée française. Seconde édition dédiée au Roi. Tome vingt-troisième. Histoire naturelle. Zoologie. Animaux invertébrés (suite). Animaux vertébrés [Text]. C.L.F. Panckoucke, Paris, 456 pp. [reissue of Audouin (1826)]
- Audouin JV, Bourdon J-B-I, Brongniart A, Candolle A-P de, Férussac AE d'Audebard de, Deshayes G-P, Deslongchamps JAE, Geoffroy Saint-Hilaire É, Geoffroy Saint-Hilaire I, Guérin F-É, Guillemin JBA, Jussieu A de, Kunth KS, Delafosse G, Lamouroux J-V-F, Latreille P-A, Prévost C, Richard A, Bory de Saint-Vincent J-B-G-M (1826) Dictionnaire classique d'Histoire naturelle, Ouvrage dirigé par ce dernier collaborateur, et dans lequel on a ajouté, pour le porter au niveau de la science, un grand nombre de mots qui n'avaient pu faire partie de la plupart des Dictionnaires antérieurs. Tome neuvième. IO–MACIS. Rey et Gravier / Baudouin Frères, Paris, 596 pp. [25 Feb 1826 (Evenhuis 1997: 111)]
- Barel CDN, Kramers PGN (1977) A survey of the echinoderm associates of the north-east Atlantic area. Zoologische Verhandelingen Leiden 156 (1): 1–159.
- Bayer FM, Grasshoff M (1997) Case 2940. *Riisea* and *riisei* Duchassaing & Michelotti, 1860 (Cnidaria, Anthozoa): proposed conservation as the correct original spellings of generic and specific names based on the surname Riise. Bulletin of Zoological Nomenclature 54 (1): 11–13. [26 Mar 1997 (issue front cover)] <https://doi.org/10.5962/bhl.part.55>
- Bell FJ (1891) Contributions to our knowledge of the antipatharian corals. Transactions of the Zoological Society of London 13 (2): 87–92, pls. 11–12. [Apr 1891 (p. 87)]

- Bell FJ (1906) Crustacea, echinoderms, peripatus, worms, and Anthozoa. pp. 731–764. In: The History of the Collections Contained in the Natural History Departments of the British Museum. Vol. II. Separate Historical Accounts of the Several Collections Included in the Department of Zoology. By Order of the Trustees of the British Museum, London, [1] + 782 pp.
- Beneden E van (1897) Die Anthozoen der Plankton-Expedition. Ergebnisse der Plankton-Expedition der Humboldt-Stiftung. Lipsius & Tischer, Kiel and Leipzig, Band 2, 1–222 pp., pls. 1–16, 1 map. [Also published with French title “Les Anthozoaires de la ‘Plankton-Expedition’” in “*Résultats de la ‘Plankton-Expedition der Humboldt-Stiftung’*”]
- Bertolonii A (1819) Amoenitates italicae, sistentes opuscula ad rem herbariam et zoologiam Italiae spectantia. Bononiae, [1] + 472 pp., pls. 1–6. [Apr 1891 (Sayre 1959: 60)]
- Blainville HMD de (1830) Zoophytes, pp. 1–546. In: Cuvier G-F (Ed). Dictionnaire des Sciences Naturelles, dans lequel on traite méthodiquement des différens êtres de la nature, considérés soit en eux-mêmes, d’après l’état actuel de nos connoissances, soit relativement a l’utilité qu’en peuvent retirer la médecine, l’agriculture, le commerce et les arts. Suivi d’une biographie des plus célèbres naturalistes. Ouvrage destiné aux médecins, aux agriculteurs, aux commercans, aux artistes, aux manufacturiers, et à tous ceux qui ont intérêt à connoître les productions de la nature, leurs caractères génériques et spécifiques, leur lieu natal, leurs propriétés et leurs usages. Tome soixantième. ZOOPH–ZYT. F. G. Levrault, Strasbourg et Paris / Le Normant, Paris, [vi] + 631 pp. [Jun 1830 (Evenhuis 1997: 183)]
- Blainville HMD de (1834–1837) Manuel d’actinologie ou de zoophytologie. F. G. Levrault, Paris, viii + 695 pp. [1834 (pp. 1–644), 1837 (pp. 645–694) (Sherborn, 1922: xxv; 1932: 113)]
- Blanchard R (1892) Deuxième rapport sur la nomenclature des êtres organisés; rapport présenté au deuxième Congrès international de zoologie réuni à Moscou du 10/22 au 18/30 août 1892. Mémoires de la Société zoologique de France 6: 126–201.
- Bo M, Lavorato A, Di Camillo CG, Polisenio A, Baquero A, Bavestrello G, Irei Y, Reimer JD (2012) Black coral assemblages from Machalilla National Park (Ecuador). Pacific Science 66 (1): 63–81. <https://doi.org/10.2984/66.1.4>
- Bourne GC (1900) The Anthozoa. pp. 184. In: Lankester ER (Ed). A Treatise on Zoology. Part II. The Porifera and Coelentera. With an Introduction by E. Ray Lankester. Adam & Charles Black, London, vi + 178 + 81 + 84 + 25 pp. [> Aug 1900 (preface, p. vi)]
- Bowley DR, Smith HM (1968) The dates of publication of Louis Agassiz’s *Nomenclator Zoologicus*. Journal of the Society for the Bibliography of Natural History 5 (1): 35–36. <https://doi.org/10.3366/jsbnh.1968.5.1.35>
- Brandt JF (1835) Prodromus descriptionum animalium ab H. Mertensio in orbis terrarium circumnavigatione observatorum. Fascic. I. Polypos, acalephas discophoras et siphonophoras nec non echinodermata continens. Recueil des Actes de la Séance Publique de Académie Impériale des Sciences de St.-Petersbourg tenue le 29 décembre 1834, 200–273, [1]. [May 1835 (*verso* of title-page); also issued as a separate with pagination 1–73]
- Brockhaus FA (1862) Schmidt, Osc. Die Spongien des adriatischen Meeres. Mit 7 Kpfrtaf. Leipzig, 1862. Engelmann. Fol. VIII, 88 pp. Allgemeine Bibliographie, Monatliches Verzeichniss der wichtigern neuen Erscheinungen der deutschen und ausländischen Literatur, 1862 (7) [Jul 1862]: 96.

- Brook G (1889) Report on the Antipatharia collected by H.M.S. *Challenger*. In: Report of the Scientific Results of the Voyage of H.M.S. Challenger During the Years 1873–76, Under the Command of Captain George S. Nares, R.N., F.R.S. and the Late Captain Frank Tourle Thomson, R.N. Prepared Under the Superintendence of the Late Sir C. Wyville Thomson, Knt., F.R.S., &c. Regius Professor of Natural History in the University of Edinburgh Director of the Civilian Scientific Staff on Board and Now of John Murray, LL.D., Ph.D., &c. One of the the Naturalists of the Expedition. Zoology—Vol. XXXII. Her Majesty's Government, Edinburgh, part 80: 1–222, pls. 1–15. [8 Nov 1889 (Low and Evenhuis, 2013: 419)]
- Bruguère J-G (1791) Tableau encyclopédique et méthodiques des trois règnes de la nature. Contenant l'helminthologie, ou les vers infusoires, les vers intestins, les vers mollusques, &c. Pancoucke, Paris, viii + 84 pp., 95 pls. [30 Jul 1791 (Evenhuis, 2003a: 37)]
- Burnett WJ, Benzie JAH, Beardmore JA, Ryland JS (1994) High genetic variability and patchiness in a common Great Barrier Reef zoanthid (*Palythoa caesia*). *Marine Biology* 121 (1): 153–160. <https://doi.org/10.1007/BF00349484>
- Burnett WJ, Benzie JAH, Beardmore JA, Ryland JS (1995) Patterns of genetic subdivision in populations of a clonal cnidarian, *Zoanthus coppingeri*, from the Great Barrier Reef. *Marine Biology* 122 (4): 665–673. <https://doi.org/10.1007/BF00350688>
- Burnett WJ, Benzie JAH, Beardmore JA, Ryland JS (1997) Zoanthids (Anthozoa, Hexacoralia) from the Great Barrier Reef and Torres Strait, Australia: systematics, evolution and a key to species. *Coral Reefs* 16 (1): 55–68. <https://doi.org/10.1007/s003380050060>
- Calder DR, Stephens LD, Sanders AE (1992) Comments on the date of publication of John McCrady's hydrozoan paper *Gymnophthalmata of Charleston Harbor*. *Bulletin of Zoological Nomenclature*, 49 (4): 287–288. [17 Dec 1992 (issue front cover)]
- Cantraine F (1838) Notices sur un genre de la famille Ostracés. *Bulletins de l'Académie Royale des Sciences et Belles-Lettres de Bruxelles* 5 (3): 111–116, 1 pl.
- Carlgren O (1895) Ueber de Gattung *Gerardia* Lac.-Duth. *Öfversigt af Kongl. Vetenskaps-Akademiens Förhandlingar* 52 (5): 319–334.
- Carlgren O (1898) Zoantharien. In: *Ergebnisse der Hamburger Magalhaensische Sammelreise 1892/93*. Herausgegeben vom Naturhistorischen Museum zur Hamburg. I. Band. Allgemeines, Chordonier, Echinodermen und Coelenteraten. L. Friederichsen & Co., Hamburg, 48 pp., 1 pl.
- Carlgren O (1900) Ostafrikanische Actinien, gesammelt von Herrn Dr. F. Stuhlmann 1898 und 1899. *Mitteilungen aus dem Naturhistorischen Museum in Hamburg*, 17 (Supplement 2): 21–144, pls. 1–7. [28 Dec 1900 (title-page); also issued as separate with pagination pp. 1–124]
- Carlgren O (1905) Bottom Life. pp. 155–225. In: Nordgaard O (Ed). *Hydrographical and Biological Investigations in Norwegian Fiords. The Protist Plankton and the Diatoms in Bottom Samples*. John Grieg, Bergen, 254 pp., 21 pls.
- Carlgren O (1913) Zoantharia. *The Danish Ingolf Expedition* 5 (4): [1], 1–62, [1], pls. 1–7.
- Carlgren O (1923) Ceriantharia und Zoantharia der Deutschen Tiefsee-Expedition. *Deutsche Tiefsee-Expedition auf dem Dampfer „Valdivia“ 1898–1899* 19 (7): 243–338, pls. 24–32. [Also issued as a separate with pagination pp. 1–98, pls. 1–9]

- Carlgren O (1924) Die Larven der Ceriantharien, Zoantharien und Aktinarien, mit einem Anhang zu den Zoantharien. Wissenschaftliche Ergebnisse der Deutschen Tiefsee auf dem Dampfer „Valdivia“ 1898–1899 19 (8): 342–476, pls. 33–38.
- Carlgren O (1949) A survey of the Ptychodactiaria, Corallimorpharia and Actiniaria. Kungliga Svenska vetenskapsakademiens handlingar (4) 1 (1): 1–121. [24 Mar 1949 (p. 12)]
- Carlgren O (1954) Actiniaria and Zoantharia from South and West Australia with comments upon some Actiniaria from New Zealand. Arkiv för Zoologi (2) 6 (34): 571–595. [10 Aug 1954 (p. 595)]
- Carlos AA, Baillie BK, Kawachi M, Maruyama T (1999) Phylogenetic position of *Symbiodinium* (Dinophyceae) isolates from tridacnids (Bivalvia), cardiids (Bivalvia), a sponge (Porifera), a soft coral (Anthozoa), and a free-living strain. Journal of Phycology 35 (5): 1054–1062. <https://doi.org/10.1046/j.1529-8817.1999.3551054.x>
- Carreiro-Silva M, Braga-Henriques A, Sampaio I, Matos V de, Porteiro FM, Ocaña O (2011) *Isozoanthus primnoidus*, a new species of zoanthid (Cnidaria: Zoantharia) associated with the gorgonian *Callogorgia verticillata* (Cnidaria: Alcyonacea). ICES Journal of Marine Science 68 (2): 408–415. <https://doi.org/10.1093/icesjms/fsq073>
- Carter HJ (1870) On *Anthozoanthus parasiticus*, Deshayes, MS. (Algiers). Annals and Magazine of Natural History (4) 5 (30): 449. [1 Jun 1870 (Evenhuis 2003b: 25)]
- Cavallari DC, Gonçalves EP, Amaral VS do (2012) New occurrences of *Myoforceps aristatus* (Bivalvia: Mytilidae) in the Brazilian coast. Strombus 19 (1–2): 23–27.
- Cavolini F (1785) Memorie per servire alla storia d e' polipi marini. Napoli [= Naples], 115 pp. 4 pls.
- Cha H (2007) Systematics of the Order Corallimorpharia (Cnidaria: Anthozoa). PhD. Dissertation, University of Kansas, Lawrence, 196 pp.
- Chun C (1903) Aus den Tiefen des Weltmeeres. Schilderungen von der Deutschen Tiefsee-Expedition. Zweite umgearbeitete und Stark vermehrte Auflage. Gustav Fischer, Jena, ix + 592 pp., 482 figs., 46 pls., 3 maps. [15 Apr 1903 (Hallier 1903: 63)]
- Cornelius PF, Wells JW (1988) Ellis & Solander's 'Zoophytes', 1786: six unpublished plates and other aspects. Bulletin of the British Museum of Natural History (Hist. ser.), 16(1): 17–87.
- Costello MJ, Bouchet P, Boxshall G, Fauchald K, Gordon D, Hoeksema BW, Poore GCB, Soest RWM van, Stöhr S, Walter TC, Vanhoorne B, Decock W, Appeltans W (2013) Global coordination and standardisation in marine biodiversity through the World Register of Marine Species (WoRMS) and related databases. PLoS ONE 8 (1): e51629 [21 pp.]. <https://doi.org/10.1371/journal.pone.0051629>
- Couch RQ (1844) Cornish Fauna; Being a Compendium of the Natural History of the County, Intended to Form a Companion to the Collection in the Museum of the Royal Institution of Cornwall. Part 3: Zoophytes and Calcareous Corallines. L.E. Gillet, Truro, xvii + 164 + [8] pp., pls. 1–23. <https://doi.org/10.5962/bhl.title.87783>
- Cowie RH (1995) Catalog of the Native Land and Freshwater Molluscs of the Hawaiian Islands. Backhuys Publishers, Leiden, 248 pp.
- Cretella M (2010) The complete collation and dating of the section *Zoologie* of the *Coquille* voyage. Bolletino Malacologico 46 (2): 83–103.
- Cuvier G (1800) Classification des Zoophytes. [unnumbered tables 9, 10]. Leçons d'Anatomie comparée. Tome I. Contenant les organes du mouvement. Baudouin, Paris, 522 pp. [Feb 1800 (Duméril 1860: iv)]

- Cuvier G (1816) Les Zoanthes. (*Zoanthus*. Cuv.). p. 53. In: Le règne animal distribué d'après son organisation, pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée. Avec figures, dessinées d'après nature. Tome. IV, contenant les zoophytes, les tables, et les planches. Deterville, Paris, vii + 255 pp., pls. 1–15. ["1817", 2 Dec 1816 (Evenhuis 1997: 173)]
- Daly M, Brugler MR, Cartwright P, Collins AG, Dawson MN, Fautin DG, France SC, McFadden CS, Opresko DM, Rodriguez E, Romano SL, Stake JL (2007) The phylum Cnidaria: a review of phylogenetic patterns and diversity 300 years after Linnaeus. *Zootaxa* 1668: 127–182.
- Danielssen DC (1890) Actinida. In: Den Norske Nordhavs-Expedition 1876–1878. Zoologi. Grøndahl & Søn, Christiania, v + 184 pp., pls. 1–25, 1 map.
- Delage Y, Hérouard E (1901) Zoanthidés. – Zoanthidae. pp. 654–667, pls. 65. In: *Traité de Zoologie concrète*. Tome II – 2^{me} Partie. Les Coelentérés. C. Reinwald, Paris, x + [xi] + 848 pp., pls. 1–72. [< 15 Nov 1901 (Friedländer 1901: 579)]
- De la Cruz-Francisco V, González-González M, Morales-Quijano I (2016) Inventario taxonómico de Hydrozoa (orden Anthoathecata) y Anthozoa (subclases: Hexacorallia y Octocorallia) del Arrecife Enmedio, Sistema Arrecifal Lobos-Tuxpan. *CICIMAR Océanides* 31 (1): 23–34.
- Del Mónaco C, Narciso S, Alfonso F, Giménez E, Bustillos F (2010) Evaluación de las comunidades de corales y peces de algunos arrecifes de la isla La Tortuga y cayos adyacentes, Venezuela. *Boletín del Centro de Investigaciones Biológicas* 44 (3): 353–376.
- Delphy J (1938) [Les actinies athénaires (*Actiniaria Athenaria*) de la faune Française. *Bulletin du Muséum national d'Histoire naturelle*, Paris 10 (6): 619–622.
- Delphy J (1939) Présence sur la côte française de la Manche de l'Actinie *Bolocera Tuediae* (Johnston)? et quelques remarques sur d'autres Anthozoaires. *Bulletin du Muséum national d'Histoire naturelle* 11 (2): 267–270.
- Dickinson EC, Overstreet LK, Dowsett RJ, Bruce MD (2011) Priority! The Dating of Scientific Names in Ornithology. A Directory to the Literature and its Reviewers. Aves Press Limited, Northampton, 319 pp, CD-ROM.
- Donati V (1765) Storia naturale dell' Antipate, o corallo nero dell'Adriatico. *Giornale d'Italia, spettante all scienza naturale, e principalmente all' agricoltura, alle aru, ed al commercio*, 1: 51–56, 60–64, 2 pl. [pp. 51–56, 1 pl. (issue no. 6, session of 18 August 1764); pp. 60–64, 1 pl. (issue no. 8, session of 25 August 1764)]
- Düben MW, Koren J (1847) Om nogle norske Actinier. *Forhandlinger ved de Skandinaviske Naturforskere*, [den 11–18 July 1844], pp. 266–268.
- Duchassaing de Fombressin P (1850) Animaux radiaires des Antilles. Plon Frères, Paris, 33 + [1] pp., pls. 1, 2. [10 Aug 1850 (Anonymous 1850: 401)]
- Duchassaing de Fombressin P, Michelotti J (1860) Mémoire sur les Coralliaires des Antilles. Extrait des Mémoires de l'Académie des Sciences de Turin. Série II. Tome XIX. L'Imprimerie Royale, Turin, 89 pp., pls. 1–10. [Separate of Duchassaing de Fombressin and Michelotti (1861)]
- Duchassaing de Fombressin P, Michelotti J (1861) Mémoire sur les Coralliaires des Antilles. *Memoria della Reale Accademia delle Scienze di Torino* (2) 19: 279–365, pls. 1–10. [First issued as a separate as Duchassaing de Fombressin and Michelotti (1860)]
- Duchassaing de Fombressin P, Michelotti J (1864) Supplément au Mémoire sur les Coralliaires des Antilles. Extrait des Mémoires de l'Académie des Sciences de Turin. Série II. Tome

- XXIII. L'Imprimerie Royale, Turin, 112 pp., pls. 1–11. [Separate of Duchassaing de Fombressin and Michelotti (1866)]
- Duchassaing de Fombressin P, Michelotti J (1866) Supplément au Mémoire sur les Coralliaires des Antilles. Extrait des Mémoires de l'Académie des Sciences de Turin. Memoria della Reale Accademia delle Scienze di Torino (2) 23: 97–206, pls. 1–11, [First issued as a separate as Duchassaing de Fombressin and Michelotti (1864)]
- Duerden JE (1898) Jamaican Actiniaria. Part I.—Zoantheae. Scientific Transactions of the Royal Dublin Society (2) 6 (14): 329–384, pls. 17a, 18a, 19, 20. [Apr 1898 (contents and title-page)]
- Duerden JE (1903) West Indian sponge-incrusting actinians. Bulletin of the American Museum of Natural History 19 (18): 495–503, pls. 44–47.
- Duméril C (1860) Histoire générale, classification naturelle et méthodique des Insectes à l'aide de tableaux synoptiques. Bulletin Société entomologique de France (3) 8 (1): i–v.
- Duncan FM (1937) On the dates of publication of the Society's 'Proceedings,' 1859–1926. With an appendix containing the dates of publication of 'Proceedings,' 1830–1858, compiled by the late F. H. Waterhouse, and of the 'Transactions,' 1833–1869, by the late Henry Peavot, originally published in P. Z. S. 1893, 1913. Proceedings of the Zoological Society of London (A – General and Experimental) 1937 (1): 71–84. <https://doi.org/10.1111/j.1469-7998.1937.tb08500.x>
- Ehrenberg CG (1828–1831) Phytozoa. Symbolae physicae animalia evertebrata exclusis insectis. Series prima cum tabularum decade prima continent animalia Africana et Asiatica. In: Symbolae physicae, seu Icones adhuc ineditae corporum naturalium novorum aut minus cognitorum, quae ex itineribus per Libyam, Aegyptum, Nubiam, Dengalam, Syriam, Arabiam et Habessiniam. Pars Zoologica, 4. Officina Academica, Berolini, [65] pp., 6 pls.
- Ehrenberg CG (1834a) Beiträge zur physiologischen Kenntniss der Corallenthiere im allgemeinen, und besonders des rother Meeres, nebst einem Versuche zur physiologischen Systematik derselbe. Abhandlungen der Königlischen Akademie der Wissenschaften zu Berlin 1832 (1): 225–380, 1 table. [Also issued as separate as Ehrenberg (1834b)]
- Ehrenberg CG (1834b) Über die Natur und Bildung der Coralleninseln und Corallenbänke im rothen Meere. Eine in der Königlischen Akademie der Wissenschaften am 22. März 1832 gelesene, im Februar 1834 revidirte und gedruckte Abhandlung. Königlischen Akademie der Wissenschaften, Berlin, 156 pp., 1 table. [Separate issue of Ehrenberg (1834a)]
- Ehrenberg CG (1839) Über die Bildung der Kreidelfsen und des Kreidemergels durch unsichtbare Organismen. Abhandlungen der Königlischen Akademie der Wissenschaften zu Berlin 1838 (1): 59–147, pls. 1–4.
- Ellis J (1768) An account of the *Actinia sociata*, or clustered animal-flower, lately found on the sea-coasts of the new-ceded islands. Philosophical Transactions of the Royal Society of London 57 (2): 428–437, pl. 19. [> 12 Nov 1767 (p. 428); 1 Jan 1768 (Fautin 2016: 430)]
- Ellis J, Solander D (1786) The Natural History of Many Curious and Uncommon Zoophytes, Collected from Various Parts of the Globe. Benjamin White and Son, London, xii + 206 pp., pls. 1–63. [before 23 Feb 1786 (Cornelius and Wells, 1988: 22); also noticed in Jul 1786 (Anonymous, 1786: 1–6); authorship of the new taxa in this publication have been debated with Haszprunar and Spies (2014: 284) recently advancing the view that the book

was authored by Watt and that new taxa should be cited as “Ellis and Solander, in Watt, 1786”; we herein follow the opinion of Cornelius and Wells (1988) and cite new taxa as “Ellis, in Ellis and Solander, 1786”]

- Esper EJC (1805) Die Pflanzenthier in Abbildungen nach der Natur mit Farben erleuchtet nebst Beschreibungen. Dritter Theil. Dreyzehende Lieferung. Raspe, Nürnberg, pp. 3–24, 2 pls. [1805 (Yang et al. 2014: 6)]
- Evenhuis NL (1997) Litteratura taxonomica dipterorum (1758–1930). Being a Selected List of the Books and Prints of Diptera Taxonomy From the Beginning of Linnaean Zoological Nomenclature to the End of the Year 1930; Containing Information on the Biographies, Bibliographies, Types, Collections, and Patronymic Genera of the Authors Listed in This Work; Including Detailed Information on Publication Dates, Original and Subsequent Editions, and Other Ancillary Data Concerning the Publications Listed Herein. Backhuys Publishers, Leiden, viii + 871 pp.
- Evenhuis NL (2003a) Dating and publication of the *Encyclopédie Méthodique* (1782–1832), with special reference to the parts of the *Histoire Naturelle* and details on the *Histoire Naturelle des Insectes*. *Zootaxa* 166: 1–48. <https://doi.org/10.11646/zootaxa.166.1.1>
- Evenhuis NL (2003b) Publication and dating of the journals forming the *Annals and Magazine of Natural History* and the *Journal of Natural History*. *Zootaxa* 385: 1–68. <https://doi.org/10.11646/zootaxa.385.1.1>
- Fautin DG (2016) Catalog to families, genera, and species of orders Actiniaria and Corallimorpharia (Cnidaria: Anthozoa). *Zootaxa* 4145 (1): 1–449. [29 Jul 2016 (pp. 1, 2)] <https://doi.org/10.11646/zootaxa.4145.1.1>
- Fautin DG, Buddemeier RW (2008) Biogeoinformatics of the Hexacorals. <http://www.kgs.ku.edu/Hexacoral/>
- Fautin DG, Daly M (2009) Actinaria, Corallimorpharia, and Zoanthidea (Cnidaria) of the Gulf of Mexico. pp. 349–357. In: Felder DL, Camp DK (Eds). Gulf of Mexico. Origins, Waters and Biota. Volume 1, Biodiversity. Texas A & M University Press, College Station, Texas, xix + 1393 pp.
- Fautin DG, Zelenchuk T, Raveendran D (2007) Genera of orders Actiniaria and Corallimorpharia (Cnidaria, Anthozoa, Hexacorallia), and their type species. *Zootaxa*, 1668: 183–244.
- Fautin DG, Williams RB, Molodtsova T (2012) Case 3574. *Cereus* Ilmoni, 1830 (Cnidaria, Anthozoa): proposed designation of a new type species. *Bulletin of Zoological Nomenclature* 69 (1): 20–23. [31 Mar 2012 (issue front cover)]
- Fisher I (ed) (1893) Bibliographies of the Present Officers of Yale University. Together with the Bibliography of the Late President Porter. Tuttle, Morehouse & Taylor, New Haven, 160 pp.
- Forbes E (1840) On the British Actiniadae. *Annals of Natural History* 5 (30): 180–184, pl. 3. [1 May 1840 (Evenhuis 2003b: 14)]
- Friedländer R (1888) Hatschek, B., Lehrbuch der Zoologie. Eine morphologische Uebersicht des Thierreiches zur Einführung in das Studium dieser Wissenschaft. Lieferung 1. Jena 1888. gr. 8. pg. 4 u. 1–144 m. 155 Abbildungen. Naturae novitates, Bibliographie neuer Erscheinungen aller Länder auf dem Gebiete der Naturgeschichte und der exacten Wissenschaften 10 (24), 378. [1 Dec 1888 (Schmid 1984: 651)]
- Friedländer R (1889) Hatschek, B., Lehrbuch der Zoologie. Eine morphologische uebersicht des Thierreiches zur Einführung in das Studium dieser Wissenschaft. Lieferung 2. Jena

1889. gr. 8. pg. 145–304 m. 444 Abbildungen. Liefg. 1. 1888. m. 455 Abbildungen. *Naturae novitates, Bibliographie neuer Erscheinungen aller Länder auf dem Gebiete der Naturgeschichte und der exacten Wissenschaften* 11 (17–18), 279. [15 Sep 1889 (Schmid 1984: 651)]
- Friedländer R (1891) Hatschek, B., *Lehrbuch der Zoologie. Eine morphologische Uebersicht der Thierreiches zur Einführung in das Studium dieser Wissenschaft. Lieferung 3.* Jena 1891. gr. 8. pg. 305–432 m. 111 Abbildungen. Liefg. 1 u. 2. 1888–1889. m. 296 Abbildungen. *Naturae novitates, Bibliographie neuer Erscheinungen aller Länder auf dem Gebiete der Naturgeschichte und der exacten*, 13 (16): 305. [1 Aug 1891 (Schmid, 1984: 651)]
- Friedländer R (1901) Delage, Y. et Hérouard, E., *Les Coelentérés.* *Naturae novitates, Bibliographie neuer Erscheinungen aller Länder auf dem Gebiete der Naturgeschichte und der exacten Wissenschaften* 23 (22), 579. [15 Nov 1901 (Schmid 1984: 651)]
- Fujii T, Reimer JD (2011) Phylogeny of the highly divergent zoanthid family Microzoanthidae (Anthozoa, Hexacorallia) from the Pacific. *Zoologica Scripta* 40 (4): 418–431. <https://doi.org/10.1111/j.1463-6409.2011.00479.x>
- Fujii T, Reimer JD (2013) A new family of diminutive zooxanthellate zoanthids (Hexacorallia: Zoantharia). *Zoological Journal of the Linnean Society* 169 (5): 509–522. <https://doi.org/10.1111/zoj.12075>
- Fujii T, Reimer JD (2016) A new solitary free-living species of the genus *Sphenopus* (Cnidaria: Anthozoa: Zoantharia: Sphenopidae) from Okinawa-jima Island, Japan. *ZooKeys* 606: 11–24. <https://doi.org/10.3897/zookeys.606.9310>
- Garcieres T (1676) *The Admirable Virtues, and Wonderful Effects of the True and Genuine Tincture of Coral, in Physick; Grounded Upon Reason, Established by Experience, and Confirmed by Authentical Authors in All Ages. Never Found Plainly Till Now.* Samuel Sprint, London, [vii] + 83 pp.
- Gazave E, Carteron S, Chenuil A, Richelle-Maurer E, Boury-Esnault N, Borchellini C (2010) Polyphyly of the genus *Axinella* and of the family Axinellidae (Porifera: Demospongiae). *Molecular Phylogenetics and Evolution* 57 (1): 35–47. <https://doi.org/10.1016/j.ympev.2010.05.028>
- Gegenbaur C (1856) *Versuch eines Systemes der Medusen, mit Beschreibung neuer oder wenig gekannter Formen; zugleich ein Beitrag zur Kenntnis der Fauna des Mittelmeeres.* *Zeitschrift für wissenschaftliche Zoologie* 8 (2): 202–273, pls. 7–10. [12 Jul 1856 (contents, p. iii)]
- Gentry A (1992) Comments on the date of publication of John McCrady's hydrozoan paper *Gymnophthalmata of Charleston Harbor*. *Bulletin of Zoological Nomenclature*, 49 (4): 288–289. [17 Dec 1992 (issue front cover)]
- Gistel J (1848) *Naturgeschichte des Thierreichs für höhere Schulen.* Hoffmann, Stuttgart, xvi + 216 + [4] pp., 32 pls. [15 Aug 1848 (Naumann 1848: 120)]
- Gmelin JF (1791) *Caroli a Linné Systema Naturae per regna tria naturae. Edition decima tertia. Tom. I. Pars VI.* Lipsiae, 3021–3910 pp. [14 May 1791 (Hopkinson, 1908: 1036)]
- Gosse PH (1853) Notes on some new or little-known marine animals (No. 2). *Annals and Magazine of Natural History* (2) 12 (69): 153–159, pl. 10. [1 Sep 1853 (Evenhuis 2003b: 19)]

- Gosse PH (1855) Description of *Peachia hastata*, a new genus and species of the class Zoophyta; with observations on the family Actiniadae. Transactions of the Linnean Society of London 21 (4): 267–276, pls. 28, 29. [10 Nov 1855 (Raphael 1970: 75)]
- Gosse PH (1858–1859) Actinologia Britannica. A History of the British Sea-Anemones and Corals. Van Voorst, London, xl + [1] + 362 pp., pls. 1–12. [1859 for part 12 containing zoanthid names (Williams RB, *in litt.*, 21 Jun 2012); reissued as a completed work as Gosse (1859)]
- Gosse PH (1859) Actinologia Britannica. A History of the British Sea-Anemones and Corals. Van Voorst, London, xl + [1] + 362 pp., 12 pls. [1859 (Williams RB, *in litt.*, 21 Jun 2012)]
- Gravier C (1918) Note sur une Actinie (*Thoracactis* n. g., *Topsenti* n. sp.) et un Annélide Polychète (*Hermadion Fauveli* n. sp.), commensaux d'une Éponge siliceuse (*Sarostegia oculata* Topsent). Bulletin de l'Institut océanographique 15 (344): 1–20. [30 Sep 1918 (p. 1)]
- Gray JE (1828–1830) Spicilegia zoologica; or Original Figures and Short Systematic Descriptions of New and Unfigured Animals. Treüttel, Würtz and Co., London, 12 pp., 11 pls. [Dates of publication from Petit (2012: 19): part 1, pp. 1–8, pls. 1–6 (1 July 1828); part 2, pp. 9–12, pls. 7–11 (August 1830)]
- Gray JE (1832) [T]he Animal Flowers, or Zoantharia, and the stellated corals. pp. 94–106. In: Synopsis of the Contents of the British Museum. Twenty-Sixth Edition. G. Woodfall and Son, London, [iv] + 236 pp.
- Gray JE (1840) Radiated Animals. pp. 55–73. In: Synopsis of the Contents of the British Museum. Forty-First Edition. G. Woodfall and Son, London, [iv] + 370 pp. [27 Jun 1840 (Petit, 2012: 30)]
- Gray JE (1858) Note on *Dysidea papillosa*, Johnston. Annals and Magazine of Natural History (3) 2 (12): 489. [1 Dec 1858 (Evenhuis, 2003b: 21)]
- Gray JE (1859) On the *Dysidea papillosa* of Dr. Johnston. Proceedings of the Zoological Society of London 1858 (380): 531–532. [Jan–May 1859 (Duncan, 1937: 83)]
- Gray JE (1867) Notes on Zoanthinae, with descriptions of some new genera. Proceedings of the Zoological Society of London 1867 (1): 233–240. [May 1867 (Duncan, 1937: 72)]
- Haddon AC (1895) Branched worm-tubes and *Acrozoanthus*. Scientific Proceedings of the Royal Dublin Society 8 (4): 344–346. [27 Jul 1895 (p. 344)]
- Haddon AC (1898) The Actiniaria of Torres Straits. Scientific Transactions of the Royal Dublin Society (2) 6 (16): 393–520, pls. 22–32. [Aug 1898 (contents and title-page)]
- Haddon AC, Shackleton AM (1891a) A revision of the British Actiniae. Part II. The Zoantheae. In: Reports on the zoological collections made in the Torres Straits by A.C. Haddon, 1888–1889. Scientific Transactions of the Royal Dublin Society (2) 4 (12): 609–672, pls. 58–60. [Nov 1891 (wrapper)]
- Haddon AC, Shackleton AM (1891b) Actiniae: I. Zoantheae. In: Reports on the zoological collections made in the Torres Straits by Professor A.C. Haddon, 1888–1889. Scientific Transactions of the Royal Dublin Society (2) 4 (13): 673–701, pls. 61–64. [Dec 1891 (wrapper)]
- Haeckel E (1875) Arabische Korallen. Ein Ausflug nach den Korallenbänken des rothen Meeres und ein Blick in das leben der Korallenthier. Populäre Vorlesung mit wissenschaftlichen Erläuterungen. Georg Reimer, Berlin, 48 pp., pls. 1–6, frontispiece. [imprint 1876 but published 1875 (Leuckart, 1875: 463)]

- Haeckel E (1896) Systematische Phylogenie der Wirbellosen Thiere (Invertebrata). Zweiter Theil. Des Entwurfs einer systematischen Stammesgeschichte. Georg Reimer, Berlin, xviii + 720 pp.
- Haime J (1849) Note sur le polypiéroïde d'un *Leiopathes glaberrima*. Annales des Sciences Naturelles, comprenant la zoologie, la botanique, l'anatomie et la physiologie comparées des deux règnes, et l'histoire des corps organisés fossiles (3) 12: 224–226. [Oct ? 1849 (signature, p. 225)]
- Hallier E (1903) Carl Chun, Aus den Tiefen Weltmeeres. Schilderungen von der deutschen Tiefsee-Expedition. 2. umgearbeitete u. stark vermehrte Aufl. Jena, Gustav Fischer. (VI, 592 S.) 18 M., geb. 20 M. p. 63. In: Blätter für Volksbibliotheken und Lesehallen, Beiblatt zum Centralblatt für Bibliothekswesen 4 (3–4) [15 Apr 1903]: 41–72.
- Hartog JC den (1980) Caribbean shallow water Corallimorpharia. Zoologische Verhandelingen, Leiden 176: 2–83, pls. 1–14. [14 Mar 1980 (issue title-page)]
- Hatschek B (1888–1891) Lehrbuch der Zoologie, eine morphologische Übersicht des Thierreiches zur Einführung in das Studium dieser Wissenschaft. Gustav Fischer, Jena, iv + 432 pp. [< 1 Dec 1888 (*Lieferung* 1, pp. i–iv, 1–144) (Friedländer, 1888: 378); < 15 Sep 1889 (*Lieferung* 2, pp. 145–304) (Friedländer, 1889: 279); < 1 Aug 1891 (*Lieferung* 3, pp. 305–432) (Friedländer, 1891: 305)]
- Heider AR v (1899a) Über zwei Zoanthen. Zeitschrift für Wissenschaftliche Zoologie 66 (2): 269–288, pls. 16–17. [21 Jul 1899 (contents, p. [iii]); issued as a separate as Heider (1899b)]
- Heider AR v (1899b) Über zwei Zoanthen. Arbeiten aus dem Zoologischen Institut zu Graz, 6(3): 119–138, pls. 16–17. [< 16 Sep 1899 (date-stamp on p. 119 of Biodiversity Heritage Library copy); separate issue of Heider (1899a)]
- Herberts C (1972) Etude systématique de quelques zoanthaires tempérés et tropicaux. Tethys, (Supplément) 3: 69–156.
- Hertwig R (1882) Report on the Actinaria dredged by H.M.S. *Challenger* during the years 1873–1876. In: Report of the Scientific Results of the Voyage of H.M.S. *Challenger* During the Years 1873–76, Under the Command of Captain George S. Nares, R.N., F.R.S. and the Late Captain Frank Tourle Thomson, R.N. Prepared Under the Superintendence of the Late Sir C. Wyville Thomson, Knt., F.R.S., &c. Regius Professor of Natural History in the University of Edinburgh Director of the Civilian Scientific Staff on Board and Now of John Murray, LL.D., Ph.D., &c. One of the the Naturalists of the Expedition. Zoology—Vol. VI. Her Majesty's Government, Edinburgh, part 15: 1–136, pls. 1–14. [31 Dec 1882 (Low and Evenhuis 2013: 409)]
- Hertwig R (1888) Report on the Actinaria dredged by H.M.S. *Challenger* during the years 1873–1876. Supplement. In: Report of the Scientific Results of the Voyage of H.M.S. *Challenger* During the Years 1873–76, Under the Command of Captain George S. Nares, R.N., F.R.S. and the Late Captain Frank Tourle Thomson, R.N. Prepared Under the Superintendence of the Late Sir C. Wyville Thomson, Knt., F.R.S., &c. Regius Professor of Natural History in the University of Edinburgh Director of the Civilian Scientific Staff on Board and Now of John Murray, LL.D., Ph.D., &c. One of the the Naturalists of the Expedition. Zoology—Vol. XXVI. Her Majesty's Government, Edinburgh, part 73: 1–56, pls. 1–4. [Nov 1888 (Low and Evenhuis 2013: 418)]

- Hibino Y, Todd PA, Yang S-Y, Benayahu Y, Reimer JD (2014) Molecular and morphological evidence for conspecificity of two common Indo-Pacific species of *Palythoa* (Cnidaria: Anthozoa). *Hydrobiologia* 733 (1): 31–43. <https://doi.org/10.1007/s10750-013-1587-5>
- Hoeksema B, Reimer JD (2013) Zoantharia. In: Fautin, Daphne G (2013) Hexacorallians of the World. Accessed through: World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&cid=607338> on 2016-08-12
- Holdsworth EWH (1861a) On an undescribed species of British *Zoanthus*. *Proceedings of the Zoological Society of London* 1861 (1): 99–100. [May 1861 (Duncan, 1937: 72)]
- Holdsworth, EWH (1861b) On an undescribed species of British *Zoanthus*. *Annals and Magazine of Natural History* (3) 7 (42): 484–485. [1 Jun 1861 (Evenhuis, 2003b: 22)]
- Holthuis LB (1955) A supplementary note on the carcinological work of C.S. Rafinesque. *Zoologische Mededelingen, Leiden* 32 (26): 279–281.
- Holthuis LB (1969) Thomas Say as a carcinologist. pp. v–xv. In: Say T. *An Account of the Crustacea of the United States*. *Historiae naturalis classica*, 73: i–xv, 1–134.
- Hopkinson J (1908) Dates of publication of the separate parts of Gmelin's edition (13th) of the 'Systema Naturae' of Linnaeus. *Proceedings of the Zoological Society of London* 1907 (4): 1035–1037.
- Hughes G (1750) *The Natural History of Barbados*. In *Ten Books*. The Author, London, [vii] + [1] + 314 + [7] + [7] pp., 29 pls., 1 map.
- Haszprunar G, Spies M (2014) An integrative approach to the taxonomy of the crown-of-thorns starfish species group (Asteroidea: *Acanthaster*): A review of names and comparison to recent molecular data. *Zootaxa* 3841 (2): 271–284. <https://doi.org/10.11646/zootaxa.3841.2.6>
- ICZN (International Commission on Zoological Nomenclature) (1969) Opinion 883. *Cavolinia* Abilgaard, 1791 (Gastropoda): grant under Plenary Powers of precedence over *Cavolinia* Bruguière, 1791. *Bulletin of Zoological Nomenclature* 26 (1): 28–31. [12 May 1969 (issue front cover)]
- ICZN (International Commission on Zoological Nomenclature) (1987) Opinion 1461. A ruling on the authorship and dates of the text volumes of the *Histoire naturelle* section of Savigny's *Description de l'Égypte*. *Bulletin of Zoological Nomenclature* 44 (3): 219–220. [25 Sep 1987 (issue front cover)]
- ICZN (International Commission on Zoological Nomenclature) (1992) Opinion 1689. *Epizoanthus* Gray, 1867 (Cnidaria, Anthozoa): conserved. *Bulletin of Zoological Nomenclature* 49 (3): 236–237. [30 Sep 1992 (issue front cover)]
- ICZN, International Commission on Zoological Nomenclature (1998) Opinion 1895. *Riisea* and *riisei* Duchassaing & Michelotti, 1860 (Cnidaria, Anthozoa): conserved as the correct original spellings of generic and specific names based on the surname Riise. *Bulletin of Zoological Nomenclature* 55 (2): 121–122. [30 Jun 1998 (issue front cover)]
- ICZN (International Commission on Zoological Nomenclature) (1999) *International Code of Zoological Nomenclature*. Fourth Edition. The International Trust for Zoological Nomenclature, London, xxix + 306 pp.

- ICZN (International Commission on Zoological Nomenclature) (2014) Opinion 2334 (Case 3574). *Cereus* Ilmoni, 1830 (Cnidaria, Anthozoa): a new type species designated. Bulletin of Zoological Nomenclature 71 (2): 136–137. [30 Jun 2014 (issue front cover)]
- Ilmoni I (1830) Dr. Ilmoni aus Finnland schickte folgende Beiträge zur Naturgeschichte der Actinien ein. Isis von Oken 23 (6): 694–699.
- Imperato F (1599) Dell'istoria natvrale di Ferrante Imperanto Napolitano. Libri XXVIII. Nella qvale ordinatamenta si tratta della diusersa condition di miniere, e pietre. Con alcune historie di Piante, & Animali; sin'hera non date in luce. Con privilegio. Stamparia à Porta Reale, Napoli, [xxii] + 791 pp., frontispiece.
- Irei Y, Sinniger F, Reimer JD (2015) Descriptions of two azooxanthellate *Palythoa* (Order Zoantharia: Subclass Hexacorallia) species from the Ryukyu Archipelago, southern Japan. ZooKeys 478: 1–26. <https://doi.org/10.3897/zookeys.478.8512>
- Johnston G (1842) History of the British Sponges and Lithophytes. W.H. Lizars, Edinburgh / S. Highley, London / W. Curry, Jun and Co., Dublin, xii + 264 pp., pls. 1–25. [6 Feb 1842–6 Mar 1842 (Anonymous 1842: 63)]
- Karlson RH (1980) Alternative competitive strategies in a periodically disturbed habitat. Bulletin of Marine Science 30 (4): 894–900.
- Kayal E, Roure B, Philippe H, Collins AG, Lavrov DV (2013) Cnidarian phylogenetic relationships as revealed by mitogenomics. BMC Evolutionary Biology 13: 5. <https://doi.org/10.1186/1471-2148-13-5>
- Kenny JS (2008) The Biological Diversity of Trinidad and Tobago. A Naturalist's Notes. Prospect Press, Port of Spain, xiv + 265 pp.
- Koch CL (1850) Uebersicht des Arachnidensystems. Fünftes Heft. J. L. Lotzbeck, Nürnberg, 8–104 pp.
- Koren J, Danielssen DC (1877) Description of some new Norwegian coelenterates. In: Koren J, Danielssen DC (eds). Fauna littoralis Norvegiae. Part Three. J.D. Beyer, Bergen, pp. 77–81, pl. 9. [> 25 Apr 1877 (unnumbered preface)]
- Koupaei AN, Mostafavi PG, Mehrabadi JF, Fatemi SMR (2014) Molecular diversity of coral reef-associated zoanths off Qeshm Island, northern Persian Gulf. International Aquatic Research 6: 64 [10 pp.]. <https://doi.org/10.1007/s40071-014-0064-8>
- Krishna JM, Gophane A (2013) Cnidarian from the coast of Goa – identified to the species level. Journal of Pharmacognosy and Phytochemistry 2 (1): 209–218.
- Lacaze-Duthiers H (1864a) Mémoire sur les Antipathaires, genre *Gerardia* (L.D.). In: Séance du lundi 11 juillet 1864. Présidence de M. Morin. Comptes Rendus hebdomadaires des Séances de l'Académie des Sciences 59 (2): 86–89. [> 11 Jul 1864 (date of *Séance*, p. 86); < 17 Aug 1864 (p. 333 of same volume states that issue no. 4 of the same volume had already been published; see also Dickinson et al. 2011: 193, 194)]
- Lacaze-Duthiers H (1864b) On the antipatharian genus *Gerardia*. Annals and Magazine of Natural History (3) 14 (82): 241–243. [1 Oct 1864 (Evenhuis 2003b: 23)]
- Lacaze-Duthiers H (1864c) Mémoire sur les anthipathaires (genre *Gerardia*, L.D.). Annales des Sciences Naturelles, Zoologie et paléontologie comprenant l'anatomie, la physiologie, la classification et l'histoire naturelle des animaux (5) 2: 169–239, pls. 13–18. [31 Dec 1864 (Article 21.3.2 of the Code, ICZN 1999: 22)]

- Lamarck JBP (1801) Zoanthe. *Zoantha*. p. 363. In: Système des animaux sans vertèbres, ou Tableau général des classes, des ordres et des genres de ces animaux; Présentant leurs caractères essentiels et leur distribution, d'après la considération de leurs rapports naturels et leur organisation, et suivant l'arrangement établis dans les galeries du Muséum d'Histoire Naturelle, parmi leur dépouilles conservées; Précédé du discours d'ouverture du cours de zoologie, donné dans le Muséum National d'Histoire Naturelle l'an 8 de la République. L'Auteur et Déterville, Paris, viii + 432 pp. [21 Jan 1801 (Evenhuis and Cowie in Cowie, 1995: 216); 25 Jan 1801 (Evenhuis 1997: 430)]
- Lamouroux JVF (1812) Extrait d'un mémoire sur la classification des Polypiers coralligènes non entièrement pierreux. *Nouveaux Bulletin des Sciences*, par la Société Philomathique de Paris 3: 181–188.
- Lamouroux JVF (1816) Histoire des polypiers coralligènes flexibles, vulgairement nommés zoophytes. F. Poisson, Caen, lxxxiv + [1] + 559 + [1] pp., pls. 1–19. [Oct 1816 (Sherborn 1922: lxxvii)]
- Lamouroux JVF (1821) Exposition méthodique des genres de l'ordre des polypiers, avec leur description et celle des principales espèces, figurés dans 84 planches; les 63 premières appartenant à l'histoire naturelle des zoophytes d'Ellis et Solander. Veuve Agasse, Paris, ix + 115 pp., pls. 1–84. [Apr 1821 (Sherborn 1922: lxxvii)]
- Le Sueur CA (1817) Observations on several species of the genus *Actinia*; illustrated by figures (concluded). *Journal of the Academy of Natural Sciences of Philadelphia* 1 (6): 169–189, pls. 7, 8. [after 9 Dec 1817 (p. 169) and before 31 Dec 1817 (Holthuis 1969: vi, vii; Phillips 1841: 58)]
- Lesson RP (1830) Description des animaux de la famille des Actiniées. Chapitre XVI. Pp. 67–83. In: Duperrey LI, Voyage autour du monde, exécuté par Ordre du Roi, sur la Corvette de Sa Majesté, la Coquille, pendant les années 1822, 1823, 1824 et 1825, sous le Ministère et Conformément aux Instructions de S.E.M. le Marquis de Clermont-Tonnerre, Ministre de la Marine; et publié sous les Auspices de son Excellence M^{gr} le C^{te} de Chabrol, Ministre de la Marine et des Colonies. Zoologie. Tome second.—2^e partie. Arthus Bertrand, Paris, [2] + xii + [8] + 9–56. [10 Nov 1838 (Sherborn and Woodward 1901a: 392, 1906: 336); < 31 Dec 1838 (Cretella 2010: 94)]
- Leuckart R (1875) Bericht über die wissenschaftlichen Leistungen in der Naturgeschichte der niederen Thiere während der Jahre 1872–1875. (Zweite Hälfte). *Archiv für Naturgeschichte* 41 (2): 313–517.
- Long BG, Poiner IR, Wassenberg TJ (1995) Distribution, biomass and community structure of megabenthos of the Gulf of Carpentaria, Australia. *Marine Ecology Progress Series* 129: 127–139. <https://doi.org/10.3354/meps129127>
- Longo GO, Krajewski JP, Segal B, Floeter SR (2012) First record of predation on reproductive *Palythoa caribaeorum* (Anthozoa: Sphenopidae): insights on the trade-off between chemical defences and nutritional value. *Marine Biodiversity Records* 5: e29 [3 pp.]. <https://doi.org/10.1017/S1755267212000206>
- Low MEY, Evenhuis NL (2013) Dates of publication of the Zoology parts of the Report of the Scientific Results of the Voyage of H.M.S. Challenger During the Years 1873–76. *Zootaxa* 3701 (4): 401–420. <https://doi.org/10.11646/zootaxa.3701.4.1>

- Low MEY, Reimer JD (2011a) *Parazoanthus* Haddon & Shackleton, 1891, and Parazoanthidae Delage & Hérouard, 1901: conservation of usage by reversal of precedence with *Bergia* Duchassing & Michelotti, 1860, and Bergiidae Verrill, 1869 (Cnidaria: Anthozoa: Hexacorallia). *Zootaxa* 2995: 64–68. [15 Aug 2011 (p. 64)]
- Low MEY, Reimer JD (2011b) Saving an octocoral genus from a “zoanthid” genus: reversal of precedence of *Palythoe* Lamouroux, 1812, and *Muricea* Lamouroux, 1812 (Cnidaria: Anthozoa: Octocorallia). *Zootaxa* 3120: 63–66. [6 Dec 2011 (p. 63)]
- Low MEY, Reimer JD (2012a) Zoanthid housekeeping: nomenclatural notes on the Zoantharia (Cnidaria: Anthozoa: Hexacorallia). *Zootaxa* 3485: 83–88. [17 Sep 2012 (p. 83)]
- Low MEY, Reimer JD (2012b) *Isaura* Lamouroux, in Audouin, Bourdon, de Candolle, d’Aubebard de Férussac, Deshayes, Deslongchamps, É. Geoffroy Saint-Hilaire, I. Geoffroy Saint-Hilaire, Guérin, Guillemin, de Jussieu, Kunth, Delafosse, Lamouroux, Latreille, Prévost, Richard & Bory de Saint-Vincent, 1826, an available name and objective synonym of *Isaurus* Gray, 1828: reversal of precedence (Cnidaria: Anthozoa: Hexacorallia). *Zootaxa* 3532: 45–50. [31 Oct 2012 (p. 45)]
- Lwowsky F (1913) Revision der Gattung *Sidisia* Gray (*Epizoanthus* auct.). Ein Beitrag zur Kenntnis der Zoanthiden. *Zoologische Jahrbücher (Abteilung für Systematik, Ökologie und Geographie der Tiere)* 34 (5–6): 557–614. [30 May 1913 (contents, p. iv); also issued as a separate thesis with separate title-page]
- Manuel RL (1981) On the identity of the sea anemone *Scolanthus callimorphus* Gosse, 1853 (Actiniaria: Edwardsiidae). *Journal of Natural History* 15 (2): 265–276. [6 Feb 1981 (Evenhuis, 2003b: 59)] <https://doi.org/10.1080/00222938100770201>
- McCrary J (1857–1859) Gymnophthalmata of Charleston Harbor. *Proceedings of the Elliott Society of Natural History of Charleston* 1: 103–221, pls. 8–12. [15 Sep 1857 (pp. 103–104), 11 May 1859 (pp. 105–221) (Calder et al. 1992: 288; Gentry 1992: 288, 289); also issued as a separate with pagination pp. i–iii, 1–119 (Gentry 1992: 289)]
- McMurrich JP (1889) A contribution to the actinology of the Bermudas. *Proceedings of the Academy of Natural Sciences of Philadelphia* 41 (1): 102–126, pls. 6–7. [18 Jun 1889 (*verso* of title-page; also Dickinson et al. 2011: table 58)]
- McMurrich JP (1904) The Actiniae of the Plate Collection. *Fauna Chilensis. Abhandlungen zur Kenntniss der Zoologie Chiles nach den Sammlungen von Dr. L. Plate. Dritter Band. Zoologische Jahrbücher, supp.* 6 (2): 215–306, pls. 14–19. [9 May 1904 (contents, p. [iii])]
- Meigen JG (1800) Nouvelle classification des mouches à deux ailes (Diptera L.) d’après un plan tout nouveau. J. J. Fuchs, Paris, 40 pp. <https://doi.org/10.5962/bhl.title.119764>
- Milliman JD, Müller G, Förstner U (1974) Eulittoral (shallow-water) sedimentation. In: Milliman JD (ed). *Recent Sedimentary Carbonates. Part 1. Marine Carbonates*. Springer, Berlin, pp. 153–205. https://doi.org/10.1007/978-3-642-65528-9_6
- Milne Edwards H (1857) Histoire naturelle des Coralliaires ou Polypes proprement dits. Tome premier. Roret, Paris, xxxiv + 326 pp. [5 Dec 1857 (Anonymous, 1857: 634)]
- Milne-Edwards H, Haime J (1851) Monographie des polypiers fossiles des terrains palaeozoïques, précédée d’un tableau général de la classification des polypes. *Archives du Muséum d’Histoire naturelle* 5: 1–502, pls. 1–20.

- Montenegro J, Low MEY, Reimer JD (2015a) The resurrection of the genus *Bergia* Duchassaing and Michelotti, 1860 (Anthozoa, Zoantharia, Parazoanthidae). Systematics and Biodiversity, 14 (1): 63–73. <https://doi.org/10.1080/14772000.2015.1101028>
- Montenegro J, Sinniger F, Reimer JD (2015b) Unexpected diversity and new species in the sponge-Parazoanthidae association in southern Japan. Molecular Phylogenetics and Evolution 89: 73–90. <https://doi.org/10.1016/j.ympev.2015.04.002>
- Muirhead A, Ryland JS (1985) A review of the genus *Isaurus* Gray, 1828 (Zoanthidea), including new records from Fiji. Journal of Natural History 19 (2): 323–335. [23 May 1985 (Evenhuis 2003b: 60)] <https://doi.org/10.1080/00222938500770241>
- Nagabhushanam AK, Jothinayagam JT (1982) Observations on the zoanthid, *Sphenopus marsupialis* (Gmelin) (Anthozoa: Zoanthidea: Sphenopidae), along the southeast coast of India, with notes on its behaviour in aquaria. Records of the Zoological Survey of India 80 (1–2): 17–26.
- Nardo GD (1844) Distribuzione naturale in ordini, famiglie e generi della Classe dei Zoofitarj (Blainville). In: Sezione di Zoologica, Anatomia comparata e Fisiologia. Atti della Quinta Unione degli Scienziati italiani, tenuta in Lucca, nel settembre del MDCCCXLIII. Giusti, Lucca, pp. 430–435.
- Nardo GD (1877) Sull' Antipate dell' Adriatico. Memoria postuma del Celebre Vitaliano Donati sfuggia all' Occhio dei Naturalisti. Atti del Reale Istituto Veneto di Scienze, Lettere ed Arti (5) 3: 673–678.
- Naumann R (1848) Gistel, Dr. Johs., Naturgeschichte des Thierreichs. Für die höheren Schulen bearbeitet. Mit einem Atlas von 32 (lith.) Taf. (darstellend 617 illum. Figuren) und mehrerer eingedruckten Xylographien. Fol. Stuttgart. 8 Thlr. In: Uebersicht der neuesten Literatur. Intelligenz-Blatt zum Serapeum, 9(15) [15 Aug 1848]: 120.
- Neave SA (1940) Nomenclator zoologicus. A List of the Names of Genera and Subgenera in Zoology from the Tenth Edition of Linnaeus 1758 to the End of 1935. In Four Volumes. Vol. III. M–P. The Zoological Society of London, London, ii + 1065 pp.
- Ng PKL, Low MEY (2010) On the generic nomenclature of nine brachyuran names, with four replacement names and two *nomina protecta* (Crustacea: Decapoda). Zootaxa 2489: 34–46.
- Ocaña O, Brito A (2004) A review of Gerardiidae (Anthozoa: Zoantharia) from the Macaronesian islands and the Mediterranean Sea with the description of a new species. Revista de la Academia Canaria de Ciencias 15 (3–4): 159–189. [Aug 2004 (p. 159)]
- Ocaña O, Brito A, Núñez J, Bacallado JJ (1995) Redescrpción de *Gerardia savaglia* (Bertoloni, 1819) (Anthozoa: Zoantharia: Gerardiidae). Vieraea 24: 153–164. [Dec 1995 (p. 153)]
- Ocaña O, Brito A, González G, Herrera R (2007) Additions in relation to Gerardiidae from the Macaronesian waters and the Mediterranean Sea (Anthozoa: Zoantharia). Vieraea 35: 163–168. [Oct 2007 (p. 163)]
- Ong CW, Reimer JD, Todd PA (2013) Morphologically plastic responses to shading in the zoanthids *Zoanthus sansibaricus* and *Palythoa tuberculosa*. Marine Biology 160:1053–1064. <https://doi.org/10.1007/s00227-012-2158-4>
- Pallas PS (1766) Elenchus zoophytorum sistens generum adumbrationes generaliores et specierum cognitarum succinctas descriptiones cum selectis auctorum synonymis. Francis-

- cum Varrentrapp, Hagrae [= The Hague], xvi + 17–28 + 451 pp. <https://doi.org/10.5962/bhl.title.6595>
- Palmer CV, Bythell JC, Willis BL (2012) Enzyme activity demonstrates multiple pathways of innate immunity in Indo-Pacific anthozoans. *Proceedings of the Royal Society B* 279 (1743): 3879–3887. <https://doi.org/10.1098/rspb.2011.2487>
- Pax F (1910a) Studien an westindischen Actinien. Habilitationsschrift, welche nebst den beigefügten Thesen mit Genehmigung der hohen philosophischen Fakultät der Königlichen Universität Breslau zur Erlangung der Venia legendi. Montag, den 17. Oktober 1910, um 12 Uhr, im Musiksaale der Universität öffentlich verteidigen wird. Gustav Fischer, Jena, 166 + [2] pp., 10 pls. [17 Oct 1910 (title-page); separate issue of Pax (1910b)]
- Pax F (1910b) Studien an westindischen Actinien. *Zoologische Jahrbücher*, supp. 11 (2): 157–330, pls. 11–19. [21 Nov 1910 (contents page); also issued as a separate as Pax (1910a)]
- Pax F, Müller I (1956) La collection de Zoanthaires du Musée océanographique de Monaco. *Bulletin de l'Institut océanographique* 53 (1076): 1–27. [22 Mar 1956 (wrapper and p. 1)]
- Pax F, Müller I (1957) Zoantharien aus Viet-Nam. *Mémoires du Museum national d'Histoire naturelle (A)* 16 (1): 1–40, pls. 1–13. [Jul 1957 (p. 40)]
- Petit RE (2012) John Edward Gray (1800–1875): his malacological publications and molluscan taxa. *Zootaxa* 3214: 1–125. [29 Feb 2012 (p. 1)]
- Phillips JS (1841) [Dates of publication of the Journal of the Academy of Natural Sciences of Philadelphia]. In: Stated meeting, July 20, 1841. Vice President Morton in the Chair. *Proceedings of the Academy of Natural Sciences of Philadelphia* 1 (3): 57–59.
- Poche F (1914) Das System der Coelenterata. *Archiv für Naturgeschichte* 80A (5): 47–128. [Jul 1914 (issue wrapper)]
- Qin X-Y, Yang K-L, Li J, Wang C-Y, Shao C-L (2015) Phylogenetic diversity and antibacterial activity of culturable fungi derived from the zoanthid *Palythoa haddoni* in the South China Sea. *Marine Biotechnology* 17 (1): 99–109. <https://doi.org/10.1007/s10126-014-9598-4>
- Quatrefages A de (1841) Zoologie : Côtes de la Manche. *L'Institut, journal universel des sciences et des sociétés savantes en France et à l'étranger (1ère. section - Sciences mathématiques, physiques et naturelles)* 9 (416): 427. [16 Dec 1841 (p. 425)]
- Quatrefages A de (1842) Memoire sur les Edwardsies (*Edwardsia*, Nob.) nouveau genre de la famille des Actinies. *Annales des Sciences Naturelles, comprenant la zoologie, la botanique, l'anatomie et la physiologie compares des deux règnes, et l'histoire des corps organisés fossiles* (2) 18: 65–109.
- Quoy J, Gaimard JP (1834) Zoophytes. In: Voyage de découvertes de l'Astrolabe exécuté par ordre du Roi, pendant les années 1826–1827–1828–1829, sous le commandement de M.J. Dumont d'Urville. Zoologie. Tome Quatrième. J. Tastu, Paris, 390 pp. (Text), Zoophytes pls. 1–26 (Atlas). [“1833” 1834 (Sherborn and Woodward, 1901b: 333)]
- Rafinesque CS (1815) Analyse de la nature ou tableau de l'univers et des corps organisés. [The Author], Palerme [= Palermo], 224 pp. <https://doi.org/10.5962/bhl.title.106607>
- Rafinesque CS (1818) [Review of the] Journal of the Academy of Natural Sciences of Philadelphia, Vol. i, Part I, Philadelphia, 8 vo., pp. 220, and 9 plates. May to December, 1817. *American Monthly Magazine and Critical Review* 3 (4): 269–274. [Aug 1818 (Holthuis, 1955: 279)]

- Raphael S (1970) The publication dates of the *Transactions of the Linnean Society of London*, Series I, 1791–1875. *Biological Journal of the Linnean Society*, 2 (1): 61–76. <https://doi.org/10.1111/j.1095-8312.1970.tb01688.x>
- Reimer JD, Fujii T (2010) Four new species and one new genus of zoanthids (Cnidaria, Hexacorallia) from the Galápagos Islands. *ZooKeys* 42: 1–36. <https://doi.org/10.3897/zookeys.42.378>
- Reimer JD, Sinniger F (2010) Unexpected diversity in Canadian Pacific zoanthids (Cnidaria: Anthozoa: Hexacorallia): a molecular examination and description of a new species from the waters of British Columbia. *Marine Biodiversity*, 40 (4): 249–260. <https://doi.org/10.1007/s12526-010-0045-0>
- Reimer JD, Ono S, Fujiwara Y, Takishita K, Tsukahara J (2004) Reconsidering *Zoanthus* spp. diversity: molecular evidence of conspecificity within four previously presumed species. *Zoological Science* 21 (5): 517–525. <https://doi.org/10.2108/zsj.21.517>
- Reimer JD, Ono S, Iwama A, Takishita K, Tsukahara J, Maruyama T (2006) Morphological and molecular revision of *Zoanthus* (Anthozoa: Hexacorallia) from southwestern Japan, with descriptions of two new species. *Zoological Science* 23 (3): 261–275. <https://doi.org/10.2108/zsj.23.261>
- Reimer JD, Hirano S, Fujiwara Y, Sinniger F, Maruyama T (2007a) Morphological and molecular characterization of *Abyssoanthus nankaiensis*, a new family, new genus and new species of deepsea zoanthid (Anthozoa: Hexacorallia: Zoantharia) from a northwest Pacific methane cold seep. *Invertebrate Systematics* 21 (3): 255–262. <https://doi.org/10.1071/IS06008>
- Reimer JD, Takishita K, Ono S, Maruyama T (2007b) Diversity and evolution in the zoanthid genus *Palythoa* (Cnidaria: Hexacorallia) utilizing nuclear ITS-rDNA. *Coral Reefs* 26 (2): 399–410. <https://doi.org/10.1007/s00338-007-0210-5>
- Reimer JD, Nonaka M, Sinniger F, Iwase F (2008a) Morphological and molecular characterization of a new genus and new species of parazoanthid (Anthozoa: Hexacorallia: Zoantharia) associated with Japanese red coral (*Paracorallium japonicum*) in southern Japan. *Coral Reefs* 27 (4): 935–949. <https://doi.org/10.1007/s00338-008-0389-0>
- Reimer JD, Ono S, Tsukahara J, Iwase F (2008b) Molecular characterization of the zoanthid genus *Isaurus* (Anthozoa: Hexacorallia) and its zooxanthellae (*Symbiodinium* spp). *Marine Biology* 153 (3): 351–363. <https://doi.org/10.1007/s00227-007-0811-0>
- Reimer JD, Hirose M, Wirtz P (2010a) Zoanthids of the Cape Verde Islands and their symbionts: previously unexamined diversity in the Northeastern Atlantic. *Contributions to Zoology* 79 (4): 147–163.
- Reimer JD, Hirose M, Nishisaka T, Sinniger F, Itani G (2010b) *Epizoanthus* spp. associations revealed using DNA markers: a case study from Kochi, Japan. *Zoological Science* 27: 729–734. <https://doi.org/10.2108/zsj.27.729>
- Reimer JD, Nakachi S, Hirose M, Hirose E, Hashiguchi S (2010c) Using hydrofluoric acid for morphological investigations of zoanthids (Cnidaria: Anthozoa): a critical assessment of methodology and necessity. *Marine Biotechnology* 12 (5): 605–617. <https://doi.org/10.1007/s10126-009-9249-3>
- Reimer JD, Hirose M, Irei Y, Obuchi M, Sinniger F (2011a) The sands of time: rediscovery of the genus *Neozoanthus* (Cnidaria: Hexacorallia) and evolutionary aspects of sand

- incrustation in brachynemic zoanths. *Marine Biology* 158 (5): 983–993. <https://doi.org/10.1007/s00227-011-1624-8>
- Reimer JD, Ishikawa SA, Hirose M (2011b) New records and molecular characterization of *Acrozoanthus* (Cnidaria: Anthozoa: Hexacorallia) and its endosymbionts (*Symbiodinium* spp.) from Taiwan. *Marine Biodiversity* 41 (2): 313–323. <https://doi.org/10.1007/s12526-010-0069-5>
- Reimer JD, Foord C, Irei Y (2012a) Species diversity of shallow water zoanths (Cnidaria: Anthozoa: Hexacorallia) in Florida. *Journal of Marine Biology* 2012: article 856079 [14 pp.]. <https://doi.org/10.1155/2012/856079>
- Reimer JD, Lin M, Fujii T, Lane DJW, Hoeksema BW (2012b) The phylogenetic position of the solitary zoanthid genus *Sphenopus* (Cnidaria: Hexacorallia). *Contributions to Zoology* 81 (1): 43–54.
- Reimer JD, Lorion J, Irei Y, Hoeksema BW, Wirtz P (2014a) Ascension Island shallow-water Zoantharia (Hexacorallia: Cnidaria) and their zooxanthellae (*Symbiodinium*). *Journal of the Marine Biological Association of the United Kingdom*. <https://doi.org/10.1017/S0025315414000654>
- Reimer JD, Polisenio A, Hoeksema BW (2014b) Shallow-water zoantharians (Cnidaria, Hexacorallia) from the Central Indo-Pacific. *ZooKeys* 444: 1–57. <https://doi.org/10.3897/zookeys.444.7537>
- Reimer JD, Uyeno D, Berumen ML (2014c) First records of Parazoanthidae and Microzoanthidae (Anthozoa: Hexacorallia: Zoantharia) from the Red Sea. *Marine Biodiversity Records* 7: e7: 1–3. <https://doi.org/10.1017/s1755267214000025>
- Reimer JD, Kawamura I, Berumen ML (2015a) First record of Nanozoanthidae from the Red Sea. *Marine Biodiversity Records* 8: e19: 1–3. <https://doi.org/10.1017/s1755267214001481>
- Reimer JD, Wee HB, Put Jr A, Hoeksema BW (2015b) Zoantharia (Cnidaria: Anthozoa: Hexacorallia) of the South China Sea and Gulf of Thailand: a species list based on past reports and new photographic records. *Raffles Bulletin of Zoology* 63: 334–356.
- Risi MM, Macdonald AHH (2016) Molecular examination of rocky shore brachynemic zoantharians (Anthozoa: Hexacorallia) and their *Symbiodinium* symbionts (Dinophyceae) in the southwest Indian Ocean. *Marine Biodiversity* 46 (1): 113–127. <https://doi.org/10.1007/s12526-015-0331-y>
- Robineau-Desvoidy JB (1830) Essai sur les myodaires. Mémoires présentés par divers savans à l'Académie Royale des Sciences de l'Institut de France, Sciences Mathématiques et Physiques, (2) 2: 1–813.
- Roche J, Tixier-Durivault A (1951) Rapports des Gerardiides avec les Zoanthides et les Antipathaires. *Bulletin du Muséum d'Histoire naturelle, Paris*, (2) 23 (4): 402–409.
- Rodríguez-Viera L, Rodríguez-Casariño J, Pérez-García JA, Olivera Y, Perera-Pérez O (2012) Invertebrados marinos de la zona central del golfo de Ana María, Cuba. *Revista de Investigaciones Marinas* 32 (2): 30–38.
- Ryland JS (1997) Reproduction in Zoanthidea (Anthozoa: Hexacorallia). *Invertebrate Reproduction and Development* 31 (1–3): 177–188. <https://doi.org/10.1080/07924259.1997.9672575>
- Ryland JS, Muirhead A (1991) *Epizoanthus* Gray, 1867 (Cnidaria, Anthozoa): proposed conservation. *Bulletin of Zoological Nomenclature* 48 (1): 19–21. [26 Mar 1991 (issue front cover)]

- Ryland JS, Lancaster JE (2003) Revision of methods for separating species of *Protopalythoa* (Hexacorallia: Zoanthidea) in the tropical West Pacific. *Invertebrate Systematics* 17 (3): 407–428. <https://doi.org/10.1071/IS02008>
- Ryland JS, Westphalen D (2004) The reproductive biology of *Parazoanthus parasiticus* (Hexacorallia: Zoanthidea) in Bermuda. In: Fautin DG, Westfall JA, Cartwright P, Daly M, Wyttenbach CR (eds). *Coelenterate Biology 2003: Trends in Research on Cnidaria and Ctenophora*. *Hydrobiologia* 530–531: 411–419.
- Ryland JS, Lancaster JE (2004) A review of zoanthid nematocyst types and their population structure. In: Fautin DG, Westfall JA, Cartwright P, Daly M, Wyttenbach CR (eds). *Coelenterate Biology 2003: Trends in Research on Cnidaria and Ctenophora*. *Hydrobiologia* 530–531: 179–187.
- Ryland JS, Putron S de, Scheltema RS, Chimonides PJ, Zhadan DG (2000) Semper's (zoanthid) larvae: pelagic life, parentage and other problems. In: Jones MB, Azevedo JMN, Neto AI, Costa AC, Frias Martins AM (Eds). *Island, Ocean and Deep-Sea Biology*. *Hydrobiologia* 440: 191–198. <https://doi.org/10.1023/A:1004171127777>
- Ryland JS, Brasseur MM, Lancaster JE (2004) Use of cnidae in taxonomy: implications from a study of *Acrozoanthus australiae* (Hexacorallia, Zoanthidea). *Journal of Natural History* 38 (10): 1193–1223. <https://doi.org/10.1080/0022293031000155179>
- Savigny J-C (1811) Description de l'Égypte ou Recueil des observations et des recherches qui ont été faites en Égypte pendant l'expédition de l'armée française, publié par les ordres de Sa Majesté l'empereur Napoléon le Grand. *Histoire naturelle*. Tome deuxième [Atlas]. Imprimerie Impériale, Paris, pls. Polypes 1–14. [Reissued as Savigny (1826)]
- Savigny J-C (1826) Description de l'Égypte ou Recueil des observations et des recherches qui ont été faites en Égypte pendant l'expédition de l'armée française. *Seconde édition, dédiée au Roi*. *Histoire naturelle*. Tome deuxième [Atlas]. C.L.F. Panckoucke, Paris, pls. Polypes 1–14. [Reissue of Savigny (1811)]
- Saville-Kent W (1893) Coral and Coral Animals. pp. 139–203, pls. 21–30. In: *The Great Barrier Reef of Australia; Its Products and Potentialities*. Containing—An Account, With Copious Coloured and Photographic Illustrations (The Latter Here Produced for the First Time), of the Corals and Coral Reefs, Pearl and Pearl-shell, Bêche-de-mer, Other Fishing Industries, and the Marine Fauna of the Australian Great Barrier Region. W.H. Allen & Co., London, xvii + [1] + 387 pp., pls. 1–48, 1–16. [> 24 Feb 1893 (preface, p. xii); < 4 Mar 1893 (Anonymous 1893: 249)]
- Saville-Kent W (1897) *The Naturalist in Australia*. Illustrated by 50 Full-Page Collotypes, 9 Coloured Plates by Keulemans and Other Artists, And Over One Hundred Illustrations in the Text. Chapman & Hall, London, xv + [1] + 302 pp., 59 pls. [< 5 Jun 1897 (Anonymous, 1897: 744)]
- Sayre G (1959) Dates of Publication Describing Musci, 1801–1821. Troy, New York, [i] + 102 pp.
- Schmid R (1984) *Naturae novitates*, 1879–1944: its publication and intercontinental transit times mirror European history. *Taxon* 33 (4): 636–654. <https://doi.org/10.2307/1220781>
- Schmidt H (1972) Die Nesselkapseln der Anthozoen und ihre Bedeutung für die phylogenetische Systematik. *Helgoländer Wissenschaftliche Meeresuntersuchungen* 23 (4): 422–458. <https://doi.org/10.1007/BF01625294>

- Schmidt O (1862) Die Spongien des Adriatischen Meeres. Wilhelm Engelmann, Leipzig, viii + 88 pp., pls. 1–7. [Jul 1862 (Brockhaus 1862: 96)]
- Schuchert P (2010) The European athecate hydroids and their medusae (Hydrozoa, Cnidaria): Capitata Part 2. *Revue Suisse de Zoologie*, 117 (3): 337–555. [Sep 2010 (p. 337)]
- Schweigger AF (1819) Ueber eine neue Familie corallenähnlicher Thierpflanzen ohne leblose Substanz. pp. 94–100. In: *Beobachtungen auf naturhistorischen Reisen. Anatomisch-physiologische Untersuchungen über Corallen; nebst einem Anhang, Bemerkungen über den Bernstein enthaltend.* Georg Reimer, Berlin, viii + 127 + [5] pp., pls. 1–12.
- Sherborn CD (1897) On the dates of the natural history portion of Savigny's 'Description de l'Égypte'. *Proceedings of the Zoological Society of London* 1897 (1): 285–288.
- Sherborn CD (1922) Index animalium sive index nominum quae ab A.D. MDCCCLVIII generibus et speciebus animalium imposita sunt. Section secunda. A Kalendis Ianuariis, MDCCCLV usque ad finem Decembris, MDCCCL. Part I. Introduction, Bibliography and Index A–Aff. pp. 1–128. 1801–1850. Trustees of the British Museum, London, cxxxvi + 128 pp.
- Sherborn CD (1932) Note on de Blainville's 'Manuel d'Actinologie', 1834–1837. *Annals and Magazine of Natural History* (10) 10 (55): 113. <https://doi.org/10.1080/00222933208673545>
- Sherborn CD, Woodward BB (1901a) Notes on the dates of publication of the natural history portions of some French voyages. Part 1. 'Amérique méridionale'; 'Indes orientales'; 'Pôle Sud' ('Astrolabe' and 'Zélée'); 'La Bonite'; 'La Coquille'; and 'L'Uranie et Physicienne'. *Annals and Magazine of Natural History* (7) 7 (40): 388–392.
- Sherborn CD, Woodward BB (1901b) Dates of publication of the zoological and botanical portions of some French voyages.—Part. II. Ferret and Galinier's 'Voyage en Abyssinie'; Lefebvre's 'Voyage en Abyssinie'; 'Exploration scientifique de l'Algérie'; Castelnau's 'Amérique du Sud'; Dumont d'Urville's 'Voyage de l'Astrolabe'; Laplace's 'Voyage sur la Favorite'; Jacquemont's 'Voyage dans l'Inde'; Tréhouart's 'Commission scientifique d'Islande'; Cailliaud, 'Voyage à Méroë'; 'Expédition scientifique de Morée'; Fabre, 'Comission scientifique du Nord'; Du Petit-Thouars, 'Voyage de la Vénus'; and on the Dates of the 'Faune Française.'. *Annals and Magazine of Natural History* (7) 8 (46): 333–336.
- Sherborn CD, Woodward BB (1906) Notes on the dates of publication of the natural history portions of some French voyages.—'Voyage autour du Monde . . . sur . . . la Coquille pendant . . . 1822–25 . . . Par L.J. Duperry [sic].' &c.—A correction. *Annals and Magazine of Natural History* (7) 17 (99): 335–336.
- Sinniger F, Häussermann V (2009) Zoanthids (Cnidaria: Hexacorallia: Zoantharia) from shallow waters of the southern Chilean fjord region, with descriptions of a new genus and two new species. *Organisms Diversity & Evolution* 9 (1): 23–36. <https://doi.org/10.1016/j.ode.2008.10.003>
- Sinniger F, Montoya-Burgos JI, Chevalloné P, Pawlowski J (2005) Phylogeny of the order Zoantharia (Anthozoa, Hexacorallia) based on the mitochondrial ribosomal genes. *Marine Biology* 147 (5): 1121–1128. <https://doi.org/10.1007/s00227-005-0016-3>
- Sinniger F, Reimer JD, Pawlowski J (2008) Potential of DNA sequences to identify zoanthids (Cnidaria: Zoantharia). *Zoological Science* 25: 1253–1260. <https://doi.org/10.2108/zsj.25.1253>

- Sinniger F, Reimer JD, Pawlowski J (2010a) The Parazoanthidae (Hexacorallia: Zoantharia) DNA taxonomy: description of two new genera. *Marine Biodiversity* 40 (1): 57–70. <https://doi.org/10.1007/s12526-009-0034-3>
- Sinniger F, Zelnio KA, Taviani M, Reimer JD (2010b) Presence of *Abyssoanthus* sp. (Anthozoa: Zoantharia) in the Mediterranean Sea: an indication of non-dependence of *Abyssoanthus* to chemosynthetic-based ecosystems? *Cahiers de biologie marine* 51 (4): 475–478.
- Sinniger F, Ocaña OV, Baco AR (2013) Diversity of zoanthids (Anthozoa: Hexacorallia) on Hawaiian seamounts: description of the Hawaiian gold coral and additional zoanthids. *PLoS ONE* 8 (1): e52607 (pp. [1]–[13]). [9 Jan 2013 (p. [1])]
- Soong K, Shiau Y-S, Chen C-P (1999) Morphological and life history divergence of the zoanthid, *Sphenopus marsupialis* off the Taiwanese coast. *Zoological Studies* 38 (3): 333–343.
- Stampar SN, Maronna MN, Kitahara MV, Reimer JD, Morandini AC (2014) Fast-evolving mitochondrial DNA in Ceriantharia: is it a reflection of its phylogenetic position? *PLoS One* 9: e86612. <https://doi.org/10.1371/journal.pone.0086612>
- Stearns REC (1873) Description of a new genus and species of alcyonoid polyp. *Proceedings of the California Academy of Sciences* 5 (1): 147–149, pl. 9. [20 Aug 1873 (separates issued in advance, p. 147)]
- Stechow E (1919) [Extract of] Neue Ergebnisse auf dem Gebiete der Hydroidenforschung. Gesellschaft für Morphologie und Physiologie in München. (Eigener Bericht.) Sitzung vom 15 Juli 1919. *Münchener Medizinische Wochenschrift* 66 (30): 852–853. [25 Jul 1919 (p. 852)]
- Stechow E (1920) Neue Ergebnisse auf dem Gebiete der Hydroidenforschung, Sitzungsberichte der Gesellschaft für Morphologie und Physiologie in München 31: 19–45.
- Steenstrup JJS (1856) *Sphenopus marsupialis* (Gmelin). Oversigt over det Kongelige danske Videnskabernes Selskabs Forhandling og dets Medlemmers Arbejder, 1856: 37–47, 1 pl. [Also issued as a separate with pagination pp. 1–11, 1 pl.]
- Stolarski J, Roniewicz E (2001) Towards a new synthesis of evolutionary relationships and classification of Scleractinia. *Journal of Paleontology* 75 (6): 1090–1108. <https://doi.org/10.1017/S0022336000017157>
- Swain TD (2009) *Isozoanthus antumbrosus*, a new species of zoanthid (Cnidaria: Anthozoa: Zoanthidea) symbiotic with Hydrozoa from the Caribbean, with a key to hydroid and sponge-symbiotic zoanthid species. *Zootaxa* 2051: 41–48. [25 Mar 2009 (p. 41)]
- Swain TD (2010) Evolutionary transitions in symbioses: dramatic reductions in bathymetric and geographic ranges of Zoanthidea coincide with loss of symbioses with invertebrates. *Molecular Ecology* 19 (12): 2587–598. <https://doi.org/10.1111/j.1365-294x.2010.04672.x>
- Swain TD, Swain LS (2014) Molecular parataxonomy as taxon description: examples from recently named Zoanthidea (Cnidaria: Anthozoa) with revision based on serial histology of microanatomy. *Zootaxa* 3796 (1): 81–107. <https://doi.org/10.11646/zootaxa.3796.1.4>
- Swain TD, Schellinger JL, Strimaitis AM, Reuter KE (2015) Evolution of anthozoan polyp retraction mechanisms: convergent functional morphology and evolutionary allometry of the marginal musculature in order Zoanthidea (Cnidaria: Anthozoa: Hexacorallia). *BMC Evolutionary Biology* 15 (123): 1–19. <https://doi.org/10.1186/s12862-015-0406-1>

- Swain TD, Strimaitis AM, Reuter KE, Boudreau W (2016) Towards integrative systematics of Anthozoa (Cnidaria): evolution of form in the order Zoanthidea. *Zoologica Scripta*. <https://doi.org/10.1111/zsc.12195>
- Tilesius GT (1809) De nova Actiniarum specie gigantea, Kamtschatica, quam in Portu Petro-Pauli ad vivum pinxit. *Memoires de l'Académie Impériale des Sciences de St. Pétersbourg* 1: 388–422, pls. 14–15.
- Untawale AG, Dhargalkar VK (2002) Marine conservation strategies for Maharashtra coast. In: Sahoo D, Pandey PC (eds). *Advances in Marine and Antarctic Science*. A.P.H. Publishing Corporation, New Delhi, pp. 107–120.
- Verrill AE (1865a) Classification of polyps: (extract condensed from a Synopsis of the Polypi of the North Pacific Exploring Expedition, under Captains Ringgold and Rodgers, U.S.N.). *Proceedings of the Essex Institute* 4: 145–152. [April 1865 (Fautin 2016: 446); reprinted as Verrill (1865b)]
- Verrill AE (1865b) Classification of polyps. (Extract condensed from a Synopsis of the Polypi of the North Pacific Exploring Expedition under Captains Ringgold and Rodgers, U.S.N.). *Annals and Magazine of Natural History* (3) 16 (93): 191–197. [1 Sep 1865 (Evenhuis 2003b: 23); reprint of Verrill (1865a)]
- Verrill AE (1868–1871) Review of the corals and polyps of the West Coast of America. *Transactions of the Connecticut Academy of Arts and Sciences* 1 (2): 377–567, 611–613, pls. 5–10. [Apr ? 1868 (pp. 377–390), Jun ? 1868 (pp. 391–398), Jul ? 1868 (pp. 399–414), Dec ? 1868 (pp. 415–422), Jan ? 1869 (pp. 423–454), Feb ? 1869 (pp. 455–478), Mar ? 1869 (pp. 479–502), Apr ? 1870 (pp. 503–518), Nov ? 1870 (pp. 519–534), Dec ? 1870 (pp. 535–558), Feb ? 1871 (pp. 559–567), Mar ? 1871 (pp. 611–613) (year-dates from Fisher (1893: 133–135); month-dates from page signatures; see also unnumbered prefatory note in Verrill 1871); reprinted with changes as Verrill (1869–1871)]
- Verrill AE (1869–1871) Notes on Radiata in the Museum of Yale College. From the *Transactions of the Connecticut Academy of Arts and Sciences*, Vol. I, March, 1871. Tuttle, Morehouse & Taylor, New Haven, pp. 247–567, 611–613, pls. 5–10. [after 15 Nov 1869 (pp. 247–502) (unnumbered preface); Apr ? 1870 (pp. 503–518), Nov ? 1870 (pp. 519–534), Dec ? 1870 (pp. 535–558), Feb ? 1871 (pp. 559–567), Mar ? 1871 (pp. 611–613) (dates of pp. 247–502 from unnumbered preface; remaining dates follow those for Verrill, 1868–1870); reprint with changes of Verrill (1868–1871) (see unnumbered preface)]
- Verrill AE (1870) Descriptions of new corals. *Contributions to Zoölogy from the Museum of Yale College*. No. 7. *American Journal of Science and Arts* (2) 49 (147): 370–375.
- Verrill AE (1882) Notice of the remarkable marine fauna occupying the outer banks off the southern coast of New England, No. 3. *American Journal of Science* (3) 23 (134): 135–142. [Apr 1882 (Fautin 2016: 447); *Epizoanthus paguriphilus* first made available in this publication; full description in Verrill (1883)]
- Verrill AE (1883) Report on the Anthozoa, and on some additional species dredged by the 'Blake' in 1877–79, and by the U.S. Fish Commission Steamer 'Fish Hawk' in 1880–82. In: *Reports on the results of dredging, under the supervision of Alexander Agassiz, on the east coast of the United States, during the summer of 1880, by the U.S. Coast Survey*

- Steamer 'Blake,' Commander J.R. Bartlett, U.S.N., commanding. Bulletin of the Museum of Comparative Zoölogy, Harvard College 11 (1): 1–72, pls. 1–8. [Jul 1883 (title-page); the name *Epizoanthus paguriphilus* was first made available in Verrill (1882)]
- Verrill AE (1900) Additions to the Anthozoa and Hydrozoa of the Bermudas. Transactions of the Connecticut Academy of Arts and Sciences 10 (2): 551–572, pls. 67–69. <https://doi.org/10.5962/bhl.part.7033>
- Verrill AE (1907) The Bermuda Islands. Part IV.—Geology and Paleontology, and Part V.—An Account of the Coral Reefs. Transactions of the Connecticut Academy of Arts and Sciences 12 (2), 45–347, pls. 16–40.
- Volpi C, Benvenuti D (2003) The Duchassaing & Michelotti collection of Caribbean corals: status of the types and location of the specimens. Atti della Societa Italiana di Scienze Naturali e del Museo Civico di Storia Naturale de Milano 144 (1): 51–74. [Jun 2003 (p. 51)]
- Walsh GE (1967) An Annotated Bibliography of the Families Zoanthidae, Epizoanthidae, and Parazoanthidae (Coelenterata, Zoantharia). Hawaii Institute of Marine Biology, Honolulu, 77 pp. [Mar 1967 (title-page)]
- Williams RB (1981a) A sea anemone, *Edwardsia meridionalis* sp. nov., from Antarctica and a preliminary revision of the genus *Edwardsia* de Quatrefages, 1841 (Coelenterata: Actiniaria). Records of the Australian Museum 33 (6): 325–360. <https://doi.org/10.3853/j.0067-1975.33.1981.271>
- Williams RB (1981b) The authorship of the family name Metridiidae (Coelentera: Anthozoa). Bulletin of Zoological Nomenclature 38 (3): 156–157. <https://doi.org/10.5962/bhl.part.8185>
- Williams RB (2000) A redescription of the zoanthid *Isozoanthus sulcatus* (Gosse, 1859), with notes on its nomenclature, systematics, behaviour, habitat and geographical distribution. Ophelia 52 (3): 193–206. <https://doi.org/10.1080/00785236.1999.10409428>
- Williams RB (2012) Identification and dating of the editions and issues of 'Le Attinie', a taxonomic monograph of the world's sea anemones (Cnidaria: Hexacorallia), by Angelo Andres. Zoological Bibliography 2 (2–3): 90–104. [26 Nov 2012 (title-page)]
- Yang S-Y, Reimer JD, Low MEY (2014) Die *Pflanzen-thiere in Abbildungen nach der Natur* and its *Fortsetzungen* by Eugenius Johann Christoph Esper: overlooked sources for its dates of publication and location of type material. Sherbornia 1 (2): 5–7. [10 Dec 2014 (p. 5)]

Appendix I

Nomenclator of family-group names in the Zoantharia Rafinesque, 1815

Names in **bold** and designated with an asterisk (*) are considered to be valid family-group names. Also refer to the genus-group names in the next section. Names marked with a † are invalid synonyms, or incorrect spellings (if further denoted by a “[sic]”). Additional names without either designation have been confused with the order Zoantharia at one time or another but have since been removed from this group.

***Abyssoanthidae** Reimer and Fujiwara, in Reimer, Sinniger, Fujiwara, Hirano and Maruyama, 2007: 258. Type genus *Abyssoanthus* Reimer and Fujiwara, in Reimer, Sinniger, Fujiwara, Hirano and Maruyama, 2007.

Bergidae Verrill, 1865. See Bergiidae Verrill, 1865.

†Bergidae [*sic*]. Incorrect subsequent spelling of Bergiidae Verrill, 1865 (e.g. Verrill 1869 [in 1868–1871]: 494; Duerden 1903: 495).

†Bergiidae Verrill, 1865a: 147. Type genus *Bergia* Duchassaing de Fombressin and Michelotti, 1860. This family-group name was first established as “Bergidae” (an incorrect original spelling. *Bergia* Duchassaing de Fombressin and Michelotti, 1860, was previously considered to be a junior subjective synonym of *Parazoanthus* Haddon and Shackleton, 1891, but has since been revalidated (see Montenegro et al. 2015a: 63–71). Bergiidae Verrill, 1865, and Parazoanthidae Delage and Hérouard, 1901, nevertheless remain subjective synonyms. Low and Reimer (2011a: 64, 65) enacted Article 23.9 (ICZN 1999: 27, 28) to reverse precedence between these family-group names, thereby making Bergiidae Verrill, 1865, a *nomen oblitum*, and Parazoanthidae Delage and Hérouard, 1901, a *nomen protectum*. Low and Reimer (2011a: 64, 65) incorrectly attributed the name Bergiidae to Verrill, 1869 (in 1868–1871), but the name should be attributed to Verrill (1865). The findings and action of Low and Reimer (2011: 64, 65) remain valid.

***Epizoanthidae** Delage and Hérouard, 1901: 664. Type genus *Epizoanthus* Gray, 1867. First described as the subfamily “Epizoanthinae”. See also Mardoellidae Danielssen, 1890.

Gerardiidae Verrill, 1865. See Gerardiidae Verrill, 1865a.

†Gerardiidae [*sic*]. Incorrect subsequent spelling of Gerardiidae Verrill, 1865 (e.g., Verrill 1869 [in 1868–1871]: 499; Duerden 1903: 495).

†Gerardiidae Verrill, 1865a: 1484. Type genus *Gerardia* Lacaze-Duthiers, 1864. This family-group name was first established as “Gerardiidae” (an incorrect original spelling). A subjective synonym of Parazoanthidae Delage and Hérouard, 1901. This family group name is a senior subjective synonym of Parazoanthidae Delage and Hérouard, 1901, and an application to the ICZN in accordance with Article 23.9.2 (ICZN 1999: 28, 29) is in preparation to suppress this name in favour of the more universally-accepted and -used name Parazoanthidae Delage and Hérouard, 1901. Also see discussion in Appendix 3.

†Heterozoanthidae Pax and Müller, 1956: 2. Type genus *Heterozoanthus* Verrill, 1870. The type genus is an objective synonym of *Parazoanthus* Haddon and Shackleton, 1891, making Heterozoanthidae Pax and Müller, 1956, a junior objective synonym of Parazoanthidae Delage and Hérouard, 1901 (Article 61.3.2 of the Code, ICZN 1999: 64) (see Low and Reimer 2012a: 83, 84).

***Hydrozoanthidae** Sinniger, Reimer and Pawlowski, 2010: 60. Type genus *Hydrozoanthus* Sinniger, Reimer and Pawlowski, 2010.

†Mardoellidae Danielssen, 1890: 116, 117. Type genus *Mardoell* Danielssen, 1890. The type genus is a subjective synonym of *Epizoanthus* Gray, 1867 (see Low and

Reimer, 2011b: 84, 85; Lwowsky, 1913: 603, 604). Mardoellidae Danielssen, 1890, is therefore a subjective synonym of Epizoanthidae Delage and Hérouard, 1901. Low and Reimer (2011b: 84) enacted Article 23.9 (ICZN 1999: 27, 28) to reverse precedence between these family-group names, thereby making Mardoellidae Danielssen, 1890, a *nomen oblitum*, and Epizoanthidae Delage and Hérouard, 1901, a *nomen protectum*.

***Microzoanthidae** Fujii and Reimer, 2011: 420, 421. Type genus *Microzoanthus* Fujii and Reimer, 2011.

***Nanozoanthidae** Fujii and Reimer, 2013: 512. Type genus *Nanozoanthus* Fujii and Reimer, 2013.

***Neozoanthidae** Herberts, 1972: 137. Type genus *Neozoanthus* Herberts, 1972.

Orinidae Verrill, 1869. See Oriniidae Verrill, 1869.

Oriniidae Verrill, 1869 (in 1868–1871): 494. Type genus *Orinia* Duchassaing de Fonbressin and Michelotti, 1860. This family-group name was first established as “Orinidae” (an incorrect original spelling). The type genus *Orinia* Duchassaing de Fonbressin and Michelotti, 1860, is a junior subjective synonym of *Rhodactis* Milne Edwards and Haime, 1851, which is currently assigned to the family Discosomidae Verrill, 1869 [Coralliomorpha] (see Fautin 2016: 25, 28, 38). Also see remarks under *Orinia* Duchassaing de Fonbressin and Michelotti, 1860, in Appendix 2.

Polythoae Duchassaing de Fonbressin and Michelotti, 1860. See Palythoidae Duchassaing de Fonbressin and Michelotti, 1860.

Palythoidae Duchassaing de Fonbressin and Michelotti, 1860: 37. Type genus *Palythoa* Lamouroux, 1816 (spelled as “*Polythoa*”). This family-group name was first established as “Polythoae” (based on “*Polythoa*”, an incorrect subsequent spelling of *Palythoa* Lamouroux, 1816, see Appendix 3). A subjective synonym of Sphenopidae Hertwig, 1882, as the type species *Sphenopus* Steenstrup, 1856, and *Palythoa* Lamouroux, 1816, as assigned to the same family (see Appendix 3). Precedence between Palythoidae Duchassaing de Fonbressin and Michelotti, 1860, and Sphenopidae Hertwig, 1882, cannot be reversed and an application to the ICZN in accordance with Article 23.9.2 (ICZN, 1999: 28, 29) is in preparation to suppress the former name in favour of the latter name that is more widely accepted and used. Also see discussion in Appendix 3.

***Parazoanthidae** Delage and Hérouard, 1901: 665. Type genus *Parazoanthus* Haddon and Shackleton, 1891. See also the synonyms Bergiidae Verrill, 1865, Gerardiidae Verrill, 1865, Heterozoanthidae Pax and Müller, 1956, Savagliidae Brook, 1889, and Savaliidae Nardo, 1844. Also see discussion in Appendix 3.

†Savagliidae Brook, 1889: 51, 74, 79. Type genus *Savaglia* Nardo, 1877. A subjective synonym of Parazoanthidae Delage and Hérouard, 1901. This family group name is a senior subjective synonym of Parazoanthidae Delage and Hérouard, 1901, and an application to the ICZN in accordance with Article 23.9.2 (ICZN, 1999: 28, 29) is in preparation to suppress this name in favour of the more widely accepted and used name Parazoanthidae Delage and Hérouard, 1901. Also see discussion in Appendix 3.

- †**Savaliidae** Nardo, 1844: 433. Type genus *Savalia* Nardo, 1844. This family-group name was first established at the rank of subfamily as “Savalini” (an incorrect original spelling). A subjective synonym of Parazoanthidae Delage and Hérouard, 1901. This family group name is a senior subjective synonym of Parazoanthidae Delage and Hérouard, 1901, and an application to the ICZN in accordance with Article 23.9.2 (ICZN, 1999: 28, 29) is in preparation to suppress this name in favour of the more universally-accepted and -used name Parazoanthidae Delage and Hérouard, 1901. Also see discussion in Appendix 3.
- †**Savaliidae** Poche, 1914: 104. Type genus *Savalia* Nardo, 1844. Proposed as a replacement name for Savagliidae Brook, 1889. A junior objective synonym of Savaliidae Nardo, 1844, and Savagliidae Brook, 1889.
- Savalini Nardo, 1844. See Savaliidae Nardo, 1844.
- ***Sphenopidae** Hertwig, 1882: 120. Type genus *Sphenopus* Steenstrup, 1856. Zoanthia Rafinesque, 1815. See Zoanthidae Rafinesque, 1815.
- ***Zoanthidae** Rafinesque, 1815: 155. Type genus *Zoanthus* Lamarck, 1801. First established at the rank of subfamily as “Zoanthia” (an incorrect original spelling). Authorship of the family-group name Zoanthidae is conventionally attributed to Gray (1832: 95; 1840: 72), however Rafinesque’s (1815) indication has priority (see Low and Reimer 2012a: 85).

Appendix 2.

Nomenclator of genus-group names in the Zoantharia Rafinesque, 1815

Names in **bold** and designated with an asterisk (*) are considered to be valid genus-group names. In accordance with Article 67.2 of the Code (ICZN 1999: 66, 67), all type species designations made herein are from among the “originally included nominal species”. Names marked with a † are invalid synonyms, or incorrect spellings (if further denoted by a “[sic]”). Additional names without either designation have been confused with the order Zoantharia at one time or another but have since been removed from this group. The identity of two possible zoantharian genus-group name-*Stephanidium* Hertwig, 1888, and *Triga* Gray, 1867-will need to be resolved.

- ****Abyssoanthus*** Reimer and Fujiwara, in Reimer, Sinniger, Fujiwara, Hirano and Maruyama, 2007: 258. Type species *Abyssoanthus nankaiensis* Reimer and Fujiwara, in Reimer, Sinniger, Fujiwara, Hirano and Maruyama, 2007, by original designation; gender masculine.
- ****Acrozoanthus*** Saville-Kent, 1893: 153, 154, unnumbered fig. Type species *Acrozoanthus australiae* Saville-Kent, 1893, by monotypy; gender masculine.
- †***Actimastus*** Rafinesque, 1818: 271. Replacement name for *Mammillifera* Le Sueur, 1817. Type species *Mammillifera auricula* Le Sueur, 1817, by subsequent designation by Haddon and Shackleton (1891a: 626) (for *Mammillifera* Le Sueur,

- 1817); gender masculine. Unnecessary replacement name for and junior objective synonym of *Mammilifera* Le Sueur, 1817. Both *Actimastus* Rafinesque, 1818, and *Mammilifera* Le Sueur, 1817, are junior subjective synonyms of *Zoanthus* Lamarck, 1801, as the type species of both genus-group names is now considered to be a species of *Zoanthus* Lamarck, 1801 (see Appendix 3).
- Anthozoanthus* Carter, 1870: 449. Type species *Anthozoanthus parasiticus*, by monotypy; gender masculine. Not a zoantharian (see Carter 1870: 449). This name appears to be Deshayes manuscript name that was published by Carter (1870).
- †*Anthozoon* Gistel, 1848: 181. Unnecessary replacement name and junior objective synonym of *Zoanthus* Lamarck, 1801; gender neuter.
- †*Antinedia* Duchassaing de Fonbressin and Michelotti, 1864: 42 (also 1866: 136). Type species *Zoanthus tuberculatus* Duchassaing de Fonbressin, 1850, by monotypy; gender feminine. The type species *Zoanthus tuberculatus* Duchassaing de Fonbressin, 1850, is a junior subjective synonym and junior secondary homonym of *Isaurus tuberculatus* Gray, 1828. A subjective junior synonym of *Isaurus* Gray, 1828 (see Muirhead and Ryland 1985: 325).
- †*Actinorhiza* Agassiz, 1846: 7. Unjustified emendation and junior objective synonym of *Actinorhyza* Blainville, 1830. See Appendix 3.
- †*Actinorhysa* [sic]. Incorrect subsequent spelling of *Actinorhyza* Blainville, 1830, by Blainville (1834: 329). See Appendix 3.
- † *Actinorhyza* Blainville, 1830: 295. Type species *Actinia sociata* Ellis, 1768 (see Appendix 3); gender feminine. An unnecessary replacement name for, and junior objective synonym of *Zoanthus* Lamarck, 1801. See also the incorrect subsequent spellings *Actinorhysa* Blainville, 1834, and *Actinorrhyza* Ehrenberg, 1834, and the unjustified emendation *Actinorhiza* Agassiz, 1844. See additional discussion in Appendix 3.
- †*Actinorrhyza* [sic]. Incorrect subsequent spelling of *Actinorhyza* Blainville, 1830, by Ehrenberg (1834a: 269). See Appendix 3.
- Actinocereus* Blainville, 1830: 294. No type species designated (see Fautin et al. 2012: 20, 21); gender masculine. Placed in synonymy (in part) of *Hughea* Lamouroux, 1821, by Andres (1881: 336). Currently a genus-group name in the Actiniaria (see Fautin et al. 2012: 20–22; Fautin 2016: 54). *Actinocereus* Blainville, 1830, is now considered a synonym of *Cereus* Ilmoni, 1830, and the International Commission for Zoological Nomenclature was petitioned Fautin et al. (2012: 22) to suppress the former name in favour of the latter, and subsequently ruled as such (ICZN 2014: 136, 137).
- **Antipathozoanthus* Sinniger, Reimer and Pawlowski, 2010: 61. Type species *Gerardia macaronesicus* Ocaña and Brito, 2003, by original designation; gender masculine.
- Axinella* O. Schmidt, 1862: 60. Type species *Axinella polypoides* O. Schmidt, 1862 (see Gazave et al. 2010). Not a zoantharian (see Gazave et al. 2010).
- **Bergia* Duchassaing de Fonbressin and Michelotti, 1860: 54. Type species *Bergia catenularis* Duchassaing de Fonbressin and Michelotti, 1860, by subsequent designation by Duerden (1903: 496); gender feminine. The type species was formerly assigned to the genus *Parazoanthus* Haddon and Shackleton, 1891a (see Duerden

- 1903: 496, Low and Reimer 2011a: 64), making *Bergia* Duchassaing de Fombressin and Michelotti, 1860, a subjective synonym of *Parazoanthus* Haddon and Shackleton, 1891a. Low and Reimer (2011a) enacted Article 23.9 (ICZN 1999: 27, 28) to reverse precedence between these genus-group names, thereby making *Bergia* Duchassaing de Fombressin and Michelotti, 1860, a *nomen oblitum*, and *Parazoanthus* Haddon and Shackleton, 1891a, a *nomen protectum*. Recent molecular and morphological work by Montenegro et al. (2015a: 63–71) have shown that the type species, *Bergia catenularis* Duchassaing de Fombressin and Michelotti, 1860, represents a generic-level monophyly and resurrected the genus-group name *Bergia* Duchassaing de Fombressin and Michelotti, 1860, for this grouping.
- ****Bullagummizoanthus*** Sinniger, Ocaña and Baco, 2013: 9. Type species *Bullagummizoanthus emilyacadiarum* Sinniger, Ocaña and Baco, 2013, by original designation; gender masculine.
- †***Carolia*** Gray, 1867: 239. Type species *Zoanthus couchii* Johnston, in Couch, 1844, by monotypy; gender feminine. The type species is now assigned to the genus *Epizoanthus* Gray, 1867 (see Haddon and Shackleton 1891a: 644, 645), making *Carolia* Gray, 1867, a subjective synonym of *Epizoanthus* Gray, 1867. The name *Carolia* Gray, 1867, is an invalid junior homonym of *Carolia* Cantraine, 1838 (Mollusca), and the *Epizoanthus* Gray, 1867, is the valid name for this genus-group. See also *Epizoanthus* Gray, 1867.
- †***Cavolinia*** Schweigger, 1819: 99. No type species designated, for originally included species, see Appendix 3; gender feminine. *Cavolinia* Schweigger, 1819, is a junior homonym of *Cavolinia* Abildgaard, 1791, and is not a valid name (see Appendix 3). See also the unnecessary replacement name *Cynicus* Gistel, 1848.
- ****Corallizoanthus*** Reimer, in Reimer, Nonaka, Sinniger and Iwase, 2008: 940. Type species *Corallizoanthus tsukaharai* Reimer, in Reimer, Nonaka, Sinniger and Iwase, 2008, by original designation; gender masculine.
- †***Corticanthus*** Andres, 1883a: 538, 541. Type species *Epizoanthus paguriphilus* Verrill, 1882, herein designated (see Appendix 3); gender masculine. First proposed as a subgenus of *Zoanthus* Lamarck, 1801. The type species is now assigned to the genus *Epizoanthus* Gray, 1867 (see Appendix 3), making *Zoanthus* (*Corticanthus*) Andres, 1883, a junior subjective synonym of *Epizoanthus* Gray, 1867.
- †***Corticifera*** Le Sueur, 1817: 178, 179. Type species *Corticifera glareola* Le Sueur, 1817, by subsequent designation by Haddon and Shackleton (1891b: 692); gender feminine. The type species is now assigned to the genus *Palythoa* Lamouroux, 1816 (see Appendix 3), making *Corticifera* Le Sueur, 1817, a junior subjective synonym of *Palythoa* Lamouroux, 1816.
- †***Corticithoa*** Andres, 1883a: 521, 535–538. Type species *Alyconium tuberculosum* Esper, 1805, herein designated (see Appendix 3); gender feminine. First proposed as a subgenus of *Palythoa* Lamouroux, 1816. The type species, *Alyconium tuberculosum*, is now assigned to the genus *Palythoa* Lamouroux, 1816 (see Appendix 3), making *Palythoa* (*Corticithoa*) Andres, 1883, a junior subjective synonym of *Palythoa* Lamouroux, 1816.

- †*Cortificera*. Incorrect spelling of *Corticifera* Le Sueur, 1817, by Brandt (1835: 208).
- †*Cynicus* Gistel, 1848: viii. Unnecessary replacement name for, and junior objective synonym of, *Cavolinia* Schweigger, 1819; gender masculine.
- Edwardsia* de Quatrefages, 1841: 427. Type species *Edwardsia beautempsii* de Quatrefages, 1842, by subsequent designation by Delphy (1938) (see Fautin 2016: 82); gender feminine. Not a zoantharian (see Fautin et al. 2007: 201, Williams 1981a: 326; Fautin 2016: 82, 83).
- †*Endeithoa* Andres, 1883a: 521, 531. Type species *Zoanthus norvegicus* Koren and Danielssen, 1877, herein designated (see Appendix 3); gender feminine. First proposed as a subgenus of *Palythoa* Lamouroux, 1816. The type species is now assigned to the genus *Epizoanthus* Gray, 1867 (see Appendix 3), making *Palythoa* (*Endeithoa*) Andres, 1883, a junior subjective synonym of *Epizoanthus* Gray, 1867.
- Epiactis* Verrill, 1869 (in 1868–1871): 492. Type species *Epiactis prolifera* Verrill, 1869, by monotypy; gender feminine. Not a zoantharian (Fautin et al. 2007: 202; Fautin 2016: 85).
- **Epizoanthus* Gray, 1867: 237. Type species *Dysidea papillosa* Johnston, 1842, by monotypy; gender masculine. Placed on *Official List* and has priority over *Sidisia* Gray, 1858 (see Opinion 1689, ICZN 1992: 236). The subsequent designation of *Mammillifera incrustatus* Düben and Koren, 1847, as the type species of *Epizoanthus* Gray, 1867, by Haddon and Shackleton (1891a: 627) is not valid as this species was not amongst the species originally included in the description of *Epizoanthus* Gray, 1867, and does not qualify for selection as the type species (see Ryland and Muirhead 1991: 19, 20; and Article 69.2 of the Code, ICZN 1999: 72, 73). See also the synonyms *Carolia* Gray, 1867, *Lirrevia* Delphy, 1939: 270, *Mardoell* Danielssen, 1890, *Marodellia* Danielssen, 1890, *Palythoa* (*Endeithoa*) Andres, 1883, *Sidisia* Gray, 1858, *Verrillia* Andres, 1883, and *Zoanthus* (*Corticanthus*) Andres, 1883,
- †*Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860: 55. Type species *Gemmaria rusei* Duchassaing de Fonbressin and Michelotti, 1860, by subsequent designation by Haddon and Shackleton (1891a: 626). The genus-group name *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860, is preoccupied by *Gemmaria* McCrady, 1859 (Hydrozoa). Herein considered to be a junior subjective synonym of *Palythoa* Lamouroux, 1816 (See Appendix 3). See also the replacement names *Haplotella* Stechow, 1919, and *Protopalythoa* Verrill, 1900.
- †*Gemmithoa* Andres, 1883a: 521, 532, 533. Type species *Mammillifera brevis* Duchassaing de Fonbressin, 1850, by monotypy; gender feminine. First proposed as a subgenus of *Palythoa* Lamouroux, 1816. The type species is now assigned to the genus *Palythoa* Lamouroux, 1816 (see Appendix 3), making *Palythoa* (*Gemmithoa*) Andres, 1883, a junior subjective synonym of *Palythoa* Lamouroux, 1816.
- †*Gerardia* Lacaze-Duthiers, 1864a: 87. Type species *Leoipathes lamarcki* Haime, 1849, by monotypy; gender feminine. A subjective junior synonym of *Savalia* Nardo, 1844 (see Appendix 3). This genus-group name was also described as new in Lacaze-Duthiers (1864c: 175, 176), and appeared as a translation in Lacaze-Duthiers (1864b: 242).

- †*Gerardina* [sic]. Incorrect spelling of *Gerardia* Lacaze-Duthiers, 1864, by H. Schmidt (1972: 452).
- †*Haplotella* Stechow, 1919: 853. Replacement name for *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860, preoccupied by *Gemmaria* McCrady, 1859 (Hydrozoa). Type species *Gemmaria rusei* Duchassaing de Fonbressin and Michelotti, 1860, by typification of the preoccupied name; gender feminine. A junior objective synonym of *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860, and a junior subjective synonym of *Palythoa* Lamouroux, 1816 (see Appendix 3). See also *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860, *Parapalythoa* Verrill, 1900, and *Protopalythoa* Verrill, 1900.
- †*Heterozoanthus* Verrill, 1870: 371. Type species *Palythoa axinellae* O. Schmidt, 1862, by subsequent designation by Low and Reimer (2012a: 84); gender masculine. As *Heterozoanthus* Verrill, 1870, and *Parazoanthus* Haddon and Shackleton, 1891, have the same type species, they are objective synonyms. Low and Reimer (2012a: 84) reversed precedence thereby making *Heterozoanthus* Verrill, 1870, a *nomen oblitum*, and *Parazoanthus* Haddon and Shackleton, 1891, a *nomen protectum*.
- Hughaea* [sic]. Incorrect spelling of *Hughea* Lamouroux, 1821, by Duchassaing de Fonbressin (1850: 9). Not a zoantharian (see *Hughea* Lamouroux, 1821).
- Hughea* Lamouroux, 1821: 89. Type species *Actinia calendula* Hughes, in Ellis and Solander, 1786, by monotypy; gender feminine. The type species, *Actinia calendula* was described by Ellis, in Ellis and Solander (1786: 7, 8, pl. 1, fig. 3) based on an unnamed cnidarian described and illustrated by Hughes (1750: 293, 294, pl. 24, fig. 1) that is clearly not a zoantharian (see also Ryland and Lancaster 2003: 411; Fautin 2016: 182). *Actinia calendula* Ellis, in Ellis and Solander, 1786, is currently considered to be assigned to the genus *Petalactis* Andres, 1883 (see Fautin 2016: 118), thereby making *Hughea* Lamouroux, 1821, a synonym of *Petalactis* Andres, 1883. Also see the unnecessary replacement name *Meto* Gistel, 1848.
- Hughuea* [sic]. Incorrect spelling of *Hughea* Lamouroux, 1821, by Gistel (1848: 181). Not a zoantharian (see *Hughea* Lamouroux, 1821).
- ****Hurlizoanthus*** Sinniger, Ocaña and Baco, 2013: 7. Type species *Hurlizoanthus parishi* Sinniger, Ocaña and Baco, 2013, by original designation; gender masculine.
- ****Hydrozoanthus*** Sinniger, Reimer and Pawlowski, 2010: 60. Type species *Parazoanthus tunicans* Duerden, 1900, by original designation; gender masculine.
- Iluanthos* Forbes, 1840: 184. Type species *Iluanthos scoticus* Forbes, 1840, by monotypy; gender masculine. Fautin (2016: 97) considers the correct original spelling of this genus-group name to be “Ilyanthus” following Article 33.3.1 of the Code (ICZN 1999: 43). Not a zoantharian (Fautin et al. 2007: 209; Fautin 2016: 97).
- †*Isaua* [sic]. Incorrect subsequent spelling of *Isaurus* Gray, 1828, by Volpi and Benvenuti (2003: 72).
- †*Isaura* Agassiz, 1844: 14. Unjustified emendation and objective synonym of *Isaurus* Gray, 1828, and junior homonym of *Isaura* Lamouroux, in Audouin, Bourdon, de Candolle, d’Aubebard de Férussac, Deshayes, Deslongchamps, É. Geoffroy Saint-Hilaire, I. Geoffroy Saint-Hilaire, Guérin, Guillemin, de Jussieu, Kunth, Dela-

fosse, Lamouroux, Latreille, Prévost, Richard and Bory de Saint-Vincent, 1826. See *Isaurus* Gray, 1828.

†*Isaura* Lamouroux, in Audouin, Bourdon, de Candolle, d'Aubebard de Férussac, Deshayes, Deslongchamps, É. Geoffroy Saint-Hilaire, I. Geoffroy Saint-Hilaire, Guérin, Guillemain, de Jussieu, Kunth, Delafosse, Lamouroux, Latreille, Prévost, Richard and Bory de Saint-Vincent, 1826: 23. Type species *Isaurus tuberculatus* Gray, 1828, by subsequent designation by Haddon and Shackleton (1891b: 682); gender masculine. An objective synonym of *Isaurus* Gray, 1828 (see Low and Reimer 2012b: 46). See *Isaurus* Gray, 1828.

**Isaurus* Gray, 1828: 8. Type species *Isaurus tuberculatus* Gray, 1828, by subsequent designation by Haddon and Shackleton (1891b: 682); gender masculine. The name *Isaurus* Gray, 1828, is a replacement name for *Isaura* Lamouroux, in Audouin, Bourdon, de Candolle, d'Aubebard de Férussac, Deshayes, Deslongchamps, É. Geoffroy Saint-Hilaire, I. Geoffroy Saint-Hilaire, Guérin, Guillemain, de Jussieu, Kunth, Delafosse, Lamouroux, Latreille, Prévost, Richard and Bory de Saint-Vincent, 1826, which Gray (1828: 8) believed to be preoccupied (see Low and Reimer 2012b: 46). Low and Reimer (2012b: 47) reversed the precedence of *Isaura* Lamouroux, in Audouin, Bourdon, de Candolle, d'Aubebard de Férussac, Deshayes, Deslongchamps, É. Geoffroy Saint-Hilaire, I. Geoffroy Saint-Hilaire, Guérin, Guillemain, de Jussieu, Kunth, Delafosse, Lamouroux, Latreille, Prévost, Richard and Bory de Saint-Vincent, 1826, and *Isaurus* Gray, 1828, thereby maintaining current and prevailing usage of the latter name. See also the synonyms *Antinedia* Duchassaing de Fombressin and Michelotti, 1864, *Pales* Gray, 1867, *Palythoa* (*Monothoa*) Andres, 1883, *Panceria* Andres, 1877, and *Zoanthus* (*Monanthus*) Andres, 1883.

**Isozoanthus* Carlgren, in Chun, 1903: 520. Type species *Isozoanthus giganteus* Carlgren, in Chun, 1903, by monotypy; gender masculine. Williams (2000) enacted Article 23.9 of the Code (ICZN 1999: 27, 28) to reverse precedence of *Palythoa* (*Taeniothoa*) Andres, 1883, and *Isozoanthus* Carlgren, in Chun, 1903. See also *Palythoa* (*Taeniothoa*) Andres, 1883.

**Kauluzoanthus* Sinniger, Ocaña and Baco, 2013: 8. Type species *Kauluzoanthus kerbyi* Sinniger, Ocaña and Baco, 2013, by original designation; gender masculine.

**Kulamanamana* Sinniger, Ocaña and Baco, 2013: 4. Type species *Kulamanamana haumaeae* Sinniger, Ocaña and Baco, 2013, by original designation; gender feminine.

†*Lirrevia* Delphy, 1939: 270. Replacement name for *Verrillia* Andres, 1883; gender feminine. *Lirrevia* Delphy, 1939, and *Verrillia* Andres, 1883, are thus subjective junior synonyms of *Epizoanthus* Gray, 1867. See also *Verrillia* Andres, 1883.

†*Mamillifera* [sic]. Incorrect spelling of *Mammillifera* Le Sueur, 1817, by Blainville (1830: 295).

†*Mammilifera* [sic]. Incorrect spelling of *Mammillifera* Le Sueur, 1817, by Gistel (1848: 181).

†*Mammillifera* Le Sueur, 1817: 177. Type species *Mammillifera auricula* Le Sueur, 1817, by subsequent designation by Haddon and Shackleton (1891a: 626); gender

- feminine. The type species is now assigned to the genus *Zoanthus* Lamarck, 1801 (see Appendix 3), making *Mammillifera* Le Sueur, 1817, a junior subjective synonym of *Zoanthus* Lamarck, 1801. See also *Actimastus* Rafinesque, 1818.
- †*Mammithoa* Andres, 1883a: 521. Type species *Mammillifera nymphaea* Le Sueur, 1817, herein designated (see Appendix 3); gender feminine. First proposed as a subgenus of *Palythoa* Lamouroux, 1816. The type species is now a junior subjective synonym of *Zoanthus pulchellus* Duchassaing de Fonbressin and Michelotti, 1864, which is currently assigned to the genus *Zoanthus* Lamarck, 1801 (see Appendix 3), making *Palythoa (Mammithoa)* Andres, 1883, a junior subjective synonym of *Zoanthus* Lamarck, 1801. See also the incorrect original spelling *Mammothoa* Andres, 1883.
- †*Mammothoa* [sic]. An incorrect original spelling of *Mammithoa* Andres, 1883, by Andres, (1883a: 533–535; 1883b: 325–327; 1884: 318–320) (see Appendix 3).
- †*Mardoell* Danielssen, 1890: 117–126. Type species *Mardoell erdmanni* Danielssen, 1890, by monotypy; gender feminine. The type species is assigned to the genus *Epizoanthus* Gray, 1867 (see Low and Reimer 2012a: 85, Lwowsky, 1913: 603, 604), making *Mardoell* Danielssen, 1890, a junior subjective synonym of *Epizoanthus* Gray, 1868. See also the incorrect emendation *Mardoellia* Blanchard, 1893.
- †*Mardoella* [sic]. Incorrect spelling of *Mardoell* Danielssen, 1890, by Bell (1906: 762) and also in Neave's (1940: 43) *Nomenclator zoologicus* entry.
- †*Mardoellia* Blanchard, 1893: 130. Type species *Mardoell erdmanni* Danielssen, 1890, by monotypy; gender feminine. Blanchard (1893: 130) incorrectly emended the genus-group name *Mardoell* Danielssen, 1890, to *Mardoellia* (see Low and Reimer 2012a: 84). *Mardoellia* Blanchard, 1893, is therefore a junior objective synonym of *Mardoell* Danielssen, 1890, and a junior subjective synonym of *Epizoanthus* Gray, 1867. See also *Mardoell* Danielssen, 1890.
- **Mesozoanthus* Sinniger and Häussermann, 2009: 31, 32. Type species *Mesozoanthus fossii* Sinniger and Häussermann, 2009, by original designation and monotypy; gender masculine.
- Meto* Gistel, 1848: 181. Replacement name for *Hughea* Lamouroux, 1821. Not a zoantharian (see remarks under *Hughea* Lamouroux, 1821).
- **Microzoanthus* Fujii and Reimer, 2011: 421. Type species *Microzoanthus occultus* Fujii and Reimer, 2011, by original designation; gender masculine.
- †*Monanthus* Andres, 1883a: 538, 540, 541, 543. Type species *Isaurus tuberculatus* Gray, 1828, herein designated (see Appendix 3); gender masculine. First proposed as a subgenus of *Zoanthus* Lamarck, 1801. The type species is also the type species of *Isaurus* Gray, 1828, making *Zoanthus (Monanthus)* Andres, 1883, a junior objective synonym of *Isaurus* Gray, 1828 (see Appendix 3).
- †*Monothoa* Andres, 1883a: 521. Type species *Panceria spongiosa* Andres, 1877, herein designated (see Appendix 3); gender feminine. The type species is now a junior subjective synonym of *Isaurus tuberculatus* Gray, 1828, (see Appendix 3), making *Palythoa (Monothoa)* Andres, 1883, a junior subjective synonym of *Isaurus* Gray, 1828.

Montlibaldia [sic]. Incorrect subsequent spelling of *Montlivaltia* Lamouroux, 1821, by Ehrenberg (1834a: 271).

Montlivaltia Lamouroux, 1821: 78. Type species *Montlivaltia caryophyllata* Lamouroux, 1821, by monotypy; gender feminine. Lamouroux (1821: 78) and Audouin (1826: 229) discussed the similarities of *Montlivaltia* Lamouroux, 1821, and *Palythoa* Lamouroux, 1816. *Montlivaltia* Lamouroux, 1821, is now considered to be a species of Scleractinia (see Stolarski and Roniewicz 2001: 1097).

**Nanozoanthus* Fujii and Reimer, 2013: 512. Type species *Nanozoanthus harenaceus* Fujii and Reimer, 2013, by original designation; gender masculine.

**Neozoanthus* Herberts, 1972: 137. Type species *Neozoanthus tulearensis* Herberts, 1972, by original designation and monotypy; gender masculine.

Orinia Duchassaing de Fonbressin and Michelotti, 1860: 54. Type species *Orinia torpida* Duchassaing de Fonbressin and Michelotti, 1860, by monotypy; gender feminine. The type species is considered to be a junior subjective synonym of *Actinia osculifera* Le Sueur, 1817 (see Cha 2007: 40; Fautin 2016: 38). As the type species of *Orinia* Duchassaing de Fonbressin and Michelotti, 1860, is currently a junior subjective synonym of a species assigned to *Rhodactis* Milne Edwards and Haime, 1851 (see Fautin 2016: 38), the former becomes a junior subjective synonym of the latter.

**Palaeozoanthus* Carlgren, 1924: 470–473. Type species *Palaeozoanthus reticulatus* Carlgren, 1924, by original designation and monotypy; gender masculine.

†*Pales* Gray, 1867: 234, 235. Type species *Pales cliftoni* Gray, 1867, by monotypy; gender masculine. A subjective junior synonym of *Isaurus* Gray, 1828 (see Muirhead and Ryland, 1985: 325). The genus-group name *Pales* Gray, 1867, is also a junior homonym of *Pales* Meigen, 1800 (Diptera), *Pales* Robineau-Desvoidy, 1830 (Diptera), and *Pales* Koch, 1850 (Arachnida).

**Palythoa* Lamouroux, 1816: 359. Type species *Palythoe* [sic] *stellata* Lamouroux, 1816 [= *Alcyonium mammillosum* Ellis, in Ellis and Solander, 1786], subsequent designation by Haddon and Shackleton (1891b: 691); gender feminine. See Appendix 3 for a discussion on the type species and its designation. Frequently incorrectly spelt as “Palythoe” (see Low and Reimer 2011b: 63). See also the synonyms *Cavolinia* Schweigger, 1819, *Corticifera* Le Sueur, 1817, *Cynicus* Gistel, 1848, *Palythoa* (*Corticithoa*) Andres, 1883, *Palythoa* (*Gemmithoa*) Andres, 1883, and the incorrect spellings *Palythoe* and *Polythoa*.

†*Palythoa* (*Corticithoa*) Andres, 1883. See *Corticithoa* Andres, 1883.

†*Palythoa* (*Endeithoa*) Andres, 1883. See *Endeithoa* Andres, 1883.

†*Palythoa* (*Gemmithoa*) Andres, 1883. See *Gemmithoa* Andres, 1883.

†*Palythoa* (*Mammithoa*) Andres, 1883. See *Mammithoa* Andres, 1883.

†*Palythoa* (*Mammothoa*) Andres, 1883. See *Mammothoa* Andres, 1883, and *Mammithoa* Andres, 1883.

†*Palythoa* (*Monothoa*) Andres, 1883. See *Monothoa* Andres, 1883.

†*Palythoa* (*Taeniothoa*) Andres, 1883. See *Taeniothoa* Andres, 1883.

- †*Palythoaster* Haeckel, 1875: 44, pl. 1, fig. 5. Type species *Palythoa savignyi* Audouin, 1826, by monotypy; gender masculine. The type species *Palythoa savignyi* Audouin, 1826, is currently assigned to the genus *Palythoa* Lamouroux, 1816, making *Palythoaster* Haeckel, 1875, a junior subjective synonym of *Palythoa* Lamouroux, 1816 (see Appendix 3).
- †*Palythoe* [sic]. An incorrect spelling of *Palythoa* Lamouroux, 1816 (see Low and Reimer 2011b: 63). See also *Palythoe* Lamouroux, 1812.
- Palythoe* Lamouroux, 1812: 188. Type species *Gorgonia muricata* Pallas, 1766, by subsequent designation by Low and Reimer (2011b: 64); gender not determined. Now a subjective synonym of *Muricea* Lamouroux, 1821 [Octocorallia] (Low and Reimer 2011b: 64). Not a zoantharian (Low and Reimer 2011b: 63). *Palythoe* is also sometimes used as an incorrect spelling of *Palythoa* Lamouroux, 1816 (see Low and Reimer 2011b: 63).
- †*Playthoa* [sic]. Incorrect spelling of *Palythoa* Lamouroux, 1816, by various authors (e.g., Milliman et al. 1974: 162).
- †*Panceria* Andres, 1877: 221–226. Type species *Panceria spongiosa* Andres, 1877, by monotypy; gender feminine. The type species is a subjective synonym of *Isaurus tuberculatus* Gray, 1828 (see Appendix 3), making *Panceria* Andres, 1877, a junior subjective synonym of *Isaurus* Gray, 1828.
- †*Parapalythoa* Verrill, 1900: 560. Type species *Parapalythoa heilprini* Verrill, 1900, by monotypy. Herein considered to be a junior subjective synonym of *Palythoa* Lamouroux, 1816 (see Appendix 3). See also *Protopalythoa* Verrill, 1900.
- ****Parazoanthus*** Haddon and Shackleton, 1891a: 653, 654. Type species *Palythoa axinellae* O. Schmidt, 1862, by original designation; gender masculine. See also the synonyms *Bergia* Duchassaing de Fonbressin and Michelotti, 1860, *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860, *Heterozoanthus* Verrill, 1870, *Parapalythoa* Verrill, 1900, and *Protopalythoa* Verrill, 1900.
- Peachia* Gosse, 1855: 270, 271. Type species *Peachia hastata* Gosse, 1855, by subsequent designation by Carlgren (1949: 32); gender feminine. Not a zoantharian (Fautin et al. 2007: 220, 221; Fautin 2016: 47, 117).
- Platyzoanthus* Saville-Kent, 1893: 155. Type species *Platyzoanthus mussoides* Saville-Kent, 1893, by monotypy; gender masculine. Not a zoantharian (see Haddon 1898: 409, den Hartog 1980: 37, 39; Fautin 2016, 25, 28).
- †*Polyphoa* [sic]. Incorrect spelling of *Palythoa* Lamouroux, 1816, by Danielssen (1890: 136). The Danish version of the same text on the same page is spelt “*Polythoa*” (see *Polythoa*).
- †*Polythea* [sic]. Incorrect spelling of *Palythoa* Lamouroux, 1816, by Gistel (1848: 181).
- †*Polythoa* [sic]. Incorrect spelling of *Palythoa* Lamouroux, 1816, by Schweigger (1819: 100). Also throughout Andres (1883a, b, 1884).
- †*Protopalythoa* Verrill, 1900: 562. Replacement name for *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860, preoccupied by *Gemmaria* McCrady, 1859 (Hydrozoa). Type species *Gemmaria rusei* Duchassaing de Fonbressin and Michelotti, 1860, by typification of the preoccupied name; gender feminine. Verrill’s (1900: 562) original designation of *Gemmaria variabilis* Duerden, 1898, as the type spe-

- cies of this genus-group name is invalid (see Appendix 3). See also *Gemmaria* Duchassaing de Fombressin and Michelotti, 1860.
- †*Rhyzanthus* Andres, 1883a: 538, 541–544. Type species *Actinia sociata* Ellis, 1768, herein designated (see Appendix 3); gender masculine. First proposed as a subgenus of *Zoanthus* Lamarck, 1801. The type species is also the type species of the genus *Zoanthus* Lamarck, 1801, making *Zoanthus* (*Rhyzanthus*) Andres, 1883, a junior objective synonym of *Zoanthus* Lamarck, 1801.
- †*Savaglia* Nardo, 1877: 674. Unjustified emendation of *Savalia* Nardo, 1844 (see Appendix 3). See *Savalia* Nardo, 1844.
- **Savalia* Nardo, 1844: 433, 434. Type species *Gorgonia savaglia* Bertolini, 1819, by monotypy; gender feminine. *Savalia savaglia* (Bertolini, 1819), and *Gerardia lamarcki* (Haime, 1849), are currently considered subjective synonyms (see Appendix 3). As each species is also the type species of its respective genus, the genera *Savalia* Nardo, 1844, and *Gerardia* Lacaze-Duthiers, 1864, are also subjective synonyms. As discussed in Poche (1914), the valid name for this taxon is *Savalia savaglia* (Bertolini, 1819), which agrees with the Principle of Priority (Article 23, ICZN 1999: 24) (see discussion in Appendix 3). The name *Savaglia* is an unjustified emendation of *Savalia* Nardo, 1844, by Nardo (1877).
- Scolanthus* Gosse, 1853: 157. Type species *Scolanthus callimorphus* Gosse, 1853, by monotypy. Not a zoantharian (see Manuel 1981: 266; Fautin 2016: 46, 129).
- †*Sidisia* Gray, 1858: 489. Type species *Sidisia barleei* Gray, 1859, by subsequent monotypy; gender feminine. *Sidisia* Gray, 1858, was first proposed without the inclusion of any nominal species. In a subsequent paper, Gray (1859: 532) included *Sidisia barleei* Gray, 1859, as the only species in *Sidisia* Gray, 1858, and *Sidisia barleei* Gray, 1859, becomes the type species of *Sidisia* Gray, 1858, by subsequent monotypy (Article 69.3, ICZN, 1999: 73). Subjective synonym of *Epizoanthus* Gray, 1867. *Sidisia* Gray, 1858, was suppressed in favour of *Epizoanthus* Gray, 1867 (see Opinion 1689, ICZN 1992: 236).
- Solanthus* [sic]. Incorrect spelling of *Scolanthus* Gosse, 1853, by Gray (1867: 240).
- †*Spenopus* [sic]. Incorrect spelling of *Sphenopus* Steenstrup, 1856, by Herberts (1972: 72, 80, 142).
- †*Sphaenopus* [sic]. Incorrect spelling of *Sphenopus* Steenstrup, 1856, by Long, Poiner and Wassenberg (1995: 134).
- **Sphenopus* Steenstrup, 1856: 37. Type species *Sabella marsupialis* Gmelin, 1791, by original designation; gender masculine.
- †*Stephanidium* Hertwig, 1888: 52. Type species *Stephanidium schulzii* Hertwig, 1888, by monotypy; gender neuter. From the original description, the type species is possibly a species of zoantharian, but the type material will need to be located and examined (also see Delage and Hérouard 1901: 662). Herein, we make no decision as to the validity of this genus and species in the event that the identification of this genus and species, but note that this genus-group name is preoccupied by *Stephanidium* Ehrenberg, 1839 (*incerta sedis*).

- †*Taeniothoa* Andres, 1883a: 521, 532. Type species *Zoanthus sulcatus* Gosse, 1859, by subsequent designation by Williams (2000: 193); gender feminine. The type species is now assigned to the genus *Isozoanthus* Carlgren, in Chun, 1903 (see Williams 2000: 195), making *Palythoa* (*Taeniothoa*) Andres, 1883, a subjective synonym of *Isozoanthus* Carlgren, in Chun, 1903. Williams (2000: 195) enacted Article 23.9 (ICZN 1999: 27, 28) to reverse precedence between these genus-group names, thereby making *Palythoa* (*Taeniothoa*) Andres, 1883, a *nomen oblitum*, and *Isozoanthus* Carlgren, in Chun, 1903, a *nomen protectum*.
- **Terrazoanthus* Reimer and Fujii, 2010: 20. Type species *Terrazoanthus onoi* Reimer and Fujii, 2010, by original designation; gender masculine.
- **Thoracactis* Gravier, 1918: 12. Type species *Thoracactis topsenti* Gravier, 1918, by monotypy; gender masculine. Gravier (1918: 12, footnote) stated that the etymology of this name was from the Greek word “ $\theta\omicron\omicron\alpha\zeta, \alpha\chi\omicron\varsigma$ ” (= breastplate). It is also clear that Gravier (1918: 12) was using this word in combination with the ending “-actis” as commonly used for many other genera of Anthozoa, sea anemones in particular (e.g., *Amphiactis*, *Calliactis*, *Epiactis*, *Monactis*, *Paractis*; see Fautin 2016 for additional examples). In this context, the genus-group names ending with this suffix are to be treated in the same way as *Zoantha* and *Zoanthus* (see Low and Reimer 2012a: 85) in which Article 30.1.3 of the Code (ICZN 1999: 35) applies. The emendation of this name to *Thoracactus* by Walsh (1967: 49) is thus not justified. Also see the unjustified emendation *Thoracactis* by Walsh (1967: 49).
- †*Thoracactus* Walsh, 1967: 49. Unjustified emendation (and junior objective synonym) of *Thoracactis* Gravier, 1918.
- †*Toracactis* [sic]. Incorrect spelling of *Thoracactis* Gravier, 1918, by Herberts (1972: 80).
- Triga* Gray, 1867: 239. Type species *Triga philippinensis* Gray, 1867, by monotypy; gender feminine. McMurrich (1889: 125) stated that “in all probability this is a *Gemmaria*”, considering it to be similar to *G. rusei* Duchassaing de Fonbressin and Michelotti, 1860, which is clearly a species of *Palythoa* Lamouroux, 1812. Heider (1899a: 283; 1899b: 133) cited McMurrich (1889) and agreed, placing this species in *Gemmaria*. Ryland and Lancaster (2003: 411) discussed that “[t]he solitary *Triga philippinensis* Gray, 1867, though tentatively referred to *Gemmaria* by McMurrich (1889), seems more likely to have been an actinian, but Gray’s [1867] two-line diagnosis is insufficient even to determine the order with any certainty”. The solitary and long polyps of this species would indicate a species belonging to the genus *Sphenopus* Steenstrup, 1856. The final identity of this genus-group name and its type species remain unresolved.
- **Umimayanthus* Montenegro, Sinniger and Reimer, 2015: 76. Type species *Umimayanthus chanpuru* Montenegro, Sinniger and Reimer, 2015, by original designation; gender masculine.
- †*Verrillia* Andres, 1883a: 520, 545. Type species *Epizoanthus crassus* Verrill, 1869, by monotypy; gender feminine. The type species is currently assigned to the ge-

nus *Epizoanthus* Gray, 1867, making *Verrillia* Andres, 1883, a junior subjective synonym of *Epizoanthus* Gray, 1867 (see Appendix 3). *Verrillia* Andres, 1883, is preoccupied by *Verrillia* Stearns, 1873 (Scleractinia), and the name *Lirrevia* was proposed by Delphy (1939: 270) was proposed as a replacement name. See also *Lirrevia* Delphy, 1939.

- ****Zibrowius*** Sinniger, Ocaña and Baco, 2013: 7. Type species *Zibrowius ammophilus* Sinniger, Ocaña and Baco, 2013, by original designation; gender masculine.
- †*Zoantha* Lamarck, 1801: 363. Type species *Actinia sociata* Ellis, 1768, by monotypy; gender feminine. This is an incorrectly Latinised spelling of a name derived from Greek and should be corrected to *Zoanthus* (see Article 30.1.3, ICZN 1999: 35). This name was first correctly emended by Tilesius (1809: 374, footnote) (see Low and Reimer 2012a: 85). See also *Zoanthus* Lamarck, 1801.
- †*Zoanthella* van Beneden, 1897: 196. No type species designated. This name was established for zoantharian larvae of and is no longer in use as genus-group name (see Ryland et al. 2000: 191). Nevertheless, this name remains nomenclaturally available despite still being used as name for larvae.
- †*Zoanthina* van Beneden, 1897: 200. No type species designated. This name was established for zoantharian larvae of and is no longer in use as genus-group name (see Ryland et al. 2000: 191). Nevertheless, this name remains nomenclaturally available despite still being used as name for larvae.
- ****Zoanthus*** Lamarck, 1801: 363. Type species *Actinia sociata* Ellis, 1768, by monotypy; gender masculine. The name *Zoanthus* is derived from the Greek words ζωο (zoo = animal) and ανθος (anthos = flower). According to Article 30.1.3 (ICZN 1999: 35), the Greek ending “-os” should be latinised to the Latin masculine *-us*. Cuvier (1800: tables 9, 10) first used the name “*Zoanthus*”, but without the inclusion of species or a description and is a *nomen nudum*. Lamarck (1801: 363) next used the name *Zoantha* with the inclusion of *Actinia sociata* Ellis, 1768, the type species by monotypy. It is possible that Lamarck (1801) assumed that the gender of the Greek word for flower was feminine. Tilesius (1809: 394, footnote) first emended Lamarck’s (1801) name to *Zoanthus*. This emendation is conventionally attributed to Cuvier (1816: 53) (see also Haddon and Shackleton 1891b, 676; Low and Reimer 2012a: 85). See also the synonyms *Actimastus* Rafinesque, 1818, *Actinorhyza* Blainville, 1830, *Anthozoon* Gistel, 1848, *Mammillifera* Le Sueur, 1817, *Palythoa* (*Mammithoa*) Andres, 1883, *Zoanthus* (*Rhyzanthus*) Andres, 1883, and incorrect original spelling *Zoantha* Lamarck, 1801.
- †*Zoanthus* (*Corticanthus*) Andres, 1883. See *Corticanthus* Andres, 1883.
- †*Zoanthus* (*Monanthus*) Andres, 1883. See *Monanthus* Andres, 1883.
- †*Zoanthus* (*Rhyzanthus*) Andres, 1883. See *Rhyzanthus* Andres, 1883.
- †*Zooanthus* [*sic*]. Incorrect spelling of *Zoanthus* Lamarck, 1801 (e.g. Carlos et al. 1999: 1057, Kenny 2008: 78, Untawale and Dhargalkar 2002: 113).

Appendix 3.

Taxonomic and nomenclatural remarks on some previous unidentified or problematic family- and genus-group names in the Zoantharia Rafinesque, 1815

In this section, previously unidentified (or problematic) genus-group names have been grouped according to the senior synonym that they have been identified with. Two problematic groups of genus-group names are discussed last: 1) *Gemmaria* Duchassaing de Fombressin and Michelotti, 1860, *Parapalythoa* Verrill, 1900, *Protopalythoa* Verrill, 1900, and *Haplotella* Stechow, 1919; and 2) *Savalia* Nardo, 1844, *Savaglia* Nardo, 1877, and *Gerardia* Lacaze-Duthiers, 1864.

Synonyms of *Epizoanthus* Gray, 1867 (Epizoanthidae) (I): *Corticanthus* Andres, 1883, and *Endeithoa* Andres, 1883

The genus-group name *Corticanthus* was proposed as a subgenus of *Zoanthus* Lamarck, 1801 by Andres (1883a: 538, 541, 544; 1883b: 330, 333, 336; 1884: 323, 326, 327, 329) with the inclusion of *Epizoanthus paguriphilus* Verrill, 1882, *Mammillifera anduzii* Duchassaing de Fombressin and Michelotti, 1860, *Mammillifera conferta* Verrill, 1869, and *Mammillifera nitida* Verrill, 1869. *Epizoanthus paguriphilus*, is herein designated as the type species of *Zoanthus (Corticanthus)* Andres, 1883, making this genus-group name a subjective junior synonym of *Epizoanthus* Gray, 1867, as the type species is now assigned to the genus *Epizoanthus* (Pax and Müller, 1956: 12, 13; Reimer et al. 2010b: 730, 733, Walsh 1967: 45, 46).

The genus-group name *Endeithoa* was proposed as a subgenus of *Palythoa* by Andres (1883a: 521, 531; 1883b: 313, 323; 1884: 307, 316, 317) with the inclusion of *Zoanthus norvegicus* Koren and Danielssen, 1877, and *Zoanthus rubricornis* Holdsworth, 1861. *Zoanthus norvegicus* is herein designated as the type species of *Polythoa (Endeithoa)* making this genus-group name a subjective junior synonym of *Epizoanthus*, as the type species is now assigned to the genus *Epizoanthus* Gray, 1867 (Carreiro-Silva et al. 2011: 408, 413, Walsh 1967: 45).

Synonyms of *Epizoanthus* Gray, 1867 (Epizoanthidae) (II): *Lirrevia* Delphy, 1939, and *Verrillia* Andres, 1883

The genus-group name *Verrillia* was proposed by Andres (1883a: 520, 545; 1883b: 312, 337; 1884: 306, 330, 331) with the inclusion of only *Epizoanthus crassus* Verrill, 1869, the type species by monotypy (Article 68.3 of the Code; ICZN 1999: 71). *Epizoanthus crassus* Verrill, 1869, is currently assigned to the genus *Epizoanthus* Gray, 1867 (Lwowsky 1913: 560, Walsh, 1967: 39), and *Verrillia* is thus a junior subjective

synonym of *Epizoanthus* Gray, 1867. *Verrillia* Andres, 1883, is not a valid name as it is a junior homonym of *Verrillia* Stearns, 1873, which was proposed for a genus of scleractinian coral. Delphy (1939: 269, 270) proposed the replacement name *Lirrevia*, which is an objective synonym of *Verrillia* Andres, 1883, and a junior subjective synonym of *Epizoanthus* Gray, 1867.

Synonyms of *Isaurus* Gray, 1828 (Zoanthidae): *Monanthus* Andres, 1883, *Monothoa* Andres, 1883, and *Panceria* Andres, 1877

Muirhead and Ryland (1985: 325) listed *Polythoa* (*Monothoa*) Andres, 1883, *Zoanthus* (*Monanthus*) Andres, 1883, and *Panceria* Andres, 1877, as synonyms of *Isaurus* Gray, 1828, but did not give details for this synonymy. Herein, type species are designated for *Polythoa* (*Monothoa*) Andres, 1883, and *Zoanthus* (*Monanthus*) Andres, 1883, to formally fix the identity of these genus-group names. The nomenclatural consequences of our type species designations are in agreement with the taxonomic conclusions of Muirhead and Ryland (1985).

The genus-group name *Monothoa* was proposed as a subgenus of *Palythoa* Lamouroux, 1816, by Andres (1883a: 521, 530, 537; 1883b: 313, 322, 329; 1884: 307, 315, 316, 322) with the inclusion of *Hughaea caraibeorum* Duchassaing de Fombressin, 1850, *Mamillifera fulva* Quoy and Gaimard, 1834, *Mamillifera vanikorensis* Quoy and Gaimard, 1834, *Mamillifera viridifusca* Quoy and Gaimard, 1834, *Panceria spongiosa* Andres, 1877, and *Triga philippinensis* Gray, 1867. *Panceria spongiosa* Andres, 1877, is herein designated as the type species of *Polythoa* (*Monothoa*) Andres, 1883. *Panceria spongiosa* Andres, 1877, is currently considered to be a junior subjective synonym of *Isaurus tuberculatus* Gray, 1828 (Muirhead and Ryland, 1985: 326), and *Polythoa* (*Monothoa*) Andres, 1883, becomes a junior subjective synonym of *Isaurus* Gray, 1828.

The genus-group name *Monanthus* was proposed as a subgenus of *Zoanthus* Lamarck, 1801, by Andres (1883a: 538, 540, 541, 543; 1883b: 330, 332, 333, 335; 1884: 323, 325, 326, 328, 329) with the inclusion of *Isaurus tuberculatus* Gray, 1828, *Isaura neglecta* Duchassaing de Fombressin and Michelotti, 1860, *Pales cliftoni* Gray, 1867, and *Palythoa savignyi* Audouin, 1826. *Isaurus tuberculatus* Gray, 1828, is herein designated as the type species of *Zoanthus* (*Monanthus*) Andres, 1883, thereby making *Zoanthus* (*Monanthus*) Andres, 1883, a junior objective synonym of *Isaurus* Gray, 1828, as they have the same type species (Article 61.3.3 of the Code, ICZN 1999: 64) (also see Low and Reimer 2012b: 47).

The genus-group name *Panceria* was proposed by Andres (1877: 221, 226) with the inclusion of *Panceria spongiosa* Andres, 1877, the type species by monotypy (Article 68.3 of the Code, ICZN 1999: 71). As discussed above, *Panceria spongiosa* Andres, 1877, is a junior subjective synonym of *Isaurus tuberculatus* Gray, 1828 and *Panceria* Andres, 1877, becomes a junior subjective synonym of *Isaurus* Gray, 1828.

Synonyms of *Palythoa* Lamouroux, 1816 (Sphenopidae) (I): *Cavolinia* Schweigger, 1819, and *Cynicus* Gistel, 1848

Schweigger (1819: 99, 100) proposed the genus-group name *Cavolinia* for two species of zoanths: *Alcyonium mammosum* Ellis, in Ellis and Solander, 1786 (incorrectly spelt as “*mamillosum*”), and *Cavolinia rosea* Schweigger, 1819 (an unnecessary replacement name for *Madrepora denudata* Cavolini, 1785). *Alcyonium mammosum* Ellis, in Ellis and Solander, 1786, and *Madrepora denudata* Cavolini, 1785 (and therefore *Cavolinia rosea* Schweigger, 1819), are now assigned to the genus *Palythoa* Lamouroux, 1816 (see Milne Edwards 1857: 301, Ryland and Lancaster 2003: 410), making *Cavolinia* Schweigger, 1819, a junior subjective synonym of *Palythoa* Lamouroux, 1816.

No type species has been designated for *Cavolinia* Schweigger, 1819, and no designation is necessary as *Cavolinia* Schweigger, 1819, is an invalid junior homonym of *Cavolinia* Abildgaard, 1791 (Mollusca), and *Cavolinia* Bruguière, 1791 (Mollusca), and has been placed on the *Official Index of Rejected and Invalid Generic Names in Zoology* (see ICZN 1969: 28).

The replacement name *Cynicus* was proposed by Gistel (1848: viii) for *Cavolinia* Schweigger, 1819. *Cynicus* Gistel, 1848, is therefore an unnecessary replacement name for, and junior objective synonym of, *Cavolinia* Schweigger, 1819.

Synonyms of *Palythoa* Lamouroux, 1816 (Sphenopidae) (II): *Corticifera* Le Sueur, 1817, *Corticithoa* Andres, 1883, and *Gemmithoa* Andres, 1883

The genus-group name *Corticifera* was proposed by Le Sueur (1817: 178, 179) with the inclusion of *Corticifera flava* Le Sueur, 1817, and *Corticifera glareola* Le Sueur, 1817. Haddon and Shackleton (1891b: 692) designated *Corticifera glareola* Le Sueur, 1817, as the type species of *Corticifera* Le Sueur, 1817. As the type species of *Corticifera*, is currently assigned to genus *Palythoa*, *Corticifera*, is now a junior subjective synonym of *Palythoa* (see also Haddon and Shackleton 1891b: 692).

The genus-group name *Corticithoa* was proposed as a subgenus of *Palythoa* by Andres (1883a: 521, 535–538; 1883b: 313, 327–330; 1884: 307, 320–323) with the inclusion of *Alyconium tuberculosum* Esper, 1805, *Corticifera aggregata* Lesson, 1830, *Corticifera glareola* Le Sueur, 1817, *Gemmaria humilis* Verrill, 1869, *Mamillifera clavata* Duchassaing de Fombressin, 1850, *Mamillifera lutea* Quoy and Gaimard, 1834, and *Palythoa cinerea* Duchassaing de Fombressin and Michelotti, 1864. *Alyconium tuberculosum* Esper, 1805, is herein designated as the type species of *Palythoa* (*Corticithoa*) Andres, 1883. *Alyconium tuberculosum*, is currently assigned to the genus *Palythoa* (see Reimer et al. 2006: 92, Ryland and Lancaster 2003: 409, 410, Walsh 1967: 19, 20). By this type species designation, *Palythoa* (*Corticithoa*) Andres, 1883, becomes a junior subjective synonym of *Palythoa*.

The genus-group name *Gemmithoa* was proposed as a subgenus of *Palythoa*, by Andres (1883a: 521, 532, 533; 1883b: 313, 324, 325; 1884: 307, 318) with the in-

clusion of *Mammillifera brevis* Duchassaing de Fombressin, 1850, the type species by monotypy (Article 68.3 of the Code; ICZN 1999: 71). *Mammillifera brevis* Duchassaing de Fombressin, 1850, is now assigned to the genus *Palythoa* (see Walsh 1967: 5), making *Polythoa* (*Gemmithoa*) Andres, 1883, a junior subjective synonym of *Palythoa* Lamouroux, 1816.

Synonyms of *Palythoa* Lamouroux, 1816 (Sphenopidae) (III): *Palythoaster* Haeckel, 1875

Haeckel (1875: 44, pl. 1, fig. 5) proposed the name *Palythoaster* for a single species, *Palythoa savignyi* Audouin, 1826, the type species by monotypy (Article 68.3 of the Code, ICZN 1999: 71). Although the date on the title-page of the work by Haeckel is “1876”, it was available in 1875 (see Leuckart 1875: 463).

Ehrenberg (1834a: 269; 1834b: 45) considered *Palythoa savignyi* Audouin, 1826, to be a valid species of zoantharian, but placed it in the genus *Hughea* Lamouroux, 1821. The genus *Hughea* Lamouroux, 1821, has long been confused for a genus of a zoantharian, which it is not (see Ryland and Lancaster 2003: 411; Appendix 2).

As discussed by Low and Reimer (2012b), the name *Palythoa savignyi* was proposed by Audouin (1826: 229) for an unnamed figured by Savigny (1811: pl. 2, fig. 1) from Egypt. Based on the figure and the other information provided by Savigny (1811) and Audouin (1826), we herein agree with the opinion of Ehrenberg (1834a: 269; 1834b: 45) in considering *Palythoa savignyi* Audouin, 1826, to be a valid species of zoantharian, but unlike this previous author, we consider it to be a species of *Palythoa* Lamouroux, 1816.

As the type species of the genus *Palythoaster* Haeckel, 1875, is considered to be a species of *Palythoa* Lamouroux, 1816, the former becomes a junior subjective synonym of the latter.

Synonyms of *Zoanthus* Lamarck, 1801 (Zoanthidae) (I): *Actimastus* Rafinesque, 1818, and *Mammillifera* Le Sueur, 1817

The genus-group name *Mammillifera* was proposed by Le Sueur (1817: 177, 178) with the inclusion of *Mammillifera auricula* Le Sueur, 1817, and *Mammillifera nymphaea* Le Sueur, 1817. *Mammillifera auricula* was designated as the type species of *Mammillifera* (see Haddon and Shackleton 1891a: 626). As the type species *Mammillifera auricula* is currently assigned to the genus *Zoanthus* Lamarck, 1801 (Duerden 1898: 334, Walsh 1967: 22, Reimer, et al. 2012a: 7), *Mammillifera* Le Sueur, 1817, and *Zoanthus* Lamarck, 1801, are subjective synonyms.

Rafinesque (1818: 271) proposed the name *Actimastus* to replace *Mammillifera* Le Sueur, 1817, stating that “*Mammillifera* of Lesueur, is rather too long; it is too much like *Mammillaria* in meaning and sense, and is composed of two Latin names united,

which are tolerated in the specific nomenclature, but not often in the generic; lastly it has too much likeness to the classical name of Mammalia to be tolerated. It must then be changed into *Actimastus*; meaning radiated mammilla”. As it is an unjustified emendation of *Mammillifera* Le Sueur, 1817 (Article 19.1 of the Code, ICZN 1999: 21), *Actimastus* Rafinesque, 1818, becomes a junior objective synonym of *Mammillifera* Le Sueur, 1817.

Synonyms of *Zoanthus* Lamarck, 1801 (Zoanthidae) (II): *Actinorbiza* Agassiz, 1846, and *Actinorhyza* Blainville, 1830

In a discussion on the genus *Zoanthus* Lamarck, 1801, Blainville (1830: 295) proposed the replacement name *Actinorhyza* for this genus-group stating only “[c]e genre, dans notre Système de nomenclature, pourroit être nommé Actinorhyse, *Actinorhyza*”. We herein consider the name *Actinorhyza* Blainville, 1830, to be an unnecessary replacement name for, and junior objective synonym of, *Zoanthus* Lamarck, 1801. The type species is thus identical to that of *Zoanthus* Lamarck, 1801 (viz., *Actinia sociata* Ellis, 1768) (Article 67.8 of the Code, ICZN 1999: 68). In a later publication, Blainville (1834: 329) spelt the name as *Actinorhyza*, an incorrect subsequent spelling. Another incorrect subsequent spelling is *Actinorrhiza*, as used by Ehrenberg (1834a: 269).

Agassiz (1846: 7) proposed the unjustified emendation *Actinorbiza* for *Actinorhyza* Blainville, 1830, and the former is a junior objective synonym of the latter (Article 19.1 of the Code, ICZN, 1999: 21).

Synonyms of *Zoanthus* Lamarck, 1801 (Zoanthidae) (III): *Mammithoa* Andres, 1883, “*Mammothoa*” Andres, 1883, and *Rhyzanthus* Andres, 1883

The genus-group name *Mammithoa* was proposed as a subgenus of *Palythoa* Lamouroux, 1816, by Andres (1883a: 521, 533–535; 1883b: 313, 325–327; 1884: 307, 318–320). In the preliminary introduction to the newly proposed genus-group names, Andres (1883a: 521; 1883b: 313; 1884: 307) listed the name as *Polythoa* (*Mammithoa*) and included six species under this grouping: *Mammillifera auricula* Lesueur, 1817, *Mammillifera cingulata* Quoy and Gaimard, 1834, *Mammillifera nymphaea* Le Sueur, 1817, *Mammillifera univittata* Lorenz, 1860, *Mammillifera viridis* Quoy and Gaimard, 1834.

In a more detailed listing of the species included in the genus-group, the name was spelt as “*Mammothoa*” (Andres 1883a: 533–535; 1883b: 325–327; 1884: 318–320). As the genus-group names *Mammithoa* and *Mammothoa* were both used in the same publication (Andres 1883a: 521, 533–535), they were made available simultaneously and are both available names. First reviser action is herein taken to select *Mammithoa* Andres, 1883, as the correct original spelling (Article 24.2.3 of the Code, ICZN 1999: 30).

In this detailed listing of the genus-group name *Mammithoa* (incorrectly spelt as *Mammothoa*), Andres (1883a: 534, 535) discussed the included species as discussed above but with the name “*Polythoa* (*Mammithoa*) *nymphosa* Dana”, being used in place

of *Mammillifera nymphaea* Le Sueur, 1817. From the synonymy of “*Polythoa* (*Mammithoa*) *nymphosa* Dana”, it is clear that the species-group name “*nymphosa* Dana” was a replacement for *nymphaea* Le Sueur, 1817. Although the species-group name *nymphosa* was attributed to Dana (see also Walsh 1967: 28), this species-group name was never used by Dana, and the authorship of *Polythoa* (*Mammithoa*) *nymphosa* should be attributed to Andres (1883a: 534, 535), and the name is herein considered an unnecessary replacement name for (and objective synonym of) *Mammillifera nymphaea* Le Sueur, 1817.

Mammillifera nymphaea Le Sueur, 1817, is herein designated as the type species of *Polythoa* (*Mammithoa*) Andres, 1883. *Mammillifera nymphaea* Le Sueur, 1817, is currently considered to be a subjective synonym of *Zoanthus pulchellus* Duchassaing de Fonbressin and Michelotti, 1864 (Acosta et al. 2005: 160, Duerden, 1898: 334, Reimer et al. 2012a: 7). *Zoanthus pulchellus* Duchassaing de Fonbressin and Michelotti, 1864, is currently assigned to the genus *Zoanthus* Lamarck, 1801 (Acosta et al. 2005: 160, Reimer et al. 2012a: 7, Walsh 1967: 29) and *Polythoa* (*Mammithoa*) Andres, 1883, therefore becomes a junior subjective synonym of *Zoanthus* Lamarck, 1801.

The genus-group name *Rhyzanthus* was proposed as a subgenus of *Zoanthus* Lamarck, 1801, by Andres (1883a: 538, 541–544; 1883b: 330, 334–336; 1884: 306, 330, 331) with the inclusion of *Actinia sociata* Ellis, 1768, *Zoanthus alderi* Gosse, 1859, *Zoanthus dubia* Lesueur, 1817, *Zoanthus mertensii* Brandt, 1835, and *Zoanthus solandri* Lesueur, 1817. *Actinia sociata* Ellis, 1768, is herein designated as the type species of *Zoanthus* (*Rhyzanthus*) Andres, 1883, thereby making this genus-group name an objective junior synonym of *Zoanthus* Lamarck, 1801, as they have the same type species (Article 61.3.3 of the Code, ICZN 1999: 64).

***Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860, and its replacement names *Protopalythoa* Verrill, 1900, and *Haplotella* Stechow, 1919, junior subjective synonyms of *Palythoa* Lamouroux, 1816, as well as comments on *Parapalythoa* Verrill, 1900**

The genus-group name *Gemmaria* was first proposed by Duchassaing de Fonbressin and Michelotti (1860: 55) with the inclusion of four nominal species: *Gemmaria rusei* Duchassaing de Fonbressin and Michelotti, 1860, *Gemmaria swiftii* Duchassaing de Fonbressin and Michelotti, 1860, *Mammillifera brevis* Duchassaing de Fonbressin, 1850, and *Mammillifera clavata* Duchassaing de Fonbressin, 1850. No type species was designated.

The identity of *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860, was debated by early workers. Duerden (1898: 353; 1903: 501) considered *Gemmaria swiftii* Duchassaing de Fonbressin and Michelotti, 1860, to be a species of *Parazoanthus* Haddon and Shackleton, 1891, and transferred this species to the genus. The assignment of *Gemmaria swiftii* Duchassaing de Fonbressin and Michelotti, 1860, to the genus *Parazoanthus* Haddon and Shackleton, 1891, remains the prevailing opinion

(e.g. Ryland and Lancaster 2003: 410, Reimer et al. 2014a: 1, 3–7). *Mammillifera clavata* Duchassaing de Fonbressin, 1850, was considered by Reimer et al. (2012a: 8, 9) to be a valid species of *Palythoa* Lamouroux, 1816. The assignment of *Mammillifera brevis* Duchassaing de Fonbressin, 1850, to the genus *Epizoanthus* Gray, 1867, by workers such as Andres (1884: 311) was questioned by Haddon and Shackleton (1891b: 687), who nevertheless stated that “neither *G. swiftii* nor *G. brevis* would appear to belong to the same genus as the type species, nor is it certain that *G. clavata* does either”.

It was probably for these reasons that Haddon and Shackleton (1891a: 626) designated *Gemmaria rusei* Duchassaing de Fonbressin and Michelotti, 1860, as the type species of *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860. The spelling of the type species, *Gemmaria rusei* Duchassaing de Fonbressin and Michelotti, 1860, requires some discussion. The International Commission on Zoological Nomenclature (ICZN 1998: 121, 122) ruled that the spellings of two genus- and species-group names derived from the surname Riise but earlier incorrectly spelled as Rüsei by Duchassaing de Fonbressin and Michelotti (see Bayer and Grasshoff 1997: 11) should be corrected. However, *Gemmaria rusei* Duchassaing de Fonbressin and Michelotti, 1860, was not one of the names that was ruled on. The spelling of the type species of *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860, should thus remain as *Gemmaria rusei* Duchassaing de Fonbressin and Michelotti, 1860 (but see also Ryland and Lancaster 2003: 410).

The genus-group name *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860, is however not a valid name as it is a junior homonym of *Gemmaria* McCrady, 1859. The genus-group name *Gemmaria* was conditionally proposed by McCrady (1859: 151) for a new species of hydrozoan, *Zanclaea gemmosa*, described in the same paper. Although the genus-group name *Gemmaria* McCrady, 1859, is currently considered to be a junior synonym of *Zanclaea* Gegenbaur, 1856 (see Schuchert 2010: 487), it nevertheless remains an available name. *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860, therefore remains an invalid name as it is preoccupied by *Gemmaria* McCrady, 1859 (Article 53.2 of the Code, ICZN 1999: 57).

Realising this homonymy, Verrill (1900: 562) wrote: “*Protopalpythoa* nom. nov. Type *G. variabilis* Duerden. *Gemmaria* Duch. and Mich., Corall. Antill., p. 55, 1860, (non McCready [*sic*], 1859)”. As *Protopalpythoa* Verrill, 1900, was explicitly proposed as a replacement name for *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860 (preoccupied by *Gemmaria* McCrady, 1859), the designation of *Gemmaria variabilis* Duerden, 1898, as the type species of *Protopalpythoa* Verrill, 1900, is invalid.

Article 67.8 of the Code (ICZN 1999: 68) states that “[i]f an author publishes a new genus-group name expressly as a new replacement name (*nomen novum*) for a previously established name ... both the prior nominal taxon and its replacement have the same type species, and type fixation for either applies also to the other, despite any statement to the contrary”.

The type species of *Protopalpythoa* Verrill, 1900, is therefore *Gemmaria rusei* Duchassaing de Fonbressin and Michelotti, 1860, as this is the type species of *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860 (as designated by Haddon and

Shackleton 1891a: 626). *Protopalythoa* Verrill, 1900, is therefore an objective synonym of *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860.

In their important paper on *Protopalythoa* Verrill, 1900, Ryland and Lancaster (2003: 410, 411) followed Verrill's (1900: 562) incorrect type species designation of *Gemmaria variabilis* Duerden, 1898, as the type species of *Protopalythoa* Verrill, 1900. Ryland and Lancaster (2003: 410, 411) stated: "*Gemmaria* Duchassaing and Michelotti, 1860, was introduced for *G. Rusei* nov., *Mamillifera clavata* Duchassaing, 1850, *G. swifti* nov. (= *Parazoanthus swifti*), and *M. brevis* Duchassaing, 1850, with no designation of type species. *Protopalythoa* Verrill, 1900 was a *nomen novum* because *Gemmaria* was preoccupied by *Gemmaria* McCrady, 1857 (cited as McCready, 1859), a hydroid. Verrill designated *Gemmaria variabilis* Duerden, 1898, as type species, on the grounds that *G. riisei* (sic: *G. rusei*) was unrecognisable and Duchassaing and Michelotti's other species were not congeners". Ryland and Lancaster (2003: 410, 411) considered the characters exhibited by *Gemmaria variabilis* Duerden, 1898 (and other congeners) to be sufficiently different from *Alcyonium mammillosum* Ellis, in Ellis and Solander, 1786 (an objective synonym of the type species of *Palythoa* Lamouroux, 1816), to require a genus-group of their own, for which they used *Protopalythoa* Verrill, 1900.

But what is the real identity of *Protopalythoa* Verrill, 1900? The identity of the genus-group *Protopalythoa* Verrill, 1900, then rests on the identity of its type species *Gemmaria rusei* Duchassaing de Fonbressin and Michelotti, 1860. Volpi and Benvenuti (2003: 66) listed the existence of a syntype of *Gemmaria rusei* Duchassaing de Fonbressin and Michelotti, 1860, at the Museo di Storia Naturale Università di Firenze (accession number: MZUF 847). Through the assistance of C. Volpi and S. Bambi, we have been able to examine high-resolution photographs of the syntype. The images clearly show a species of *Palythoa* Lamouroux, 1816, based on sampling information (depth, locality, etc.) and heavy sand encrustation of the polyps, and therefore *Gemmaria rusei* Duchassaing de Fonbressin and Michelotti, 1860, is referable to the genus *Palythoa* Lamouroux, 1816, as is currently defined.

Although the specific identity of *Gemmaria rusei* Duchassaing de Fonbressin and Michelotti, 1860, will require further research, we herein assign this species-group taxon to *Palythoa* Lamouroux, 1816. *Protopalythoa* Verrill, 1900, thus becomes a junior subjective synonym of *Palythoa* Lamouroux, 1816. Likewise, the preoccupied genus-group name *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860, as well as *Parapalythoa* Verrill, 1900 (discussed below), and *Haplotella* Stechow, 1919 (also discussed below), become junior subjective synonyms of *Palythoa* Lamouroux, 1816.

Verrill (1900: 562, 1907: 287) himself had already observed the close affinity between *Protopalythoa* Verrill, 1900, and *Palythoa* Lamouroux, 1816, stating "[t]he name *Gemmaria* having been preoccupied in Hydrozoa, it is necessary to give a new one to this group, if it is to be considered as really distinct from *Palythoa*, from which it seems to differ only in the fact that the zoöids are not united together laterally by coenenchyma, but only by stolons or based expansions. Some species of *Palythoa* are not thus united for more than half their height, or even less, and perhaps future discoveries may show a complete gradation between the two conditions" (Verrill 1900: 562). Similarly,

Verrill (1907: 287) also discussed that: “[s]hould they [i.e., the species assigned to *Protopalythoa*] ultimately prove to be identical, it would probably be necessary to unite the genus *Protopalythoa* (= *Gemmaria* of many authors) to *Palythoa* ... The only tangible difference between the two genera is the presence in the latter of a thick crust-like coenenchyma, uniting the polyps together laterally. But in this species they are often united for less than half their height”.

Two other genus-group names need to be discussed. *Parapalythoa* Verrill, 1900, is another genus-group name that needs to be discussed in connection with *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860. Verrill (1900: 560) proposed the name *Parapalythoa* for specimens of “*Gemmaria Rusei*” described by McMurrich (1889: 124, 125) from Bermuda. Verrill (1900: 560) considered the material studied by McMurrich (1889) to be distinct from *Gemmaria rusei* Duchassaing de Fonbressin and Michelotti, 1860, and proposed that they should be called *Parapalythoa heilprini*. Although Verrill (1907: 283) later stated that the genus-group name *Parapalythoa* was an error for *Protopalythoa*, this statement has no bearing on the availability of the genus-group name *Parapalythoa*, and the name remains an available one, with *Parapalythoa heilprini* Verrill, 1900, being the type species by monotypy. Regardless of the validity of *Parapalythoa heilprini* Verrill, 1900, it is clear that Verrill (1900: 560) was intending to establish a new genus-group for a taxon similar to, but distinct from, *Gemmaria rusei* Duchassaing de Fonbressin and Michelotti, 1860. As discussed above, *Gemmaria rusei* Duchassaing de Fonbressin and Michelotti, 1860, is herein considered to be a species of *Palythoa* Lamouroux, 1816, and as the differences in the specimens of “*Gemmaria rusei*” described by McMurrich (1889: 124, 125) not being sufficient for a distinction at the genus-group level, we herein consider *Parapalythoa* Verrill, 1900, to also be a junior subjective synonym of *Palythoa* Lamouroux, 1816.

A final name that needs to be discussed in connection with *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860, is *Haplotella* Stechow, 1919. Stechow (1919: 853) proposed the name *Haplotella* as a replacement name for *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860, and has the same type species (see discussion above). *Haplotella* Stechow, 1919, is a junior objective synonym of *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860, and a junior subjective synonym of *Protopalythoa* Verrill, 1900, and *Parapalythoa* Verrill, 1900. The date of publication of *Haplotella* is conventionally cited as “1920”, but in an abstract to the work that appeared in 1919, the following was stated: “Die Aktiniengattung “*Gemmaria*” Duchassaing et Michelotti 1861 erhält wegen Präokkupation durch ein Hydroidengenus den Namen *Haplotella*” (Stechow, 1919: 853). The date of publication of the name *Haplotella* is thus 1919.

In conclusion, the genus-group names *Gemmaria* Duchassaing de Fonbressin and Michelotti, 1860, *Parapalythoa* Verrill, 1900, *Protopalythoa* Verrill, 1900, and *Haplotella* Stechow, 1919, are all herein considered to be synonyms of *Palythoa* Lamouroux, 1816. Furthermore, the genus-group name *Protopalythoa* Verrill, 1900, can no longer be used by authors who consider *Gemmaria variabilis* Duerden, 1898 (and other related species) to be sufficiently distinct to require a genus-group of its own (i.e. *Protopalythoa* sensu Ryland and Lancaster 2003). To resolve this, a new genus-group name will need to be established.

***Savalia* Nardo, 1844, is a senior subjective synonym of *Gerardia* Lacaze-Duthiers, 1864, with comments the family-group names Gerardiidae Verrill, 1865, Savagliidae Brook, 1889, and Parazoanthidae Delage and Hérouard, 1901**

The family-, genus- and species-group names for the zoantharian living on gorgonians in the Mediterranean area has been a matter of a long-running historical debate (reviewed in Brook 1889: 79, 80, Bell 1891: 89–91; Carlgren 1895: 319–334, Roche and Tixier-Durivault 1951: 402–409).

At the species-group rank, the first available name to be applied to this taxon was “*Gorgonia Savaglia*” by Bertoloni (1819: 219). The name “savaglia” being the commonly used vernacular name for this animal (e.g., Imperato 1599: 724; Garençieres 1676: 5, Bell 1899: 90). Haime (1849: 225, note) later established “*Leiopathes Lamarcki*” (the species-group name being spelled with only one “i”).

At the genus-group rank, Nardo (1844: 433, 434) and Lacaze-Duthiers (1864a: 87) respectively established the genus-group names *Savalia* (type species *Gorgonia savaglia* Bertoloni, 1819), and *Gerardia* (type species *Leiopathes lamarcki* Haime, 1849). Each species is the type species of its respective genus-group by monotypy (Article 68.3 of the Code, ICZN, 1999: 71). As a further complication, the genus-group name *Savaglia* was used by Nardo (1877), who was clearly intending to emend the spelling of *Savalia* Nardo, 1844.

Brook (1889: 79, 80) considered the valid binomial name for the species under consideration to be *Savaglia lamarcki* (Haime, 1849). The reason for for doing was given as: “Lacaze Duthiers was the first to show the true relations of this form, and its difference from the typical Antipathidae. Nardo, in 1843 [*sic*], gave the generic name *Savaglia* to the species described as *La Savaglia* by Donati in 1765, which he says is identical with *Leiopathes lamarcki*, Haime; in this case his name has priority over that of *Gerardia*, instituted by Lacaze Duthiers in 1864. This I gather from a more recent paper; I have not seen the original, and do not know if Nardo gave the species a specific, as well as a generic, name; there is no mention of one in his recent publication. I have, therefore, retained the specific name of Haime. Although it seems highly probable that Nardo’s *Savaglia* is the same as *Gerardia*, Lacaze Duthiers, his description of the polyp does not agree with Lacaze Duthiers’ observations on living specimens. Nardo states that the polyp has only fourteen tentacles, whereas the species in question has twenty-four” (Brook 1889: 79).

Bell (1891: 89–91) took the opposite position and considered the valid binomial name to be *Gerardia savalia* (Bertoloni, 1819). Bell (1891: 89–91) argued that the establishment of the genus-group name *Savalia* in Nardo (1844: 433, 434) was not ‘legitimate’ as “Nardo does not write out the full name or names of the species to be placed in this genus, but it is clear that had he done so he would have then written *Savaglia savaglia* (or *Savalia savalia*). This use of a specific for a generic name has been forbidden by the rules of the British Association. Meanwhile this species had become famous by the researches of Lacaze-Duthiers, who, conferring on it (in 1864) the generic name of *Gerardia*, retained for it the specific name of *lamarcki* given it by Haime (in 1849) when he called it *Leiopathes lamarcki*. ... It is clear that there is no escape

from the conclusion that the proper specific name of this Coral is *savalia*, and the generic *Gerardia* ...”.

In order to resolve the matter of the correct genus- and species- group names, a more detailed analysis of the description given by Nardo (1844: 433, 434). The description is given as follows: “*Sotto famiglia II.^a Savalini — Polipi a sedici tentacoli! Gen. Savalia N. ... Fa meraviglia come sia sfuggito all’occhio degli osservatori il bel lavoro del Donati V. sull’Antipate dell’ Adriatico Gorgonia savoglia [sic] (Bertoloni) inserito nel primo volume del Giornale di Grisellino, ove vedesi esattamente descritto e figurato l’animale con i suoi sedici tentacoli. Non v’ha dubbio che una tale specie debbasi distinguere dal genere Anthipathes. Costituisee anzi a mio credere una sotto famiglia, come mostrerò in più esteso lavoro relativo ai caratteri distintivi delle famiglie dei Zoofitarj*” (Nardo 1844: 433, 434).

This is translated as: “Subfamily 2. Savalini — Polyps with 16 tentacles. Genus *Savalia* N[obis]. ... The fine work of V. Donati on Antipate [= black corals/zoophytes] of the Adriatic Sea in the first volume of the *Giornale di Grisellino* has long been overlooked by previous workers, in which *Gorgonia savoglia [sic]* (Bertoloni) was exactly described and figured as an animal with sixteen tentacles. There is not doubt that this species is deserving of a genus distinct from the genus *Anthipathes*. Indeed, as I will show in a more extensive work relating to the distinctive characteristics of the families of Zoofitarj, it constitutes a new subfamily”.

The objection raised by Brook (1889: 79) for not using *Gorgonia savaglia* Bertoloni, 1819, as the species-group name is not tenable, as the only reason given is that he did “not know if Nardo gave the species a specific name” as Brook did not have access to the 1844 paper by Nardo. From the description by Nardo (1844: 433, 434), the genus-group name *Savalia* was clearly established for *Gorgonia savaglia* Bertoloni, 1819 (incorrectly spelled as “*savoglia*”). Virtually all recent authors consider *Gorgonia savaglia* Bertoloni, 1819, to be a subjective synonym of *Leiopathes lamarcki* Haime, 1849 (e.g. Roche and Tixier-Durivault 1951: 402, Ocaña et al. 1995: 155, Ocaña and Brito 2004: 170, Sinniger et al. 2005: 1124, 1125, Ocaña et al. 2007: 163–167). *Gorgonia savaglia* Bertoloni, 1819, has priority over *Leiopathes lamarcki* Haime, 1849, and the former is the valid species-group name.

The objections raised by Bell (1891: 90) for not accepting *Savalia* Nardo, 1844, are also not justified. Firstly, Nardo (1844: 433, 434) did indeed provide the name of the species he intended to be placed in *Savalia* Nardo, 1844 (as discussed above). Secondly, as already stated by Poche 1914: 104, tautomeric names (viz., “*Savaglia savaglia* / *Savalia savalia*”) are not invalid on this basis alone (Articles 18 and 23.3.7 of the Code, ICZN 1999: 21, 26). As their respective type species are considered to be subjective synonyms (see above), the genus-group name *Savalia* Nardo, 1844, is a subjective synonym of *Gerardia* Lacaze-Duthiers, 1864 (see also Sinniger et al. 2013: [1], [2], [7]). As *Savalia* Nardo, 1844, has priority over *Gerardia* Lacaze-Duthiers, 1864, the former is the valid genus-group name.

At the family-group rank, three names have been proposed for *Savalia savaglia* (Bertoloni, 1819). As noted above, Nardo (1844: 433, 434) established the family-group name Savaliidae (as “Savalini”) based on the genus-group name *Savalia* Nardo,

1844. Later, Verrill (1869 [in 1868–1870]: 499) established the family-group name Gerardiidae (as “Gerardidae”). Brook (1889: 79) wrote “Savagliidae, n[om]. n[ov]. (Gerardidae [*sic*], Verrill)”. Brook (1889: 79, 80) based this family-group name based on the emended name *Savaglia* Nardo, 1877, as he did not have access to Nardo (1844) (discussed above). The family-group name Savagliidae proposed by Brook (1889: 79) is herein regarded to have been proposed independently of Savaliidae Nardo, 1844 (based on *Savalia* Nardo, 1844). As their respective type genera are synonyms (as discussed above), the family-group names Savaliidae Nardo, 1844, Gerardiidae Verrill, 1865, and Savagliidae Brook, 1889, are all synonyms.

The oldest and therefore valid family-group name for *Savalia savaglia* (Bertoloni, 1819) is therefore Savaliidae Nardo, 1844. The type genus of Savaliidae Nardo, 1844, (i.e., *Savalia* Nardo, 1844), is however currently assigned to the family Parazoanthidae Delage and Hérourard, 1901 (Sinniger et al. 2013: [2], [3]). This means that the family-group names Savaliidae Nardo, 1844, Gerardiidae Verrill, 1865, and Savagliidae Brook, 1889, are all subjective synonyms of Parazoanthidae Delage and Hérourard, 1901. To maintain current and widespread use of Parazoanthidae Delage and Hérourard, 1901, the best course of action would be to enact Article 23.9 of the Code (ICZN 1999: 27, 28) to reverse precedence between Parazoanthidae Delage and Hérourard, 1901, and the other three family-group names. This is not possible, however, as all three family-group names have been used after 1899, and Article 23.9.1.1 of the Code (ICZN 1999: 28) cannot be satisfied.

In accordance with Article 23.9.2 (ICZN 1999: 28, 29), an application is being prepared to request that the International Commission on Zoological Nomenclature suppress the senior subjective synonyms Savaliidae Nardo, 1844, Gerardiidae Verrill, 1865, and Savagliidae Brook, 1889, in favour of Parazoanthidae Delage and Hérourard, 1901, to maintain current and widespread usage.

Sphenopidae Hertwig, 1882, and the hitherto overlooked subjective synonym Palythoidae Duchassaing de Fonbressin and Michelotti, 1860

The family-group name Sphenopidae was established by Hertwig (1882: 120) for the genus *Sphenopus* Steenstrup, 1856. The name Sphenopidae Hertwig, 1882, had been used sporadically since it was established. Since it was reinstated by Ryland et al. (2000: 191, 192) for the genera *Sphenopus* Steenstrup, 1856, and *Palythoa* Lamouroux, 1816, however, the name Sphenopidae Hertwig, 1882, is now in current and widespread use (see references cited below).

This current and widespread use of the family-group name Sphenopidae Hertwig, 1882, is threatened by the hitherto overlooked subjective synonym Palythoidae Duchassaing de Fonbressin and Michelotti, 1860 (type genus *Palythoa* Lamouroux, 1816).

This family-group name was first established as “Polythoae” for “*Polythoa*”, “*Bergia* n. g.” and “*Gemmaria* n. g.” (Duchassaing de Fonbressin and Michelotti, 1860: 37). That Duchassaing de Fonbressin and Michelotti (1860: 37) intended this to be

a family-group name is evidenced by their use of “Zoanthidae” (containing “*Zoanthus*”, “*Isaura*”, “*Mammillifera*” and “*Orinia* n. g.”) as a counterpart to “Polythoae”. The name “*Polythoa*” is an incorrect subsequent spelling of *Palythoa* Lamouroux, 1816 (the first usage of which appears to be by Schweigger 1819: 100).

As type genus of Sphenopidae Hertwig, 1882, and of Palythoidea Duchassaing de Fombressin and Michelotti, 1860, are considered to belong to the same family grouping (e.g., Ryland et al. 2000: 192; Reimer et al. 2012b: 45, 47, 49; Irei et al. 2015: 2, 3, 14–16; Reimer and Fujii 2016: 14, 19), both family-group names are therefore subjective synonyms.

To prevent the nomenclatural and taxonomic destabilisation that would result from the replacement of Sphenopidae Hertwig, 1882, with its subjective synonym Palythoidea Duchassaing de Fombressin and Michelotti, 1860, as required by the Principle of Priority (Article 23, ICZN 1999: 24) requires that the oldest available name for the taxon under consideration must be used.

To mediate the Principle of Priority, Article 23.9.1 of the Code (ICZN 1999: 27) allows for a reversal of precedence of a junior synonym when the senior synonym has not been used as a valid name after 1899 (Article 23.9.1.1) and the junior synonym “has been used for a particular taxon, as its presumed valid name, in at least 25 works, published by at least 10 authors in the immediately preceding 50 years and encompassing a span of not less than 10 years” (Article 23.9.1.2).

Since 1899, the name Palythoidea has been used in three publications. The first usage was by Barel and Kramers (1977: 32) used the term “species of Palythoidea” for an unidentified zoantharian associated with echinoderms. From the context, it appears that Barel and Kramers (1977: 32) were reporting on material with close affinities to *Palythoa* Lamouroux, 1816.

As Ng & Low (2010:37, 38) have argued, “valid usage” of a name must be unambiguous and show clearly that the author[s] both considered it the correct name to be used and adopted the name. According to these criteria, the usage of “Palythoidea” by Barel and Kramers (1977: 32) cannot be considered to be valid.

The second usage was by Herbets (1972: 125) who stated: “La disposition brachynémique des mésentères est un autre caractère des Palythoidea”. That Herbets (1972: 125) was not using “Palythoidea” as a valid family-group name is evidenced by the fact that the section in which the term “Palythoidea” appears is headed by “Zoanthidae” (p. 106), and in a diagram detailing the classification of zoantharians, “*Palythoa*”, “*Isaurus*”, “*Spenopus* [sic]” and “*Zoanthus*” are clearly placed in “Zoanthidae” (p. 142).

The third usage is by Pax and Müller (1957: 3, 4) in which they recognise the “Unterfamilie Palythoinae” under Epizoanthidae, and further provide a key for identifying this subfamily. Clearly, Pax and Müller (1957: 3, 4) were recognising the family-group name Palythoidea as valid.

The name Palythoidea Duchassaing de Fombressin and Michelotti, 1860, thus has been used as the valid name for the taxon is denotes since 1899 (and Article 23.9.1.1 of the Code cannot be fulfilled).

Article 23.9.1.2 of the Code is fulfilled as the family-group name Sphenopidae Hertwig, 1882, is in current and widespread usage, as evidenced by the 30 publications by 99 different authors over the past 34 years using Sphenopidae as a valid name for the taxon it denotes (viz., Nagabhushanam and Jothinayagam 1982: 17; Ryland et al. 2000: 191, 192; Ryland and Lancaster 2003: 407, 409, 415; Ryland and Lancaster 2004: 180; Ryland and Westphalen 2004: 411; Ryland et al. 2004: 1195, 1197; Acosta et al. 2005: 147–149, 151, 154, 160; Sinniger et al. 2005: 1122, 1125, 1126; Daly et al. 2007: 144; Sinniger et al. 2008: 1254, 1256, 1257; Fautin and Daly 2009: 356; Del Mónaco et al. 2010: 360; Reimer and Sinniger 2010: 251; Reimer et al. 2010c: 606, 616; Swain 2010: 2592; Reimer et al. 2011a: 983, 985, 987, 989, 991, 992; Cavallari et al. 2012: 25; Longo et al. 2012: [1]; Palmer et al. 2012: 3880; Reimer et al. 2012b: 43, 45, 47, 49; Rodríguez-Viera et al. 2012: 32; Costello et al. 2013: [2]; Fujii and Reimer 2013: 510, 516; Krishna and Gophane 2013: 210; Koupaei et al. 2014: 64; Alencar et al. 2015: 1113, 1114, 1121; Irei et al. 2015: 1, 2, 4, 6, 14–16, 20; Qin et al. 2015: 100; De la Cruz-Francisco et al. 2016: 24; Fujii and Reimer 2016: 11, 12, 14, 17, 19, 20; Risi and Macdonald 2016: 113).

As precedence of the family-group names Sphenopidae Hertwig, 1882, and Palythoidea Duchassaing de Fonbressin and Michelotti, 1860, cannot be replaced, the former will need to be replaced by the latter. To prevent resulting nomenclatural instability, and in accordance with Article 23.9.2 (ICZN 1999: 28, 29), an application is being prepared to request that the International Commission on Zoological Nomenclature suppress the senior subjective synonym Palythoidea Duchassaing de Fonbressin and Michelotti, 1860, in favour of Sphenopidae Hertwig, 1882, to maintain current and widespread usage.

The type species of *Palythoa* Lamouroux, 1816

The genus-group name *Palythoa* (with the simultaneous French vernacular ‘Palythoé’) was established by Lamouroux (1816: 359–362) with the inclusion of two species, which were rendered French vernacular: ‘Palythoé Etoilée’ and ‘Palythoé Ocellée’. The second species causes no problems “*P. Ocellata* ... Sol. et Ell., p. 180, n. 6, tab. 1, fig. 6 (*A. Ocellatum*) ...” is listed. This is *Alcyonium ocellatum* Ellis, in Ellis and Solander, 1786, a synonym of *Alcyonium mammillosum* Ellis, in Ellis and Solander, 1786 (see Pax 1910a: 102; 1910b: 258; Acosta et al. 2005: 159). The French vernacular name ‘Palythoé Etoilée’, however, is not the translation of the name “*P[alythoa] Mamillosa* [sic] ... Sol. et Ell., p. 179, n. 5, tab. 1, figs. 4–5 (*A. Mamillosum* [sic]) ...” given in the synonymy.

On an unnumbered page of errata after page 559 of Lamouroux (1816) gives “[page] 361 ... [line] 12 ... *Mamillosa*, lisez *Stellata*”, and the captions on page 558 to plate 14 (and on the plate itself) of Lamouroux (1816) gives “*Palythoe* [sic] *stellata*, p. 361. figure copiée dans Ellis”. It is thus clear that it was the intention of Lamouroux (1816: 361, 558, unnumbered errata page, pl. 14, fig. 2, caption) to rename *Alcyonium*

mammillosum Ellis, in Ellis and Solander, 1786, *Palythoa stellata*. *Palythoe* [sic] *stellata* Lamouroux, 1816, is also a junior objective synonym of *Alcyonium mammillosum* Ellis, in Ellis and Solander, 1786, as Lamouroux (1816: 361, 558, unnumbered errata page, pl. 14, fig. 2, caption) proposed the former as a replacement name for the latter.

Alcyonium ocellatum Ellis, in Ellis and Solander, 1786, and *Palythoa stellata* Lamouroux, 1816, are thus to be considered species originally included in the genus *Palythoa* Lamouroux, 1816, and were both eligible for fixation (Article 67.2 of the Code, ICZN 1999: 67).

The type species of the genus *Palythoa* Lamouroux, 1816, is accepted as *Alcyonium mammillosum* Ellis, in Ellis and Solander, 1786, by subsequent designation by Haddon and Shackleton (1891b: 691) (see Low and Reimer 2011b: 63). As Lamouroux (1816: 361, 558, unnumbered errata page, pl. 14, fig. 2, caption) proposed the replacement name *Palythoa stellata* for *Alcyonium mammillosum* Ellis, in Ellis and Solander, 1786, the latter cannot be considered an originally included species in the sense of Article 67.2 of the Code (ICZN 1999: 67).

Article 69.2.2 of the Code (ICZN 1999: 72) states that “[i]f an author designates as type species a nominal species that was not originally included (or accepts another’s such designation) and if, but only if, at the same time he or she places that nominal species in synonymy with one and only one of the originally included species (as defined in Article 67.2), that act constitutes fixation of the latter species as type species of the nominal genus or subgenus”.

Haddon and Shackleton (1891b: 691) wrote that “*Palythoa mammillosa* is evidently regarded by Lamouroux [1816] as the type species of the genus. He reproduces Solander’s figure of this species, but not that of *P. ocellata*, of which he merely gives a description. Unfortunately a Latinized version of the French name ‘*Palythoé Etoillée*,’ given by Lamouroux to *P. mammillosa*, has been added at the bottom of his plate—a circumstance which has given rise to some confusion”.

Clearly, Haddon and Shackleton (1891b: 691) considered *Alcyonium mammillosum* Ellis, in Ellis and Solander, 1786, to be a synonym of *Palythoe* [sic] *stellata* Lamouroux, 1816, as well as the type species of *Palythoa* Lamouroux, 1816. The conditions of Article 69.2.2 of the Code (ICZN 1999: 72) are met and Haddon and Shackleton (1891b: 691) are deemed to have designated *Palythoe* [sic] *stellata* Lamouroux, 1816, as the type species of *Palythoa* Lamouroux, 1816. As both names are objective synonyms, the valid name for the species under discussion is *Palythoa mammillosa* (Ellis, in Ellis and Solander, 1786).

Table 1. An updated supraspecific classification of the Zoantharia. All valid genera and their type species are given. The numbers given in parentheses after each suborder and family represent the total number of families and genera, respectively, in each grouping. A total of one order, three suborders, nine families and twenty-seven genera are currently recognised in the Zoantharia.

ORDER ZOANTHARIA RAFINESQUE, 1815

1. SUBORDER BRACHYCNEMINA HADDON AND SHACKLETON, 1891 (3)

1. Neozoanthidae Herberts, 1972 (1)

1. *Neozoanthus* Herberts, 1972
Neozoanthus tulearensis Herberts, 1972

2. Sphenopidae Hertwig, 1882 (2)

2. *Sphenopus* Steenstrup, 1856
Sabella marsupialis Gmelin, 1791
3. *Palythoa* Lamouroux, 1816
Palythoe [sic] *stellata* Lamouroux, 1816

3. Zoanthidae Rafinesque, 1815 (3)

4. *Zoanthus* Lamarck, 1801
Actinia sociata Ellis, 1768
5. *Acrozoanthus* Saville-Kent, 1893
Acrozoanthus australiae Saville-Kent, 1893
6. *Isaurus* Gray, 1828
Isaurus tuberculatus Gray, 1828

2. SUBORDER MACROCNEMINA HADDON AND SHACKLETON, 1891 (5)

4. Epizoanthidae Delage and Hérouard, 1901 (3)

7. *Epizoanthus* Gray, 1867
Dysidea papillosa Johnston, 1842
8. *Paleozoanthus* Carlgren, 1924
Paleozoanthus reticulatus Carlgren, 1924
9. *Thoracactis* Gravier, 1918
Thoracactis topsenti Gravier, 1918

5. Hydrozoanthidae Sinniger, Reimer and Pawlowski, 2010 (2)

10. *Hydrozoanthus* Sinniger, Reimer and Pawlowski, 2010
Parazoanthus tunicans Duerden, 1900
11. *Terrazoanthus* Reimer and Fujii, 2010
Terrazoanthus onoi Reimer and Fujii, 2010

6. Microzoanthidae Fujii and Reimer, 2011 (1)

12. *Microzoanthus* Fujii and Reimer, 2011
Microzoanthus occultus Fujii and Reimer, 2011

7. Nanozoanthidae Fujii and Reimer, 2013 (1)

13. *Nanozoanthus* Fujii and Reimer, 2013
Nanozoanthus harenaceus Fujii and Reimer, 2013

8. Parazoanthidae Delage and Hérouard, 1901 (13)

14. *Parazoanthus* Haddon and Shackleton, 1891
Palythoa axinella Haddon and Shackleton, 1891
15. *Antipathozoanthus* Sinniger, Reimer and Pawlowski, 2010
Gerardia macaronesisicus Ocaña and Brito, 2003
16. *Bergia* Duchassaing de Fombressin and Michelotti, 1860
Bergia catenularis Duchassaing de Fombressin and Michelotti, 1860

2. SUBORDER MACROCNEMINA HADDON AND SHACKLETON, 1891 (5) (Continued)

17. *Bullagummizoanthus* Sinniger, Ocaña and Baco, 2013
Bullagummizoanthus emilyacardiarum Sinniger, Ocaña and Baco, 2013
18. *Corallizoanthus* Reimer, in Reimer, Nonaka, Sinniger and Iwase, 2008
Corallizoanthus tsukaharai Reimer, in Reimer, Nonaka, Sinniger and Iwase, 2008
19. *Hurlizoanthus* Sinniger, Ocaña and Baco, 2013
Hurlizoanthus parrishii Sinniger, Ocaña and Baco, 2013
20. *Isozoanthus* Carlgren, in Chun, 1903
Isozoanthus giganteus Carlgren, in Chun, 1903
21. *Kauluzoanthus* Sinniger, Ocaña and Baco, 2013
Kauluzoanthus kerbyii Sinniger, Ocaña and Baco, 2013
22. *Kulamanamana* Sinniger, Ocaña and Baco, 2013
Kulamanamana haumeaee Sinniger, Ocaña and Baco, 2013
23. *Mesozoanthus* Sinniger and Häussermann, 2009
Mesozoanthus fossii Sinniger and Häussermann, 2009
24. *Savalia* Nardo, 1844
Gorgonia savaglia Bertolini, 1819
25. *Umimayanthus* Montenegro, Sinniger and Reimer, 2015
Umimayanthus chanpuru Montenegro, Sinniger and Reimer, 2015
26. *Zibrowius* Sinniger, Ocaña and Baco, 2013
Zibrowius ammophilus Sinniger, Ocaña and Baco, 2013

3. SUBORDER INCERTA SEDIS (1)**9. Abysoanthisidae Reimer and Fujiwara, in Reimer, Sinniger, Fujiwara, Hirano and Maruyama, 2007 (1)**

27. *Abysoanthis* Reimer and Fujiwara, in Reimer, Sinniger, Fujiwara, Hirano and Maruyama, 2007
Abysoanthis nankaiensis Reimer and Fujiwara, in Reimer, Sinniger, Fujiwara, Hirano and Maruyama, 2007
-