

Updated checklist of the extant Chondrichthyes within the Exclusive Economic Zone of Mexico

Nicolás Roberto Ehemann¹, Lorem del Valle González-González¹,
Jorge Guillermo Chollet-Villalpando², José De La Cruz-Agüero¹

1 Instituto Politécnico Nacional – Centro Interdisciplinario de Ciencias Marinas (CICIMAR–IPN), Colección Ictiológica, Avenida IPN s/n, Colonia Playa Palo Santa Rita, La Paz, Baja California Sur, 23096, México

2 Instituto de Ecología, A.C. – INECOL, Biodiversidad y Sistemática, Carretera Antigua a Coatepec 351, Colonia El Haya, Xalapa, 91070, Veracruz, México

Corresponding author: José De La Cruz-Agüero (jcruz@ipn.mx)

Academic editor: N. Bogutskaya | Received 14 March 2018 | Accepted 22 May 2018 | Published 12 July 2018

<http://zoobank.org/5C1CDCE3-B07A-4E15-B6B7-2C5F36F8C568>

Citation: Ehemann NR, González-González LV, Chollet-Villalpando JG, Cruz-Agüero JDL (2018) Updated checklist of the extant Chondrichthyes within the Exclusive Economic Zone of Mexico. ZooKeys 774: 17–39. <https://doi.org/10.3897/zookeys.774.25028>

Abstract

The checklist presented in this study includes the latest taxonomic and systematic modifications and updates (early 2018) for the Chondrichthyes that inhabit the Exclusive Economic Zone (EEZ) of Mexico. The list is based on a literature review of field-specific books, scientific publications and database information from collections and museums worldwide available online such as, the Ocean Biogeographic Information System (OBIS), Global Biodiversity Information Facility (GBIF), Encyclopedia of Life (EOL), iSpecies, FishBase and the National Biodiversity Information System (SNIB–CONABIO). Information was cross-referenced with digital taxonomic systems such as the Catalog of Fishes of the California Academy of Sciences, the World Register of Marine Species (WoRMS), and the Integrated Taxonomic Information System (ITIS). There is a total of two subclasses two divisions, 13 orders, 44 families, 84 genera, and 217 species that represent approximately 18% of all living and described species of chondrichthyans worldwide. For the Mexican Pacific and the Gulf of California, 92 species of chondrichthyans are listed compared to 94 species for the Gulf of Mexico and the Caribbean Sea. Additionally, 31 species listed occur on both coasts of Mexico. The species richness of the Mexican chondrichthyans will surely continue to increase, due to the exploration of deep-water fishing areas in the EEZ.

Keywords

chimaeras, elasmobranchs, rays, sharks, systematics, taxonomy

Introduction

The natural history of the Chondrichthyes (chimaeras, sharks, skates, and rays) inhabiting the waters of the maritime territory and the adjacent and oceanic zones of Mexico, (referred hereinafter as the Exclusive Economic Zone (EEZ)) has always been a difficult issue to address. The causes of the underestimation or overestimation of these species can be diverse (Del Moral-Flores et al. 2015a) and are typically attributed to i) incorrect taxonomic identifications, ii) undetected synonyms, iii) inaccurate or incorrect curatorial records (e.g., date of collection, location, coordinates), iv) use of obsolete information and, v) absence of updated taxonomic lists.

Recent chondrichthyans studies have led to various taxonomic and systematic readjustments, name substitutions, new gender-specific combinations and description of new species. New taxonomic arrangements are based on combined conventional morphology, geometric morphometrics and DNA studies (e.g., Del Moral-Flores et al. 2015b, Acero et al. 2016, Last et al. 2016a, Last et al. 2016b, Last et al. 2016c, Last et al. 2016d, White and Naylor 2016).

The current systematic checklist of chimaeras, sharks, rays, and skates from the Mexican EEZ, incorporates the newest taxonomic and systematic proposals for chimaeras and sharks (Ebert et al. 2013, Weigmann 2016, 2017), skates and rays (Last et al. 2016a, b, c, d, Weigmann 2016, 2017) and follows, with some nomenclatural modifications, the Chondrichthyes classifications made by Weigmann 2016. The present inventory also includes information provided since last century until last year regarding the chondrichthyans of Mexico and Mexican endemic species (e.g., Castro-Aguirre 1965, Castro-Aguirre et al. 1996, Espinosa-Pérez et al. 2004, Palacios-Salgado et al. 2012, Del Moral-Flores and Pérez-Ponce de León 2013, Del Moral-Flores et al. 2015a, Del Moral-Flores et al. 2016).

In Mexico, an updated inventory of the natural resources relevant to Chondrichthyes is necessary, and many ecological relationships are also unknown. This information is imperative for developing environmental or biogeographic theory and for explaining scientific implications of the availability of resources (Siqueiros-Beltrones and De La Cruz-Agüero 2004). Systematic lists are thus a fundamental requirement for the decision-making process in the evaluation of the changes in biodiversity derived from anthropogenic factors and consequently, for the establishment of regional systems and priority areas with regard to the conservation of Mexican chondrichthyans. Due to the constant updating of taxonomic and systematic information for Chondrichthyes, the primary objective of this study is to provide an updated list (early 2018) of the chimaeras, sharks, skates, and rays, living in the EEZ of Mexico.

Materials and methods

First, the Weigmann taxonomic checklists (2016, 2017) were consulted to select the species that were registered in the areas defined by the latter author (see also IHO 1953)



Figure 1. Map showing the Exclusive Economic Zone (EEZ) of Mexico. The zone comprises, including islands and territorial sea, approximately 3,150,000 km². Modified from CONABIO (2011), available at: http://www.conabio.gob.mx/informacion/gis/layouts/contdv250_zeemgw.png

as northeastern Pacific (NEP, Canada to Panama) and northwestern Atlantic (NWA, eastern United States of America to the southern Caribbean Sea). In this geographical framework, all chondrichthyan species with a distribution identified within the Mexican EEZ were included. The selected taxocenoses were contrasted and compared with the biological stocks recorded in the Ichthyological Collection (IC) (<http://coleccion.cicimar.ipn.mx/>) of the Centro Interdisciplinario de Ciencias Marinas (marked with an asterisk in Table 1). The Mexican EEZ is the area of ocean extending 200 nautical miles (370.4 km) from the coast (DOF 1976). This geographic boundary refers to a political delineation rather than an ecological boundary (Figure 1).

Field-specific books were consulted (e.g., Compagno 1999, Castro 2011a, Ebert et al. 2013, Last et al. 2016a) for comparative purposes and several recent scientific publications were reviewed in a deliberate manner (e.g., Ruiz-Campos et al. 2010, Hoyos-Padilla et al. 2013, Del Moral-Flores et al. 2015a, Del Moral-Flores et al. 2016, Villalobos et al. 2016, Weigmann 2016, 2017). In this way, a clear, cumulative database was compiled with all existing records for species within the Mexican EEZ. This database was cross-referenced with information available online from biological collections and museums worldwide. Global databases consulted included the Ocean Biogeographic Information System (OBIS 2017), the Global Biodiversity Informa-

Table 1. A systematic listing of the chondrichthyans inhabiting the Exclusive Economic Zone of Mexico (EEZ). The checklist was depurated from references, field books, catalogued specimens and online databases (see text) about the distribution of Chondrichthyes in the littoral areas of the Pacific Ocean (including the Gulf of California) and the Gulf of Mexico (including the Caribbean Sea). The systematic arrangement is based on Weigmann (2016), with some modifications for the batomorphs cited by Last et al. (2016a). The species are listed in alphabetical order. (*) = Species with specimens cataloged in the Ichthyological Collection (IC) of CICIMAR–IPN (<http://coleccion.cicimar.ipn.mx/>). (†) = New valid taxa according to Weigmann (2017) and Last et al. (2016a), taking into account that when a genus is designated, all species are included. (?) = Species whose validity requires further investigation. (West C.) = west coast of the EEZ of Mexico. (East C.) = east coast of the EEZ of Mexico.

Subclass	Order	Family	Genus	Species	West C.	East C.
Holocephali	Chimaeriformes	Rhinochimaeridae	<i>Harriotta</i>	<i>Harriotta baecckeli</i> Karrer, 1972	✓	
			<i>Rhinochimaera</i>	<i>Harriotta raleighana</i> Goode & Bean, 1895	✓	
		Chimaeridae		<i>Rhinochimaera atlantica</i> Holte & Byrne, 1909		✓
				<i>Hydrolagus alberti</i> Bigelow & Schroeder, 1951		✓
				<i>Hydrolagus collieri</i> * (Lay & Bennett, 1839)		✓
				<i>Hydrolagus macrophthalmus</i> De Buen, 1959		✓
	<i>Hydrolagus melanophasma</i> * James, Ebert, Long & Didier, 2009		✓			
			<i>Hydrolagus minabilis</i> (Collett, 1904)		✓	
Euselachii						
Infraclass						
Elasmobranchii						
Division	Heterodontiformes	Heterodontidae	<i>Heterodontus</i>	<i>Heterodontus francisci</i> * (Girard, 1855)	✓	
				<i>Heterodontus mexicanus</i> * Taylor & Castro-Aguirre <i>Ginglymostoma cirratum</i> (Bonnaterre, 1788)	✓	✓
Selachii	Orectolobiformes	Ginglymostomatidae	<i>Ginglymostoma</i>	<i>Ginglymostoma unami</i> * Del Moral-Flores, Ramírez-Antonio, Angulo & Pérez-Ponce de León, 2015	✓	
				<i>Rhincodon typus</i> Smith, 1828	✓	✓
	Lamniformes	Alopiidae	<i>Alopius</i>	<i>Alopius pelagicus</i> Nakamura, 1935	✓	✓
<i>Alopius superciliosus</i> Lowe, 1841 <i>Alopius vulpinus</i> (Bonnaterre, 1788)				✓	✓	

Subclass	Order	Family	Genus	Species	West C.	East C.
Lamniformes	Cetorhinidae	<i>Cetorhinus</i>	<i>Cetorhinus maximus</i> (Gunnerus, 1765)	✓	✓	
			<i>Carcharodon carcharias</i> (Linnaeus, 1758)	✓	✓	
	Lamnidae	<i>Isurus</i>	<i>Isurus paucus</i> Rafinesque, 1810	✓	✓	
			<i>Isurus paucus</i> Guitart, 1966	✓	✓	
			<i>Lamna ditropis</i> Hubbs & Follett, 1947	✓	✓	
	Megachasmaidae	<i>Megachasma</i>	<i>Megachasma pelagios</i> Taylor, Compagno & Struhsaker, 1983	✓	✓	
			<i>Carcharias taurus</i> Rafinesque, 1810	✓	✓	
	Odontaspidae	<i>Odontaspis</i>	<i>Odontaspis ferox</i> (Risso, 1810)	✓	✓	
			<i>Odontaspis noronhai</i> (Maul, 1955)	✓	✓	
			<i>Pseudocarcharias kamoharui</i> * (Matsubara, 1936)	✓	✓	
	Pseudocarchariidae	<i>Pseudocarcharias</i>	<i>Carcharhinus acronotus</i> (Poey, 1860)	✓	✓	
			<i>Carcharhinus albimarginatus</i> (Rüppell, 1837)	✓	✓	
			<i>Carcharhinus altimus</i> (Springer, 1950)	✓	✓	
			<i>Carcharhinus brachyurus</i> (Günther, 1870)	✓	✓	
			<i>Carcharhinus brevipinna</i> (Müller & Henle, 1839)	✓	✓	
			<i>Carcharhinus cerdale</i> Gilbert, 1898	✓	✓	
			<i>Carcharhinus falciformis</i> (Müller & Henle, 1839)	✓	✓	
			<i>Carcharhinus galapagensis</i> (Snodgrass & Heller, 1905)	✓	✓	
			<i>Carcharhinus isodon</i> (Müller & Henle, 1839)	✓	✓	
			<i>Carcharhinus leucas</i> (Müller & Henle, 1839)	✓	✓	
<i>Carcharhinus limbatus</i> * (Müller & Henle, 1839)			✓	✓		
<i>Carcharhinus longimanus</i> (Poey, 1861)			✓	✓		
Carcharhiniformes	Carcharhinidae	<i>Carcharhinus</i>	<i>Carcharhinus melanopterus</i> (Quoy & Gaimard, 1824)	✓	✓	
			<i>Carcharhinus obscurus</i> (Lesueur, 1818)	✓	✓	
			<i>Carcharhinus perezii</i> (Poey, 1876)	✓	✓	
			<i>Carcharhinus plumbeus</i> (Nardo, 1827)	✓	✓	
			<i>Carcharhinus porosus</i> * (Ranzani, 1839)	✓	✓	
			<i>Carcharhinus signatus</i> (Poey, 1868)	✓	✓	
			<i>Galeocerdo cuvier</i> (Péron & Lesueur, 1822)	✓	✓	
			<i>Nasolamia velox</i> * (Gilbert, 1898)	✓	✓	

Subclass	Order	Family	Genus	Species	West C.	East C.
Selachii	Carcharhiniformes	Carcharhinidae	<i>Negaprion</i>	<i>Negaprion brevirostris</i> (Poey, 1868)	✓	✓
			<i>Prionace</i>	<i>Prionace glauca</i> * (Linnaeus, 1758)	✓	✓
			<i>Rhizoprionodon</i>	<i>Rhizoprionodon longurio</i> * (Jordan & Gilbert, 1882)	✓	
				<i>Rhizoprionodon porosus</i> (Poey, 1861)	✓	
				<i>Rhizoprionodon terraenovae</i> (Richardson, 1836)	✓	
				<i>Triacnodon obesus</i> (Rüppell, 1837)	✓	
			<i>Apristurus</i>	<i>Apristurus brunneus</i> (Gilbert, 1892)	✓	
				<i>Apristurus kampae</i> Taylor, 1972	✓	
				<i>Apristurus laurussonii</i> (Saemundsson, 1922)	✓	
				<i>Apristurus nasutus</i> de Buen, 1959	✓	
		Pentanchidae	<i>Apristurus parvipinnis</i> Springer & Heemstra, 1979	✓		
			<i>Apristurus riveri</i> Bigelow & Schroeder, 1944	✓		
			<i>Cephalurus cephalus</i> (Gilbert, 1892)	✓		
			<i>Galeus anae</i> (Nichols, 1927)	✓		
		<i>Galeus</i>	<i>Galeus piperatus</i> Springer & Wagner, 1966	✓		
			<i>Parmaturus campechianus</i> Springer, 1979	✓		
			<i>Parmaturus xaniurus</i> * (Gilbert, 1892)	✓		
			<i>Cephaloscyllium ventriosum</i> * (Garman, 1880)	✓		
		Scyliorhinidae	<i>Scyliorhinus hesperius</i> Springer, 1966	✓		
			<i>Scyliorhinus meadi</i> Springer, 1966	✓		
			<i>Scyliorhinus retifer</i> (Garman, 1881)	✓		
			<i>Sphyrna corona</i> Springer, 1940	✓		
		<i>Sphyrna</i>	<i>Sphyrna lewini</i> * (Griffith & Smith, 1834)	✓		
			<i>Sphyrna media</i> Springer, 1940	✓		
			<i>Sphyrna mokarran</i> (Rüppell, 1837)	✓		
			<i>Sphyrna tiburo</i> (Linnaeus, 1758)	✓		
		<i>Galeorhinus</i>	<i>Sphyrna zygaena</i> * (Linnaeus, 1758)	✓		
<i>Galeorhinus galeus</i> (Linnaeus, 1758)	✓					
Triakidae	<i>Mustelus</i>	<i>Mustelus albipinnis</i> * Castro-Aguirre, Antuna-Mendiola, González-Acosta & De La Cruz-Agüero, 2005	✓			

Subclass	Order	Family	Genus	Species	West C.	East C.				
Carcharhiniformes	Triakidae		<i>Mustelus</i>	<i>Mustelus californicus</i> * Gill, 1864	✓					
				<i>Mustelus canis</i> (Mitchill, 1815)		✓				
				<i>Mustelus dorsalis</i> Gill, 1864		✓				
				<i>Mustelus henlei</i> * (Gill, 1863)		✓				
				<i>Mustelus higmani</i> Springer & Lowe, 1963		✓				
				<i>Mustelus lunulatus</i> * Jordan & Gilbert, 1882		✓				
				<i>Mustelus norrisi</i> Springer, 1939		✓				
				<i>Mustelus sinuomexicanus</i> Heemstra, 1997		✓				
				<i>Triakis semifasciata</i> * Girard, 1855		✓				
				<i>Chlamydoselachus</i>				✓		
	Hexanchiformes	Hexanchidae		<i>Hexanchus</i>	<i>Heptranchias</i>		✓			
					<i>Hexanchus griseus</i> * (Bonnaterre, 1788)		✓			
					<i>Hexanchus nakamurai</i> Teng, 1962		✓			
					<i>Hexanchus vitulus</i> Springer & Waller, 1969		✓			
					<i>Notorynchus</i>		✓			
					<i>Notorynchus cepedianus</i> (Péron, 1807)		✓			
					<i>Centrophorus</i>				✓	
					<i>Centrophorus granulosus</i> (Bloch & Schneider, 1801)				✓	
					<i>Centrophorus uyato</i> (Rafinesque, 1810)				✓	
					<i>Dalatius</i>				✓	
Squaliformes	Etmopteridae		<i>Etmopterus</i>	<i>Etmopterus bispinatus</i> (Quoy & Gaimard, 1824)		✓				
				<i>Isistius brasiliensis</i> (Quoy & Gaimard, 1824)		✓				
				<i>Isistius plutodus</i> Garrick & Springer, 1964		✓				
				<i>Squaliolus</i>		✓				
				<i>Squaliolus laticaudus</i> Smith & Radcliffe, 1912		✓				
				<i>Centrosyllium</i>		✓				
				<i>Centrosyllium nigrum</i> * Garman, 1899		✓				
				<i>Etmopterus bullisi</i> Bigelow & Schroeder, 1957		✓				
				<i>Etmopterus billianus</i> (Poey, 1861)		✓				
				<i>Etmopterus schultzei</i> Bigelow, Schroeder & Springer, 1953		✓				
Oxynoridae			<i>Oxynotus</i>	<i>Etmopterus virens</i> Bigelow, Schroeder & Springer, 1953		✓				
				<i>Oxynotus caribbaeus</i> Cervigón, 1961		✓				
Somniosidae			<i>Centroscyrmus</i>	<i>Centroscyrmus coelepis</i> Barbosa du Bocage & de Brito Capello, 1864		✓				

Selachii

Subclass	Order	Family	Genus	Species	West C.	East C.		
Selachii	Squaliformes	Somniosidae	<i>Centroscymnus</i>	<i>Centroscymnus austonii</i> Garman, 1906		✓		
			<i>Somniosus</i>	<i>Somniosus microcephalus</i> (Bloch & Schneider, 1801)		✓		
				<i>Somniosus pacificus</i> Bigelow & Schroeder, 1944	✓			
			<i>Zameus</i>		<i>Zameus squamulosus</i> (Günther, 1877)		✓	
			<i>Cirrhigaleus</i>		<i>Cirrhigaleus asper</i> (Merrett, 1973)		✓	
			Squalidae		<i>Squalus acanthias</i> * Linnaeus, 1758	✓	✓	
				<i>Squalus cubensis</i> Howell Rivero, 1936		✓		
				<i>Squalus mitsukurini</i> Jordan & Snyder, 1903		✓		
				<i>Squalus suckleyi</i> (Girard, 1855)	✓			
			Echinorhinidae	<i>Echinorhinus</i>	<i>Echinorhinus brucus</i> (Bonnaterre, 1788)		✓	
					<i>Echinorhinus cookei</i> Pletschmann, 1928	✓		
					<i>Squatina californica</i> * Ayles, 1859	✓		
			Squatiniiformes	Squatina		<i>Squatina dumeril</i> Lesueur, 1818		✓
						<i>Squatina heteroptera</i> ? Castro-Aguirre, Espinosa Pérez & Huidobro Campos, 2007		✓
						<i>Squatina mexicana</i> ? Castro-Aguirre, Espinosa Pérez & Huidobro Campos, 2007		✓
Batomorphi	Anacanthobatidae	Springeria †		<i>Springeria foliostris</i> (Bigelow & Schroeder, 1951)		✓		
				<i>Springeria longirostris</i> Bigelow & Schroeder, 1962		✓		
				<i>Bathyrhaja abyssicola</i> (Gilbert, 1896)	✓			
	Arhynchobatidae	Bathyrhaja		<i>Bathyrhaja interrupta</i> (Gill & Townsend, 1897)		✓		
				<i>Bathyrhaja spinosissima</i> (Beebe & Tee-Van, 1941)	✓			
				<i>Bathyrhaja trachura</i> (Gilbert, 1892)	✓			
	Rajiformes	Pseudonaja		<i>Pseudonaja fischeri</i> Bigelow & Schroeder, 1954		✓		
				<i>Cruriraja poeyi</i> Bigelow & Schroeder, 1948		✓		
				<i>Cruriraja rugosa</i> Bigelow & Schroeder, 1958		✓		
	Gurgessiellidae †	Fenestraria		<i>Fenestraria ishiyamai</i> (Bigelow & Schroeder, 1962)		✓		
				<i>Fenestraria plutonia</i> (Garman, 1881)		✓		
				<i>Fenestraria sinuata</i> (Bigelow & Schroeder, 1950)		✓		

Subclass	Order	Family	Genus	Species	West C.	East C.	
Batomorphi	Rajiformes	Gurgesiellidae †	<i>Gurgesiella</i>	<i>Gurgesiella atlantica</i> (Bigelow & Schroeder, 1962)		√	
			<i>Amblyraja</i>	<i>Amblyraja badia</i> (Garman, 1899)	√		
					<i>Amblyraja hyperborea</i> (Collett, 1879)	√	
					<i>Beringaja binoculata</i> (Girard, 1855)	√	
					<i>Beringaja cortezensis</i> † (McEachran & Miyake, 1988)	√	
				<i>Beringaja</i>	<i>Beringaja inornata</i> †* (Jordan & Gilbert, 1881)	√	
					<i>Beringaja rhina</i> † (Jordan & Gilbert, 1880)	√	
					<i>Beringaja stellulata</i> † (Jordan & Gilbert, 1880)	√	
				<i>Brevinaja</i>	<i>Brevinaja colesi</i> Bigelow & Schroeder 1948		√
					<i>Brevinaja spinosa</i> Bigelow & Schroeder, 1950		√
					<i>Dactylobatus armatus</i> Bean & Weed, 1909		√
				<i>Dactylobatus</i>	<i>Dactylobatus clar-kii</i> (Bigelow & Schroeder, 1958)		√
					<i>Dipturus bullisi</i> (Bigelow & Schroeder, 1962)		√
					<i>Dipturus garricki</i> (Bigelow & Schroeder, 1958)		√
				<i>Dipturus</i>	<i>Dipturus olseni</i> (Bigelow & Schroeder, 1951)		√
					<i>Dipturus oregoni</i> (Bigelow & Schroeder, 1958)		√
					<i>Dipturus teevani</i> (Bigelow & Schroeder, 1951)		√
					<i>Leuconaja garmani</i> (Whitley, 1939)		√
				<i>Leuconaja</i>	<i>Leuconaja lentiginosa</i> (Bigelow & Schroeder, 1951)	√	
					<i>Leuconaja yucatanensis</i> (Bigelow & Schroeder, 1950)		√
					<i>Rajella fuliginea</i> (Bigelow & Schroeder, 1954)		√
				<i>Rajella</i>	<i>Rajella purpuriventralis</i> † (Bigelow & Schroeder, 1962)		√
					<i>Rostronaja ackleyi</i> (Garman, 1881)		√
					<i>Rostronaja eglanteria</i> (Bosc, 1800)		√
				<i>Rostronaja</i> †	<i>Rostronaja equatorialis</i> * (Jordan & Bollman, 1890)	√	
					<i>Rostronaja texana</i> (Chandler, 1921)		√
					<i>Rostronaja velezi</i> (Chirichigno, 1973)	√	
		<i>Diplobatis</i>	<i>Diplobatis ommata</i> * (Jordan & Gilbert, 1890)	√			
		<i>Narcine</i>	<i>Narcine bancroftii</i> (Griffith & Smith, 1834)		√		
			<i>Narcine brasiliensis</i> (Olfers, 1831)		√		
		Narcinidae					
	Torpediniformes						

Subclass	Order	Family	Genus	Species	West C.	East C.
Torpediniformes	Narcinidae		<i>Narcine</i>	<i>Narcine entemedor</i> * Jordan & Starks, 1895	✓	
				<i>Narcine vermiculatus</i> Breder, 1928	✓	
	Torpedinidae		<i>Torronare</i>	<i>Torronare californica</i> (Ayres, 1855)	✓	
				<i>Torronare nobiliana</i> (Bonaparte, 1835)		✓
				<i>Torpedo andersoni</i> Bullis, 1962		✓
				<i>Platyrhinoidis triseriata</i> * (Jordan & Gilbert, 1880)	✓	
	Pristidae		<i>Pristis</i>	<i>Pristis pectinata</i> Latham, 1794		✓
				<i>Pristis pristis</i> (Linnaeus, 1758)	✓	✓
				<i>Pseudobatos glaucostigmus</i> * (Jordan & Gilbert, 1883)	✓	
				<i>Pseudobatos lentiginosus</i> (Garman, 1880)		✓
<i>Pseudobatos leucorhynchus</i> * (Günther, 1866)				✓		
<i>Pseudobatos percellens</i> (Walbaum, 1792)					✓	
Rhinobatiformes †	Rhinobatidae		<i>Pseudobatos planiceps</i> (Garman, 1880)	✓		
			<i>Pseudobatos prahli</i> (Acero & Franke, 1995)	✓		
			<i>Pseudobatos productus</i> * (Ayres, 1854)	✓		
			<i>Pseudobatos spinosus</i> ? (Günther, 1870)	✓		
			<i>Zapteryx exasperata</i> * (Jordan & Gilbert, 1880)	✓		
			<i>Zapteryx xyster</i> Jordan & Evermann, 1896	✓		
Myliobatiformes	Trygonorhinidae †		<i>Zapteryx</i>	<i>Aetobatus narinari</i> (Euphrasen, 1790)	✓	
				<i>Aetobatus laticeps</i> * † (Gill, 1865)	✓	
	Aetobatidae †		<i>Aetobatus</i>	<i>Hypanus americanus</i> (Hildebrand & Schroeder, 1928)		✓
				<i>Hypanus dipterurus</i> * (Jordan & Gilbert, 1880)	✓	
	Dasyatiidae		<i>Hypanus</i> †	<i>Hypanus guttatus</i> (Bloch & Schneider, 1801)		✓
				<i>Hypanus longus</i> (Garman, 1880)	✓	
				<i>Hypanus sabinus</i> (Lesueur, 1824)		✓
				<i>Hypanus say</i> (Lesueur, 1817)		✓
				<i>Pteroplatyrygon violacea</i> (Bonaparte, 1832)	✓	
				<i>Syracura pacifica</i> (Beebe & Tee-Van, 1941)	✓	
Potamotrygonidae		<i>Syracura schmarda</i> (Werner, 1904)	✓			

Subclass	Order	Family	Genus	Species	West C.	East C.
		Gymnuridae	<i>Gymnura</i>	<i>Gymnura crebripunctata</i> * (Peters, 1869)	✓	
				<i>Gymnura marmorata</i> * (Cooper, 1864)	✓	
				<i>Gymnura lessae</i> Yokota & De Carvalho, 2017		✓
				<i>Mobula alfredi</i> † (Krefft, 1868)		✓
				<i>Mobula birostris</i> † (Walbaum, 1792)	✓	✓
				<i>Mobula hypostoma</i> (Bancroft, 1831)		✓
		Mobulidae	<i>Mobula</i>	<i>Mobula mobular</i> (Bonnaterre, 1788)	✓	
				<i>Mobula munkiana</i> Notarbartolo Di Sciara, 1987	✓	
				<i>Mobula tarapacana</i> (Philippi, 1892)	✓	
				<i>Mobula thurstoni</i> (Lloyd, 1908)	✓	
				<i>Myliobatis californica</i> * Gill, 1865	✓	
		Myliobatidae	<i>Myliobatis</i>	<i>Myliobatis longirostris</i> * Applegate & Fitch, 1964	✓	
			<i>Aetomylaeus</i>	<i>Aetomylaeus asperrimus</i> (Gilbert, 1898)	✓	
				<i>Rhinoptera bonasus</i> (Mitschill, 1815)		✓
		Rhinopteridae	<i>Rhinoptera</i>	<i>Rhinoptera brasiliensis</i> Müller, 1836		✓
				<i>Rhinoptera steindachneri</i> * Evermann & Jenkins, 1891	✓	
				<i>Urobatis concentricus</i> * Osburn & Nichols, 1916	✓	
				<i>Urobatis halleri</i> * (Cooper, 1863)	✓	
			<i>Urobatis</i>	<i>Urobatis jamaicensis</i> (Cuvier, 1816)		✓
				<i>Urobatis maculatus</i> * Garman, 1913	✓	
				<i>Urotrygon aspidura</i> * (Jordan & Gilbert, 1882)	✓	
		Urotrygonidae		<i>Urotrygon chilensis</i> * (Günther, 1872)	✓	
				<i>Urotrygon cimar</i> López & Bussing, 1998	✓	
			<i>Urotrygon</i>	<i>Urotrygon munda</i> Gill, 1863	✓	
				<i>Urotrygon nana</i> Miyake & McEachran, 1988	✓	
				<i>Urotrygon rogersi</i> * (Jordan & Starks, 1895)	✓	
				<i>Urotrygon simulatrix</i> Miyake & McEachran, 1988	✓	

tion Facility (GBIF 2017), the Encyclopedia of Life (EOL 2017), iSpecies (2017), the FishBase Project (Froese and Pauly 2017) and the National Biodiversity Information System (SNIB–CONABIO, <https://goo.gl/9aNLSv>).

Chondrichthyes taxa were verified using published references from online resources (i.e., the Internet) such as the Catalog of Fishes of the California Academy of Sciences (<https://goo.gl/S792vp>), World Register of Marine Species (<http://www.marine-species.org/>), Integrated Taxonomic Information System (<https://www.itis.gov/>), and Chondrichthyan Tree of Life (<https://sharksrays.org/>). Online database searches were carried out between June 22 and November 15 of 2017. The systematic arrangement (Table 1) is organized in a manner based on the proposal in Weigmann (2016) with some modifications for the group of skates and rays species (Last et al. 2016a). Finally, the spellings in the citations of the common names of the species (sensu Froese and Pauly 2017) follows Nelson et al. (2002).

Results

The detailed literature review involving the species of chimaeras, sharks, skates, and rays that currently exist worldwide includes 1,212 species (Weigmann 2017). Taxa delineations are in accordance with the phylogenetic system published by Weigmann (2016) for the Chondrichthyes and include two subclasses (Holocephali and Euselachii), one infraclass (Elasmobranchii) with two divisions (Selachii and Batomorphi), 14 orders, 64 families, and 215 genera. The phylogeny includes one order, three families, six genera, and 49 species of chimaeras; nine orders, 34 families, 105 genera, and 517 species of sharks; and four orders, 27 families, 104 genera, and 646 species of skates and rays. For a comparison of the number of species of Chondrichthyes recognized worldwide between 2005 and 2015, see Table 1 of Weigmann (2016).

It should be noted that the specific taxonomic richness of the Chondrichthyes of Mexico may vary slightly within the recent literature (e.g., Weigmann 2016, Last et al. 2016a, Del Moral-Flores et al. 2015a) (approximately ten or fewer species). However, at the family level for the batomorphs, there are important taxonomic implications with a difference of up to nine families. Thus, the proposed classification for the skates and rays by Last et al. (2016a) was included in this systematic listing for the Mexican EEZ, regarding the newest taxa in the Dasyatidae family (Last et al. 2016b), the resurrection of the Aetobatidae family (White and Naylor 2016), the new order Rhinopristiformes (Last et al. 2016c), and nomenclatural changes made in the order Rajiformes (Last et al. 2016d). For a detailed description of the taxonomic changes in Chondrichthyes in general, see Weigmann (2017).

The dynamism (or uncertainty) of the classification of Chondrichthyes can be exemplified by the Thornback guitarfish, *Platyrrhinoidis triseriata* (Jordan & Gilbert, 1880), which is currently recognized in two families, Rhinobatidae (sensu Froese and Pauly 2017) and Platyrrhinidae (sensu Last et al. 2016a), and in four different orders such as, Rajiformes (Froese and Pauly 2017), Rhinobatiformes (Weigmann, 2016), Torpediniformes (Last et al. 2016a) and Myliobatiformes (Nelson et al. 2016). Herein,

for the species mentioned, we follow the criteria for family allocation of Naylor et al. (2012) based on DNA sequences, which concurs with the criteria that the American zoologist Samuel W. Garman suggested more than a century ago (Garman 1913: 290). At the order level, we adopted the proposal depicted in the phylogenetic tree by Yang et al. (unpublished) cited in Last et al. (2016a) followed too, in the Chondrichthyan Tree of Life Project (<https://sharksrays.org/>).

Finally, according to the worldwide species richness of Chondrichthyes, in Mexico, there are no records of one family of Holocephali, the Plownose chimaeras, Callorhynchidae (distributed in the south of the American continent); neither for 11 families of sharks: Parascylliidae, Brachaeluridae, Orectolobidae, Hemiscylliidae, Stegostomatidae, Leptochariidae, Hemigaleidae, Pseudotriakidae, Pristiophoridae, Proscylliidae, and Mitsukurinidae (the last three families having species that inhabit the American continent and are likely to be found in the future within the Mexican EEZ, such as the Goblin shark *Mitsukurina owstoni* Jordan, 1898); and finally not for eight families of skates and rays: Glaucostegidae, Hexatrygonidae, Hypnidae, Rhinidae, Narkidae, Plesiobatidae, Urolophidae and Zanobatidae (none of which are distributed in the American continent).

The Mexican Exclusive Economic Zone (EEZ)

A total of 217 species of chimaeras, sharks, skates, and rays was recorded and classified into two subclasses, one infraclass, two divisions, 13 orders, 44 families, and 84 genera (Table 1). This species richness corresponds to 17.9% of the total of current species of Chondrichthyes worldwide, according to Weigmann (2017). The Euselachii subclass was the most representative in terms of species richness at 96.3%, while the Holocephali subclass was the least represented at only 3.7%. The chimaeras were represented in the Mexican EEZ exclusively by a single order (Chimaeriformes) with two families. Of these, Rhinochimaeridae was the best represented with two genera, while *Hydrolagus* (Chimaeridae) was the genus that had the largest number of species (five species). In terms of sharks, the Carcharhiniformes and Squaliformes orders were represented by six families each. Within Carcharhiniformes, the Carcharhinidae family contained seven genera, and *Carcharhinus* had 18 species resulting with the highest species diversity. Finally, for rays and skates, the order Myliobatiformes was represented by eight families, where the genera *Pseudobatos* (eight species), *Mobula* (seven species) and *Urotrygon* (seven species) were the most diverse at the species level. On the other hand, the family Rajidae (order Rajiformes) contained eight different genera with 27 species (Table 1).

The Pacific Ocean and the Gulf of California (west coast)

For this area of the Mexican EEZ, Chondrichthyes was represented by 92 species found only in this area and 31 species distributed on both coasts (i.e., with an amph-Amer-

ican distribution), for 123 species. These species belong to two subclasses (Table 1), with sharks (Selachii division) representing 51.2% of the species (63 species), followed by rays (Batomorphi division) representing 44.7% of the species (55 species) and finally chimaeras (subclass Holocephali), with 4.1% of the species (five species). Considering both coasts of Mexico, the species richness for this zone represents 49.59% of the total of Chondrichthyes currently recorded for the EEZ (of a total of 248 species including the amphi-American species).

The order Chimaeriformes was represented by two families (Chimaeridae and Rhinochimaeridae) with two genera and five species, representing 62.5% of the chimaera species reported for Mexico (Table 1). The two species of chimaeras from the genus *Harriotta* (Rhinochimaeridae) were distributed exclusively on this basin of the Mexican EEZ.

For the group of sharks, eight different orders were recorded, including 22 families, 36 genera, and 63 species. The Carcharhiniformes were the best represented with six families. Specifically, Carcharhinidae was the most taxonomically diverse with seven genera and 17 species. Of these species, eleven belong to the genus *Carcharhinus*. The Triakidae family includes three genera and seven species, which five species belonged to the genus *Mustelus* (Table 1). The horn sharks, Heterodontidae family, were distributed exclusively within the Pacific Mexican EEZ. On the other hand, the batomorphs were represented by four orders, 16 families, 23 genera, and 55 species (Table 1). The genus *Urotrygon* (Myliobatiformes: Urotrygonidae) contained the highest diversity of batomorph, with seven species (Table 1). The Banded guitarfish *Zapteryx exasperata* (Jordan & Gilbert, 1880) and the Witch guitarfish *Z. xyster* (Jordan & Evermann, 1896) are now members of the new family Trygonorrhinidae (sensu Last et al. 2016c), and they are exclusively distributed in the western slope of the Mexican EEZ.

In this area, the following species were identified as endemic: the Whitemargin smoothhound *Mustelus albiginnis* Castro-Aguirre, Antuna-Mendiola, González-Acosta & De La Cruz-Agüero, 2005 (Carcharhiniformes: Triakidae), the Spotted round ray *Urobatis maculatus* Garman, 1913 (Myliobatiformes: Urotrygonidae), the Cortez skate *Beringraja cortezensis* (McEachran & Miyake, 1988) (Rajiformes: Rajidae) and the Spiny guitarfish *Pseudobatos spinosus* (Günther, 1870) (Rhinopristiformes: Rhinobatidae). For the latter species, see the Discussion.

The Gulf of Mexico and the Caribbean Sea (east coast)

The Exclusive Economic Zone of the eastern slope of Mexico was represented by 94 chondrichthyans species that occur only in this area and 31 amphi-American species (125 in total), belonging to two subclasses (Table 1). Of these, 75 species belong to sharks (60.0%), 47 (37.6%) to skates and rays, and three (2.4%) to chimaeras. Considering both coasts of Mexico, the species richness for this zone represents 50.41% of the total chondrichthyans currently recorded (of 248 species including the amphi-Americans species).

The Holocephali were grouped into one order (Chimaeriformes), two families, two genera, and three species that represent approximately 40% of the species recorded to date for the Mexican EEZ (Table 1). Within the sharks, the Squaliformes and Lamniformes orders contained the most number of families with six families. However, the Carcharhinidae family (order Carcharhiniformes) had the highest number of genera with five genera, and the *Carcharhinus* genus had the highest species richness with 14 species (Table 1). The sharks belonging to the family Centrophoridae were distributed exclusively in this slope of the EEZ of Mexico. For the batomorphs, the order Myliobatiformes contained the largest number of families with seven, where Rajidae with six genera was the most diverse, as its genus *Dipturus* is the taxon with the highest species richness with five species (Table 1).

For the Gulf of Mexico and the Caribbean Sea there were three endemic species of sharks: Campeche catshark *Parmaturus campechiensis* Springer 1979 (Carcharhiniformes: Scyliorhinidae), the Disparate angel shark *Squatina heteroptera* Castro-Aguirre, Espinosa Pérez & Huidobro Campos, 2007 the Mexican angel shark *S. mexicana* Castro-Aguirre, Espinosa Pérez & Huidobro Campos, 2007 (Squatiniiformes: Squatinidae) and one species of skate, the Yucatan skate *Leucoraja yucatanensis* (Bigelow & Schroeder, 1950). However, see in Discussion about these Angel shark species.

Amphi-American Chondrichthyes in the EEZ of Mexico

Thirty-one species of chondrichthyans were recorded on both oceanic basins of the EEZ of Mexico. Sharks, 27 species, constituted approximately 87% of the total of species followed by four species of rays. Amphi-American chimaeras were not recorded.

Among the sharks, the genus *Carcharhinus* (Carcharhiniformes: Carcharhinidae) is the richest with seven species, the Bull shark *C. leucas* (Müller & Henle, 1839), the Oceanic whitetip shark *C. longimanus* (Poey, 1861), the Blacktip shark *C. limbatus* (Müller & Henle, 1839), the Silky shark *C. falciformis* (Müller & Henle, 1839), the Dusky shark *C. obscurus* (Lesueur, 1818), the Smalltail shark *C. porosus* (Ranzani, 1839) and the Bignose shark *C. altimus* (Springer, 1950). The hammerhead sharks of the Sphyrnidae family with distributions on both coasts are the Scalloped hammerhead *Sphyrna lewini* (Griffith & Smith, 1834), the Great hammerhead *S. mokarran* (Rüppell, 1837), the Bonnethead *S. tiburo* (Linnaeus, 1758) and the Smooth hammerhead *S. zygaena* (Linnaeus, 1758).

In the case of the batomorphs, only four species are distributed on both coasts of the country, the Pelagic stingray *Pteroplatytrygon violacea* (Bonaparte, 1832), the Chupare stingray *Styracura schmardae* (Werner, 1904), the Giant manta *Mobula birostris* (Walbaum, 1792) and the Common sawfish *Pristis pristis* (Linnaeus, 1758). The amphi – American distribution of the species *Ginglymostoma cirratum* Bonnaterre, 1788 (Nurse shark) and *Aetobatus narinari* (Euphrasen, 1790) (the Spotted eagle ray) is not considered in the present study because of the reasons stated in the Discussion.

Discussion

The species richness of Mexican chimaeras, sharks, skates, and rays, when compared to other Latin American countries, is above the 165 species reported for Brazil (Rosa and Gadig 2014); the 117 from Colombia (62 Pacific coast species and 75 for the Caribbean Sea; Navia et al. 2016); the 99 for Costa Rica (12 Caribbean Sea, 75 Pacific, and 12 amphi-American; Bussing and López 1999,2010, Espinoza et al. 2018); the 98 for Venezuela (60 sharks, 37 rays, and one chimera; Cervigón and Alcalá 1999, Tavares and López 2009), and the 38 species for Ecuador (Jiménez-Prado and Beárez 2004, Estupiñan-Montaño et al. 2016). This diversity of Chondrichthyes inhabiting the EEZ of Mexico (approximately 3,000,150 km²) makes it a megadiverse country for this group of species.

The total numbers of chondrichthyans fishes herein reported to the species, genus, family, and order levels in this study are 217, 84, 44, and 13, respectively. These numbers are similar to those reported for Mexico by Del Moral-Flores et al. (2015a) and Del Moral-Flores et al. (2016). However, some of the discrepancies can be attributed to recent taxonomic readjustments, especially for the Division Batomorphi.

Sharks, in general, are the group with the highest diversity of species in the EEZ of Mexico with 111 species (51%). These results are consistent with a previous study by Del Moral-Flores et al. (2016), who reported the same number of species; however, the systematic inventories are not equivalent because the present study does not include records of species reported by those authors as *aff. sp.* or species herein recognized as synonyms (e.g., *Sphyrna vespertina* Springer, 1940; *Negaprion fronto* (Jordan & Gilbert, 1882); *Centrophorus niaukang* Teng, 1959) or subspecies (e.g. *Leucoraja garmani caribbaea* (McEachran, 1977)).

The group of the skates and rays contained 98 species and constituted 45% of the total diversity recorded for the EEZ of Mexico. This figure is very similar to that reported by Del Moral-Flores et al. (2016), who listed 95 species. The species additions to the Mexican chondrichthyans for this group are the Longsnout butterfly ray *Gymnura crebripunctata* (Peters, 1869), the Lessa's butterfly ray *Gymnura lessae* Yokota & De Carvalho, 2017 (see below), the Brazilian cownose ray *Rhinoptera brasiliensis* Müller, 1836, the Fake round ray *Urotrygon simulatrix* Miyake & McEachran, 1988, and the Pacific guitarfish *Pseudobatos planiceps* (Garman, 1880) (the latter species is placed in a new genus, sensu Last et al. 2016c). The eight species of chimaeras did not present any differences among the bibliographic sources and the databases consulted.

Although the Batomorphi constitutes approximately 53% (633) of the total living species of the Chondrichthyes class worldwide, with an addition of at least fifty detected species and yet to be described (Last et al. 2016a), it is not surprising that sharks are the most representative group in the present study. This phenomenon is considered to be related to a historical preference for studying sharks, due to their charismatic characteristics (sensu Isasi-Catalá 2011), and a greater interest in fishing for them, in comparison with the rays and skates. However, currently, the number of scientists referencing batomorpha is increasing, allowing us to consider a scenario where skates

and rays species might be studied with a greater emphasis at national and international levels (see Last et al. 2016e).

The new families identified and restored by Last et al. (2016c) and White and Naylor (2016) (Trygonorrhinidae and Aetobatidae, respectively) are represented within the EEZ of Mexico, with the records of the species of guitarfish *Zapteryx exasperata* and *Z. xyster* (Rhinopristiformes: Trygonorrhinidae) for the coasts of the Pacific Ocean and the Eagle rays *Aetobatus narinari* and *A. laticeps* (Gill, 1865) (Myliobatiformes: Aetobatidae) for the Atlantic and Pacific Oceans, respectively. Recently, the existence of genetic and morphometric variability in a latitudinal gradient for the guitar species of the *Zapteryx* genus has been identified and should be studied in more detail (Castillo-Páez et al. 2017).

According to the information in OBIS (2017) and GBIF (2017), the Reef stingray *Urobatis concentricus* Osburn & Nichols, 1916 (Myliobatiformes: Urotrygonidae) is exclusive to the western Mexican seas of the Mexican EEZ except for three records from Costa Rica. We considered that these latter records are misidentification or a data capture error in the computer platforms previously mentioned. Thus, in the study conducted by Palacios-Salgado et al. (2012), they do not consider this species (or any other batomorph species) as endemic to the Cortez biogeographic province (eastern Pacific Ocean Region, sensu Briggs (1974)). Nevertheless, other authors consider this species as endemic to the western coast of Mexico (Ehemann et al. 2017).

In the case of the Bat ray *Myliobatis californica* (Gill, 1865) (Myliobatiformes: Myliobatidae), a situation similar to that previously reported for *U. concentricus* occurred, because Palacios-Salgado et al. (2012) also make no reference to this species as endemic to the Cortez biogeographic province as there are records outside the study area of the authors. The source of the data consulted (i.e., OBIS 2017) reviewed all the records for this species ($n = 73$) within the continental shelf of Mexico, which may be considered as an endemic species. However, another source of data consulted (i.e., GBIF 2017) presented more than 500 records, with a few dozen registered for the area of the United States of America, and three records for Indonesia, the Maldives and Panama, which could be attributed, again, to misidentification or a capture error within the GBIF platform, which would corroborate this hypothesis with the one record of this species in the Gulf of Mexico.

The Spiny guitarfish *Pseudobatos spinosus* (Günther, 1870) (Rhinopristiformes: Rhinobatidae) had a single record in the databases consulted (i.e., OBIS 2017), which referred to the collection location of the holotype. Excluding the original description and some of the checklist and computer databases references (e.g., Del Moral-Flores et al. 2015a, Del Moral-Flores et al. 2016, Froese and Pauly 2017), there is no other information available for this species. According to F. Del Moral-Flores (UNAM–Campus Iztacala, pers. comm.), the record could be related to an anomalous specimen (i.e., the holotype) of *Pseudobatos* spp., described as *Rhinobatos spinosus* by Günther (1870: 518). Another possibility that has been cited for this record is that it may be a juvenile specimen of another Rhinobatidae species (Compagno 1999: 471–498). Taking into consideration the conditions of the holotype (a dissected and unrecognizable specimen

of 33 cm in length, deposited in the British Museum, BMNH: 1870.6.20.2) and the total absence of records since its diagnosis, the exclusion of this species from the EEZ of Mexico could be considered. Thus, the taxonomic lists and identification guides would avoid, to some extent, the overestimation of the chondrichthyans species from the EEZ of Mexico.

For the Disparate angel Shark (*Squatina heteroptera*) and the Mexican angel shark (*S. mexicana*) (both referred as endemic species from the east coast of Mexico), recently Weigmann (2016) and Vaz and De Carvalho (2018), have treated these species as junior synonyms of *Squatina dumeril* Lesueur, 1818. Despite the fact that those authors came to the same preliminary conclusion, in this manuscript each *Squatina* species mentioned, are retained as valid species. Nevertheless, it is highly recommended to do further investigation to demonstrate its taxonomic validity. Such as the Spiny guitarfish, these two species are marked in the checklist with a question mark (i.e. species whose validity requires further investigation).

At present, there are recent publications that support the separation of a species considered to have an amphi-American distribution, which is the case for the Nurse shark *Ginglymostoma cirratum* Bonnaterre, 1788 and the UNAM's nurse shark *Ginglymostoma unami* Del Moral-Flores, Ramírez-Antonio, Angulo & Pérez-Ponce de León, 2015. The latter species was described from specimens collected in the tropical eastern Pacific Ocean initially identified as *G. cirratum* and recorded as this new species (*G. unami*) and as endemic to this region, excluding the presence of *G. cirratum* for the Pacific Ocean (Del Moral-Flores et al. 2015b).

Another similar case is for the species the Spotted eagle ray *Aetobatus narinari* (Euphrasen, 1790), considered as an amphi-American species (in fact cosmopolitan species). Currently, based on DNA sequences (White and Naylor 2016, Last et al. 2016a), the Pacific eagle ray *Aetobatus laticeps* is recognized as distributed exclusively on the eastern Pacific coast, while its congener *A. narinari* is an inhabitant of the western Atlantic coast including the Gulf of Mexico and the Caribbean Sea. However, Last et al. (2016e) consider “that it is necessary to do more work to distinguish morphologically the two forms”. Currently, the species is cited as an ambiguous synonym for *A. narinari* in Froese and Pauly (2017).

According to the recent morphometric and molecular results obtained by De Carvalho et al. (2016), the subfamily Styracurinae was described and relocated within the family Potamotrygonidae, which are freshwater batomorphs known only in South America until these recent results. The species *Styracura scharmdae* (Werner, 1904) and the Pacific chupare ray *S. pacifica* (Beebe & Tee-Van, 1941) were removed from *Himantura* in Dasyatidae (sensu Last et al. 2016c, De Carvalho et al. 2016) and are thus the only representatives of the potamotrygonids within the EEZ of Mexico; the first species is cited an inhabitant of both coasts, and the second species is restricted to the western basin.

The recent taxonomic relocation of the two species of the genus *Manta*, the Reef manta ray *M. alfredi* (Kreff, 1868) and the Giant oceanic manta ray *M. birostris* (Walbaum, 1792) within the genus *Mobula* (see Last et al. 2016a), and the consideration

of the Spinetail mobula *Mobula japonica* (Müller & Henle, 1841) as synonymous with the Devil fish *M. mobular* (Bonnaterre, 1788), are considered by some specialists as a taxonomic decision subject to discussion (Guy Stevens, The Manta Team: <https://goo.gl/KYbtZO>; com. pers.).

Finally, a recent taxonomic and morphological revision of butterfly rays (Gymnuridae) has limited the distribution of the Smooth butterfly ray *Gymnura micrura* (Bloch & Schneider, 1801) to the southwestern Atlantic and the new species the Lessa's butterfly ray *Gymnura lessae* Yokota & De Carvalho, 2017 occurring in the Gulf of Mexico, north, and central western Atlantic, substituting *G. micrura* records in that area (see Yokota and De Carvalho 2017).

With the increasing use of various techniques and the analysis tools currently available (e.g., molecular sequences, mitogenome analysis, geometric morphometrics), the future of the biological classification of Chondrichthyes may have higher stability, predictability, and robustness (sensu Crisci and López-Armengol 1983). Due to its geographical location, the extension of its patrimonial sea, and the increase in studies on its chondrichthyans, Mexico will undoubtedly continue to contribute to the knowledge for this group of cartilaginous fishes. At present, for the country, there are at least four species of sharks and two batomorphs that need to be formally described, which have been previously mentioned by various authors (e.g., Castro-Aguirre et al. 1996, Castro 2011a, 2011b, Del Moral-Flores et al. 2015a).

As a corollary to the above, recently published works or studies in progress can be cited. Thus, Fields et al. (2016) proposed the existence of populations with possible cryptic speciation among hammerhead sharks (Sphyrnidae) from the Gulf of Mexico and the Caribbean Sea. Another study conducted by Hinojosa-Alvarez et al. (2016) indicates the possibility of a third species of manta ray within the genus *Mobula* (sensu lato *Manta*, see Last et al. 2016a) for the Yucatan Peninsula. Finally, the mitochondrial divergence between the populations of the Cownose ray *Rhinoptera steindachneri* Evermann & Jenkins, 1891, of the Gulf of California (see also Sandoval-Castillo and Rocha-Olivares 2011) is currently being investigated (Christian Jones, NOAA-SFSC Mississippi, pers. comm.), as is the case of *Urotrygon* spp. in the southern Gulf of California by the present authors, what could result in the description of new species in the EEZ of Mexico.

Acknowledgements

The authors thank the BEIFI-IPN and CONACyT Program (NRE and LVGG), COFAA-IPN, EDI-IPN and CONACyT-SNI (JDA and JGCV), for their support. The authors thank the Research and Postgraduate Secretariat of the IPN (SIP-IPN) for its support of the project *ElasmoMex: Information System and Interactive Diagnostic Keys of the Mexican Elasmobranchs* (SIP-IPN 20161015; <https://goo.gl/hB5btX>). The authors are also very grateful for the suggestion made by Simon Weigmann -Elasmo-Lab Elasmobranch Research Laboratory- which helped to improve the understanding of this

article. This paper is in honor of the late José Luis “Doc” Castro-Aguirre (1943–2011), a mainstay of contemporary ichthyology in Mexico. At the time of his death, “Doc” Castro-Aguirre was the co-author of the first and the last species of chondrichthyans described in Mexico by national scientists, i.e., the Mexican hornshark *Heterodontus mexicanus* Taylor & Castro-Aguirre, 1972, and the Whitemargin smoothhound *Mustelus albiginnis* Castro-Aguirre, Antuna-Mendiola, González-Acosta and De La Cruz-Agüero, 2005. AJE (<https://www.aje.com/>) performed the English language editing of the manuscript.

References

- Aceró AP, Tavera JJ, Anguila R, Hernández L (2016) A new southern Caribbean species of angel shark (Chondrichthyes, Squaliformes, Squatinidae), including phylogeny and tempo of diversification of American Species. *Copeia* 104: 577–585. <https://doi.org/10.1643/CI-15-292>
- Briggs JC (1974) Marine zoogeography. McGraw-Hill, New York, 461 pp.
- Bussing WA, López MI (1999) Peces demersales y pelágicos costeros del Pacífico de Centro América meridional. Guía ilustrada. *Revista de Biología Tropical* 47: 1–164. <https://doi.org/10.15517/rbt.v47i2>
- Bussing WA, López MI (2010) Peces costeros del caribe de Centro América meridional. Guía ilustrada. *Revista de Biología Tropical* 58: 1–240. <https://doi.org/10.15517/rbt.v58i0>
- Castillo-Páez A, Sandoval-Castillo J, Corro-Espinoza D, Tovar-Ávila J, Blanco-Parra MDP, Saavedra-Sotelo NC, Sosa-Nishizaki O, Galván-Magaña F, Rocha-Olivares A (2017) Cutting through the Gordian knot: unravelling morphological, molecular, and biogeographical patterns in the genus *Zapteryx* (guitarfish) from the Mexican Pacific. *ICES Journal of Marine Science* 74: 1630–1638. <http://doi.org/10.1093/icesjms/fsx021>.
- Castro JI (2011a) *The Sharks of North America*. Oxford University Press, New York, 640 pp.
- Castro JI (2011b) Resurrection of the name *Carcharhinus cerdale*, a species different from *Carcharhinus porosus*. *Aqua International Journal of Ichthyology* 17: 1–10. <https://aquapress.com/resurrection-of-the-name-carcharhinus-cerdale-a-species-different-from-carcharhinus-porosus-2/>
- Castro-Aguirre JL (1965) Peces sierra, rayas, mantas y especies afines de México. *Anales del Instituto Nacional de Investigaciones Biológico Pesqueras* 1: 169–256.
- Castro-Aguirre JL, Espinosa-Pérez H (1996) Catálogo sistemático de las Rayas y especies afines de México. *Listados Faunísticos de México VII*. Instituto de Biología – UNAM, Ciudad de México, 75 pp.
- Cervigón F, Alcalá A (1999) *Los peces marinos de Venezuela*. Tomo V. Fondo Editorial del Estado Nueva Esparta, Isla de Margarita, 230 pp.
- Compagno LJV (1999) Checklist of living elasmobranchs. In: Hamlett WC (Ed) *Sharks, skates, and rays: the biology of elasmobranch fishes*. John Hopkins University Press, Maryland, 471–498.

- Crisci JV, López-Armengol MF (1983) Introducción a la teoría y práctica de la taxonomía numérica. Monografía científica No. 26: Serie de biología. Secretaría General de la Organización de los Estados Americanos, Programa Regional de Desarrollo Científico y Tecnológico, Washington, D.C., 132 pp.
- De Carvalho MR, Loboda TS, Da Silva JP (2016) A new subfamily, Styracurinae, and new genus, *Styracura*, for *Himantura schmardae* (Werner, 1904) and *Himantura pacifica* (Beebe & Tee-Van, 1941) (Chondrichthyes: Myliobatiformes). *Zootaxa* 4175: 201–221. <http://dx.doi.org/10.11646/zootaxa.4175.3.1>
- Del Moral-Flores LF, Pérez-Ponce de León G (2013) Tiburones, rayas y quimeras de México. *Biodiversitas* 111: 1–6. <http://www.biodiversidad.gob.mx/Biodiversitas/Articulos/biodiv111art1.pdf>
- Del Moral-Flores LF, Morrone JJ, Alcocer Durand J, Espinosa-Pérez H, Pérez-Ponce de León G (2015a) Lista patrón de los tiburones, rayas y quimeras (Chondrichthyes, Elasmobranchii, Holocephali) de México. *Arxius de Miscellània Zoològica* 13: 47–163. 163. http://amz.museucienciasjournals.cat/files/AMZ_vol_13_2015_pp_47-163_Del-Moral-Flores-et-al.pdf
- Del Moral-Flores LF, Ramírez-Antonio E, Angulo A, Pérez-Ponce de León G (2015b) *Ginglymostoma unami* sp. nov. (Chondrichthyes: Orectolobiformes: Ginglymostomatidae): una especie nueva de tiburón gata del Pacífico oriental tropical. *Revista Mexicana de Biodiversidad* 86: 48–58. <https://doi.org/10.7550/rmb.46192>
- Del Moral-Flores LF, Morrone JJ, Alcocer Durand J, Ponce de León G (2016) Diversidad y afinidades biogeográficas de los tiburones, rayas y quimeras (Chondrichthyes: Elasmobranchii, Holocephali) de México. *Revista de Biología Tropical* 64: 1469–1486. <https://doi.org/10.15517/rbt.v64i4.22774>
- DOF (1976) Diario Oficial de la Federación. Estados Unidos Mexicanos. México, D.F., February 6, 1976. Available at <http://www.dof.gob.mx/>
- Ebert DA, Fowler S, Compagno LVJ, Dando M (2013) *Sharks of the World: A Fully Illustrated Guide*. Wild Nature Press, Plymouth, 528 pp.
- Espinosa-Pérez H, Castro-Aguirre JL, Huidobro-Campos L (2004) Catálogo sistemático de Tiburones (Elasmobranchii: Selachimorpha). Listados Faunísticos de México IX. Instituto de Biología UNAM, Ciudad de México, 135 pp.
- EOL (2017) The Encyclopedia of Life. <http://eol.org/> [Reviewed: November 15, 2017]
- Ehemann NR, Perez-Palafox XA, Pabón-Aldana K, Mejía-Falla PA, Navia AF, Cruz-Escalona VH (2017) Biological notes on the reef stingray, *Urobatis concentricus*, an endemic species of Mexico. *Journal of Fish Biology* 91: 1228–1235. <http://doi.org/10.1111/jfb.13398>
- Espinoza M, Díaz E, Angulo A, Hernández S, Clarke TM (2018) Chondrichthyan diversity, conservation status, and management challenges in Costa Rica. *Frontiers in Marine Science* 5: article 85. <http://doi: 10.3389/fmars.2018.00085>
- Estupiñan-Montaño C, Galván-Magaña F, Hacothen-Domené A, Estupiñan-Ortíz JF (2016) First reports of smalltooth sand tiger sharks, *Odontaspis ferox* (Elasmobranchii: Lamniformes: Odontaspidae), off continental Ecuador. *Acta Ichthyologica et Piscatoria* 46: 251–253. <http://doi.org/10.3750/AIP2016.46.3.09>

- Fields AT, Feldheim KA, Gelsleichter J, Pfoertner C, Chapman DD (2016) Population structure and cryptic speciation in bonnethead sharks *Sphyrna tiburo* in the south-eastern USA and Caribbean. *Journal of Fish Biology* 89: 2219–2233. <https://doi.org/10.1111/jfb.13025>
- Froese R, Pauly D (2017) FishBase. World Wide Web Electronic Publication. <http://www.fishbase.org> [Reviewed: November 15, 2017]
- Garman SW (1913) The Plagiostomia (Sharks, Skates, and Rays). *Memoirs of the Museum of Comparative Zoology at Harvard College* 36: 1–528.
- GBIF (2017) Global Biodiversity Information Facility. Available at [<http://www.gbif.org/>] [Reviewed: November 15, 2017]
- Hinojosa-Alvarez S, Walter RP, Díaz-Jaimes P, Galván-Magaña F, Paig-Tran EM (2016) A potential third Manta Ray species near the Yucatán Peninsula? Evidence for a recently diverged and novel genetic *Manta* group from the Gulf of Mexico. *PeerJ* 4 e2586. <https://doi.org/10.7717/peerj.2586>
- Hoyos-Padilla M, Papastamatiou YP, O’Sullivan J, Lowe CG (2013) Observation of an attack by a Cookiecutter Shark (*Isistius brasiliensis*) on a White Shark (*Carcharodon carcharias*) *Pacific Science* 67: 129–134. <https://doi.org/10.2984/67.1.10>
- IHO (1953) Limits of Oceans and Seas, 3a Ed Monte-Carlo: International Hydrographic Organization. <https://epic.awi.de/29772/1/IHO1953a.pdf>
- Isasi-Catalá E (2011) Los conceptos de especies indicadoras, paraguas, banderas y claves: su uso y abuso en ecología de la conservación. *Interiencia* 36: 31–38. <https://www.interiencia.net/wp-content/uploads/2018/01/031-ISASI-8.pdf>
- iSpecies (2017) iSpecies. A simple mashup of species information. <http://ispecies.org> [Reviewed: June 22, 2017]
- Jiménez-Prado P, Béarez P (2004) Peces marinos del Ecuador continental II. SIMBIOE, Quito, 406 pp.
- Last PR, Naylor GJ, Séret B, White W, Stehmann M, De Carvalho MR (2016a) Rays of the World. Cornell University Press, Ithaca and London, 790 pp.
- Last PR, Naylor GJ, Manjaji-Matsumoto BM (2016b) A revised classification of the family Dasyatidae (Chondrichthyes: Myliobatiformes) based on new morphological and molecular insights. *Zootaxa* 4139: 345–368. <http://dx.doi.org/10.11646/zootaxa.4139.3.2>
- Last PR, Seret B, Naylor GJ (2016c) A new species of guitarfish, *Rhinobatos borneensis* sp. nov. with a redefinition of the family-level classification in the order Rhinopristiformes (Chondrichthyes: Batoidea). *Zootaxa* 4117: 451–475. <http://dx.doi.org/10.11646/zootaxa.4117.4.1>
- Last PR, Weigmann S, Yang L (2016d) Changes to the nomenclature of the skates (Chondrichthyes: Rajiformes). Supplementary Information. In: Last PR, Yearsley GK (Eds) Rays of the World. CSIRO Special Publication, 11–34.
- Last PR, Weigmann S, Yang L (2016e) The rays of the World Project – an explanation of nomenclatural decisions. Supplementary Information. In: Last PR, Yearsley GK (Eds) Rays of the World. CSIRO Special Publication, 1–10.
- Navia AF, Mejía-Falla PA, Hleap JS (2016) Zoogeography of Elasmobranchs in the Colombian Pacific Ocean and the Caribbean Sea. *Neotropical Ichthyology* 14: e140134. <http://doi.org/10.1590/1982-0224-20140134>
- Naylor GJ, Caira JN, Jensen K, Rosana K, White W, Last PR (2012) A DNA sequence-based approach to the identification of shark and ray species and its implications for global elas-

- mobranch diversity and parasitology. *Bulletin of the American Museum of Natural History* 367: 1–263. <http://hdl.handle.net/2246/6183>
- Nelson JS, Starnes WC, Warren ML (2002) A Capital case for common names of species of fishes – A white crappie or a White Crappie. *Fisheries* 27: 31–33. https://www.srs.fs.usda.gov/pubs/ja/ja_nelson003.pdf
- OBIS (2017) Ocean Biogeographic Information System. <http://www.iobis.org> [Reviewed: November 15, 2017]
- Palacios-Salgado DS, Burnes-Romo LA, Tavera JJ, Ramírez-Valdez A (2012) Endemic Fishes of the Cortez Biogeographic Province (Eastern Pacific Ocean). *Acta Ichthyologica et Piscatoria* 42: 153–164. <http://doi.org/10.3750/AIP2011.42.3.01>
- Rosa RS, Gadig OB (2014) Conhecimento da diversidade dos *Chondrichthyes* marinhos no Brasil: a contribuição de José Lima de Figueiredo. *Arquivos de Zoologia Museu e Zoologia da Universidade e São Paulo* 45: 89–104. <http://www.revistas.usp.br/azmz/article/view/88075>
- Ruiz-Campos G, Castro-Aguirre JL, Balart EF, Campos-Dávila L, Vélez-Marín R (2010) New specimens and records of chondrichthyan fishes (Vertebrata: Chondrichthyes) off the Mexican Pacific coast. *Revista Mexicana de Biodiversidad* 81: 363–371. <http://dx.doi.org/10.22201/ib.20078706e.2010.002.259>
- Sandoval-Castillo J, Rocha-Olivares A (2011) Deep mitochondrial divergence in Baja California Populations of an Aquilopelagic Elasmobranch: The golden Cownose Ray. *Journal of Heredity* 102: 269–274. <http://doi.org/10.1093/jhered/esr004>
- Siqueiros-Beltrones D, De La Cruz-Agüero J (2004) Examen filosófico de las listas sistemáticas como informes científicos y porqué deben ser publicados. *Oceanides* 19: 1–9. <http://oceanides.ipn.mx/index.php/cicimaroceanides/article/view/13/8>
- Tavares R, López D (2009) Tendencias de la producción pesquera de elasmobranchios en Venezuela: con énfasis en los tiburones. *Proceedings of the 62nd Gulf and Caribbean Fisheries Institute* 178–183. <http://nsgl.gso.uri.edu/flsgp/flsgpc09001/data/papers/025.pdf>
- Villalobos E, Martínez A, Lambarri C, Espinosa H (2016) New record of *Zameus squamulosus* (Chondrichthyes: Squaliformes: Somniosidae) in the Southern Gulf of Mexico. *Marine Biodiversity Records* 9: 61. <http://doi.org/10.1186/s41200-016-0047-0>
- Weigmann S (2016) Annotated checklist of the living sharks, batoids, and chimaeras (Chondrichthyes) of the world, with a focus on biogeographical diversity. *Journal of Fish Biology* 88: 837–1037. <http://doi.org/10.1111/jfb.12874>
- Weigmann S (2017) Reply to Borsa (2017): Comment on Annotated checklist of the living sharks, batoids, and chimaeras (Chondrichthyes) of the world, with a focus on biogeographical diversity by Weigmann (2016). *Journal of Fish Biology* 90: 1176–1181. <http://doi.org/10.1111/jfb.13234>
- White WT, Naylor GJ (2016) Resurrection of the family Aetobatidae (Myliobatiformes) for the pelagic eagle rays, genus *Aetobatus*. *Zootaxa* 4139: 435–438. <http://doi.org/10.11646/zootaxa.4139.3.10>
- Yokota L, De Carvalho MR (2017) Taxonomic and morphological revision of butterfly rays of the *Gymnura micrura* (Bloch & Schneider 1801) species complex, with the description of two new species (Myliobatiformes: Gymnuridae). *Zootaxa* 4332: 1–74. <https://doi.org/10.11646/zootaxa.4332.1.1>