

Ephemeroptera of Canada

Luke M. Jacobus¹

¹ *Division of Science, Indiana University Purdue University Columbus, 4601 Central Avenue, Columbus, Indiana 47203, USA*

Corresponding author: *Luke M. Jacobus* (luke.jacobus@gmail.com)

Academic editor: *D. Langor* | Received 3 May 2018 | Accepted 9 July 2018 | Published 24 January 2019

<http://zoobank.org/BDB88C5A-F09C-4F36-B55E-796DAF35B6B4>

Citation: Jacobus LM (2019) Ephemeroptera of Canada. In: Langor DW, Sheffield CS (Eds) *The Biota of Canada – A Biodiversity Assessment. Part 1: The Terrestrial Arthropods*. ZooKeys 819: 211–225. <https://doi.org/10.3897/zookeys.819.26411>

Abstract

Thus far, 335 currently valid species in 82 genera and 21 families of mayflies (Ephemeroptera) have been documented from Canada, remarkably representing a little more than half of the combined species richness of Canada, Mexico and the USA. The current known species richness for Canada represents an increase of 11.3% as compared to that reported in 1979. Species richness is greatest in the families Heptageniidae (83), Baetidae (76) and Ephemerellidae (45). A total of 328 DNA Barcode Index Numbers (BINs) are available for Canadian mayfly species. The greatest net gains anticipated for future species tallies are for Baetidae (25), Heptageniidae (10) and Leptophlebiidae (10). A total of 66 more species overall is anticipated for Canada, with greatest gains potentially coming from lentic habitats across Canada and from far eastern and far western areas in general. However, even metropolitan areas should not be overlooked for the potential of discovery.

Keywords

aquatic insects, biodiversity assessment, Biota of Canada, Ephemeroptera, mayflies

“Great strides” have been made in our understanding of the Canadian mayfly fauna since Lehmkuhl (1979) used this phrase in his review of the status of this order as part of the larger work by Danks (1979). At the time, Lehmkuhl (1979) indicated that there was “no comprehensive treatment of the Canadian fauna.” That situation has since been remedied with a continually updated Canada species list that has

been available online since 1995 (McCafferty 1996, McCafferty and Jacobus 2018). Ecological and biological information was summarized by Waltz and Burian (2008) and others in previous editions of that work. Peters (1988) provided a biogeographic discussion of the Canada fauna that remains largely applicable today. McCafferty et al. (1990) discussed the status and needs of mayfly systematics in North America north of Mexico, including Canada. A history of mayfly science in Canada was summarized by McCafferty and Randolph (1998) and further reviewed by McCafferty (2001).

Thus far, 335 currently valid species in 82 genera and 21 families have been documented from Canada, remarkably representing a little more than half the combined species richness of Canada, Mexico and the USA (McCafferty and Jacobus 2018). Two species described from Canada are considered *nomen dubia*, and thus, they are not included in species counts (McCafferty and Bae 1992, Jacobus and McCafferty 2007). At least 15 of the species documented from Canada have a Holarctic distribution (e.g., Kluge 1980, Novikova and Kluge 1987, Kluge and Novikova 2011, Bauernfeind and Soldán 2012, Kjærstad et al. 2012, Savolainen et al. 2014, Cordero et al. 2016). Eight of the Canadian species have geographic distributional ranges that extend south to Central America (McCafferty and Jacobus 2018). Ten of the Canadian species are not yet known outside Canada, but the number of truly endemic species probably is much lower, because most of these ten have been collected near the southern border and likely occur in the United States also (Jacobus 2018). Low endemism in Canada is not surprising, considering the recent glacial history of Canada and therefore the large degree to which the current fauna reflects dispersal and post-glaciation recolonization events (Peters 1988, McCafferty and Randolph 1998). No non-native species are known from Canada (Randolph et al. 2002).

The current known species richness for Canada represents an increase of 11.3% as compared to that reported in 1979. The families with the largest net increase in species numbers are Heptageniidae (10), Caenidae (6) and Baetidae (5). Ameletidae probably has nearly as great an increase as Heptageniidae (Zloty 1996, Zloty and Harper 1999), but current and 1979 species numbers could not be compared due to the way data were classified in the 1979 report (see footnote 7 in Table 1).

The current family and genus classifications used here for the Ephemeroptera of Canada are quite different from those that were hypothesized nearly 40 years ago, as reflected in the work of Lehmkuhl (1979). Current family-level classifications in North America, and Canada in particular, primarily reflect the work of McCafferty (1991b), with subsequent revisions or significant review by Landa and Soldán (1985), Peters and Peters (1993), Wang and McCafferty (1995), and McCafferty (2004). Higher classification remains tentative and contentious in many cases, as reviewed by Ogden et al. (2009), and the higher classification followed here borrows from various sources (see footnote 1 in Table 1) to best reflect the most recent phylogenetic hypotheses (Jacobus and McCafferty 2006, Ogden et al. 2009: fig. 7, Miller et al. 2018).

The classification of species into genera also has changed markedly in the last 40 years. The family Baetidae has seen the greatest number of changes, with multiple new genera described, several more recent generic revisions, and some species going

back and forth between genera several times. McCafferty and Waltz (1990) provided a summary of these changes up to that point, along with some of their own, and that work marks the starting point for Baetidae references cited in Table 1; the reference list is long but not exhaustive, providing a strong starting place for understanding the current generic classification applied to the species found in Canada. Heptageniidae also has seen considerable changes in generic classification of species, and most of the current genus-level systematics for Canada are reflected in Webb and McCafferty (2008). Sun and McCafferty (2008) made significant changes to genera of the subfamily Brachycercinae in the family Caenidae, and Jacobus and McCafferty (2008) most recently revised the generic classification of Ephemerellidae. The classification of species into genera across the order remains inconsistent and somewhat contested globally (e.g., Bauernfeind and Soldán 2012, Jacobus 2015, 2016).

The number of new species described and named from Canada has been relatively few since 1979, but the most notable gain has been in the genus *Ameletus* Eaton (Ameletidae) (Zloty 1996, Zloty and Harper 1999). In contrast to new species descriptions, a remarkable number of species synonyms have been proposed for the Canadian fauna, and these are reflected in the complete species synonymies given by McCafferty and Jacobus (2018). However, it should be noted that new evidence, especially from DNA barcoding, challenges many of these concepts of highly variable species, and some of the current concepts of single species may be split into multiple species after more research is completed (Webb et al. 2012). Recently collected mayfly specimens from Canada have played an important role in the generation of regional DNA barcode libraries (Ball et al. 2005, Zhou et al. 2009, 2010, Webb et al. 2012) and discovering trans-Atlantic species distributions (e.g., Kjærstad et al. 2012, Savolainen et al. 2014, Cordero et al. 2016).

While a list of species for Canada that reflects current species concepts has been maintained online for over 20 years (see above), only coarse geographic distributions have been indicated there, with Canada divided into eastern, western, and far northern regions, essentially following those indicated by Edmunds et al. (1976: 50) and McCafferty and Waltz (1990: fig. 1). McCafferty and Randolph (1998), however, provided a comprehensive listing of species for each of the Canadian territories and provinces at that time. Subsequent notable contributions for each include the following for Alberta (Webb and McCafferty 2003, McCafferty et al. 2004, 2012, McCafferty 2009, Webb et al. 2012), British Columbia (Jacobus and McCafferty 2001, McCafferty et al. 2012), Manitoba (Flanagan et al. 2001, Jacobus and McCafferty 2001, McCafferty et al. 2004, McCafferty 2009, Zhou et al. 2009, 2010, Webb et al. 2012, Kjærstad et al. 2012), New Brunswick (Jacobus and McCafferty 2001, McCafferty 2009, Webb et al. 2012, Burian 2017), Newfoundland and Labrador (Lomond and Colbo 2000, McCafferty 2009, 2011b, Webb et al. 2012, Burian 2017), the Northwest Territories (Randolph and McCafferty 2001, Bowman et al. 2010, Gorski et al. 2015, Burian 2017, Giberson and Burian 2017), Nova Scotia (Jacobus and McCafferty 2001, McCafferty et al. 2004), Nunavut (Randolph and McCafferty 2001, Giberson et al. 2007, McCafferty 2011a, Burian 2017), Ontario (McCafferty et al. 2008, McCafferty 2009, Webb et

Table 1. Census of Ephemeroptera in Canada.

Taxon ¹	No. species reported in Lehmkuhl (1979)	No. species currently known from Canada ²	No. BINs ³ available for Canadian species	Est. no. undescribed or unrecorded species in Canada	General distribution by ecozone ⁴	Information sources
Suborder Pisciforma⁵						
Superfamily Baetoidea						
Baetidae	71	76	87	25	most ecozones	McCafferty and Waltz 1990, Lago-Ortiz and McCafferty 1998, Randolph et al. 2002, Jacobus and McCafferty 2005, McCafferty et al. 2005, 2010, Guenther and McCafferty 2008, McCafferty 2011a, Burian and Myers 2011, Jacobus and Wiersema 2014, Cruz et al. 2017, Webb et al. 2018
Superfamily Siphonuroidea						
Acanthametropodidae ⁶	1	1	0	1	Boreal Plains, Prairies	Lehmkuhl 1976, McCafferty 1991a
Ameletidae ⁷	2	23	18	5	nearly all ecozones	Zloty and Harper 1999
Ametropodidae	2	2	1	0	central & western Boreal & Taiga ecozones	Jacobus 2013
Metropodidae	3	3	3	1	most ecozones	Berner 1978, Engblom et al. 1993, Kluge 1996
Siphonuridae ⁷	35	20	12	1	nearly all ecozones	Traver 1935, Kondratieff and Voshell 1981, Provonsha and McCafferty 1982, McCafferty and Edmunds 1997, Newell and Anderson 2009, Burian 2017
Suborder Setisura⁵						
Superfamily Oligoneuroidea⁵						
Isoneuridae ⁸	5	6	4	1	most ecozones south of taiga	Kondratieff and Voshell 1984
Oligoneuridae	1	1	0	1	Prairies	Edmunds 1951
Superfamily Heptagenioidea						
Arthropleidae ⁹	1	1	0	0	Taiga Shield, Boreal Shield, Hudson Plains, Atlantic Maritime, eastern Boreal Plains	Ide 1930, Wang and McCafferty 1995
Heptageniidae ¹⁰	73	83	82	10	nearly all ecozones	Burian et al. 2008, Webb and McCafferty 2008, 2011

Taxon ¹	No. species reported in Lehmkuhl (1979)	No. species currently known from Canada ²	No. BINs ³ available for Canadian species	Est. no. undescribed or unrecorded species in Canada	General distribution by ecoregion ⁴	Information sources
Pseudironidae ¹¹	1	1	0	0	Boreal Plains, Prairies, western Boreal Shield	Pescador 1985
Suborder Carapacea						
Baetiscidae	7	6	2	0	most ecoregions south of Taiga Plains and Taiga Shield	Pescador and Berner 1981, Baumann and Kondratieff 2000, Webb and McCafferty 2003
Suborder Furcatengalia						
Infraorder Lanceolata						
Superfamily Leptophleboidea						
Leptophlebiidae	24	27	34	8	nearly all ecoregions	Traver 1935, Allen 1973, McCafferty 1992, Burian 2000, Tiunova and Kluge 2016, McCafferty et al. 2017
Infraorder Scaphodonta						
Superfamily Potamanthoidea						
Potamanthidae	3	4	0	0	Atlantic Maritime, Mixedwood Plains, extreme southern Boreal Shield	Bae and McCafferty 1991
Superfamily Ephemerioidea						
Ephemeridae ¹²	11	9	8	2	most ecoregions south of Arctic	McCafferty 1975, 1994
Palingeniidae ¹³	1	1	0	0	southern Boreal Shield, eastern Prairies	McCafferty 2004
Polymitarcyidae	3	3	4	0	Prairies, Boreal Plains, Boreal Shield, Mixedwood Plains, Atlantic Maritime	McCafferty 1975, 1994, Molineri 2010
Infraorder Pannota						
Superfamily Caenoidea						
Caenidae	10	16	15	4	nearly all ecoregions	Provonsha 1990, Sun and McCafferty 2008

Taxon ¹	No. species reported in Lehmkuhl (1979)	No. species currently known from Canada ²	No. BINs ³ available for Canadian species	Est. no. undescribed or unrecorded species in Canada	General distribution by ecoregion ⁴	Information sources
Neophemeridae	1	1	0	0	Mixedwood Plains	Schmude et al. 2012
Superfamily Ephemeroidea						
Ephemeroidea	44	45	50	5	nearly all ecoregions	Jacobus and McCaffery 2008, Jacobus 2010
Leptophlebiidae ^{4a}	4	6	8	2	most ecoregions, except for Arctic, Taiga Cordillera, Taiga Plains, and Boreal Cordillera	Alba-Tercedor and Flannagan 1995, Wiersma and McCaffery 2000, 2005, Baumgardner et al. 2003, Baumgardner 2009
Total	301	335	328	66		

¹Mayfly higher classification is far from resolved. This classification is a synthesis based on Sun et al. (2006), Ogden et al. (2008, 2009; Figs 1, 7), Prisecaru et al. (2014), Miller et al. (2018). ²Based on Mayfly Central species list for North America (McCaffery 1996, McCaffery and Jacobus 2018). ³Barcode Index Number, as defined in Ramasingham and Hebert (2013). Only those BINs that include Canada specimens are included in counts. Counts accurate as of 30 April 2018. ⁴See figure 1 in Langor (2019) for a map of ecoregions. ⁵Taxon known to be paraphyletic, even when considering only North American fauna, north of Mexico. ⁶Recorded in 1979 as Siphonuridae, subfamily Acanthametropodinae. ⁷Due to changes in family-level classification, the 1979 report of species in the subfamily Siphonurinae now applies to species in the current families Ameletidae and Siphonuridae. For simplicity, the pooled 1979 total of 35 species has been inserted for Siphonuridae and nothing has been entered for Ameletidae. ⁸Recorded in 1979 as Siphonuridae, subfamily Isonychiinae. ⁹Recorded in 1979 as Heptageniidae, subfamily Arthropleinae. ¹⁰Today this family excludes taxa that were reported as Arthropleinae and Pseudironinae in 1979. Thus, the number of species reported herein for 1979 includes only those reported for Heptageniinae and Anepeorinae. ¹¹Recorded in 1979 as Heptageniidae, subfamily Pseudironinae. ¹²Recorded in 1979 as Ephemeroidea, subfamily Ephemeroidea. ¹³Recorded in 1979 as Ephemeroidea, subfamily Pentageniinae. ¹⁴Recorded in 1979 as Tricorythidae.

al. 2012, Klubertanz 2016b), Prince Edward Island (Jacobus and McCafferty 2001), Quebec (Jacobus and McCafferty 2001, Randolph et al. 2002, Burian 2017), Saskatchewan (McCafferty et al. 2004, Webb et al. 2004, 2007, McCafferty 2009, Miyazaki and Lehmkuhl 2011, Webb et al. 2012), and Yukon Territory (McCafferty et al. 2004, Giberson and Burian 2017). Other scattered reports of new provincial and territorial records can be found in a variety of sources, including those listed in Table 1. The general distributions detailed in Table 1 are based on Canadian ecozones, as are the general distributions given for other taxa in this series of papers. Published records of species, as reviewed above, were used to determine the ecozones inhabited by each family.

Beyond those already reported in the scientific literature, an additional 66 species of mayflies are expected to be added to the list of Canadian species (Table 1). This tally includes named species known to occur in adjacent parts of the USA that will be found predictably in adjoining Canada (L Jacobus unpubl. data, SK Burian pers. comm., JM Webb pers. comm.), and it includes known, but unnamed, species likely new to science, especially from the families Ameletidae, Baetidae, Caenidae (Brachycercinae), Ephemerellidae and Heptageniidae (Jacobus et al. unpubl. data). For most of these unnamed species, more material is needed to allow a final decision and to facilitate descriptions; others simply are awaiting completion of formal scientific descriptions. A couple of southeastern United States species likely to occur in Canada have been diagnosed tentatively, however, and given informal designations by McCafferty et al. (2017).

Lehmkuhl (1979) indicated that very few descriptions of larvae were “adequate for taxonomic purposes. Therefore the state of knowledge...[was]...not tabulated” at that time. Major advances in our knowledge of the larval stages have occurred since then (McCafferty et al. 1990), and in fact the larva is now the most commonly and widely studied metamorphic stage of mayflies. Morphological identifications of larvae of most species from Canada can be made using two recent workshop manuals (Jacobus et al. 2014, Jacobus 2017), both available by request, and the works cited therein. Jacobus (2017) relies heavily on integrating two recently published identification keys (Klubertanz 2016a, McCafferty et al. 2017). Specimens collected from Manitoba must be identified with all these sources cautiously, given the mix of eastern and western species in the province. Although the knowledge of larvae is generally good, certain taxa still require better description, and a variety of cryptic species complexes require resolution (Webb et al. 2012, 2018). Morphological identifications of mayfly subimagos and female adults to the species level are usually very difficult to impossible at the present, and morphological identification of male adults is possible only via the use of a variety of published literature and unpublished research notes, some of which remains unavailable outside private libraries.

Looking to the future, additional work is needed on lentic habitats throughout Canada because unusual new taxa continue to be discovered from lakes throughout North America (Hill et al. 2010, McCafferty 2011a). Northern Canada in general requires much more work before its fauna will be documented adequately (McCafferty et al. 1990, Cordero et al. 2016, Giberson and Burian 2017), but even less remote

areas continue to yield new records of species (e.g., Klubertanz 2016b). Provinces such as Newfoundland and British Columbia have less documented species richness than might be expected, based on richness of surrounding areas (McCafferty and Randolph 1998, Meyer and McCafferty 2007, McCafferty et al. 2012) and may provide opportunities for considerable discovery that will add to species numbers, including new country records for Canada. Although our knowledge of the larval stage of mayflies has improved drastically since Lehmkuhl's (1979) assessment, association of larvae with male adults should remain a priority, even in geographic areas that are considered easily accessible and mostly well known. The Ottawa and Montreal regions, for example, contain type localities for species still unknown as larvae (e.g., McDunnough 1925).

Acknowledgments

David Langor and Cory Sheffield helped to improve various stages of the manuscript. Steve Burian, Donna Giberson, Jason Robinson and Jeff Webb provided valuable discussion. Donna Giberson and Jason Robinson suggested revisions that improved a late version of the manuscript. Jeremy deWaard confirmed and updated some Barcode Index Number data.

References

- Alba-Tercedor J, Flannagan J (1995) Two new Canadian species of the genus *Tricorythodes* Ulmer, with additional studies on other North American species (Insecta, Ephemeroptera: Leptohyphidae). *Canadian Journal of Zoology* 73: 1588–1598. <https://doi.org/10.1139/z95-189>
- Allen RK (1973) Generic revisions of mayfly nymphs. I. *Traverella* in North and Central America (Leptophlebiidae). *Annals of the Entomological Society of America* 66: 1287–1295. <https://doi.org/10.1093/aesa/66.6.1287>
- Bae YJ, McCafferty WP (1991) Phylogenetic systematics of the Potamanthidae (Ephemeroptera). *Transactions of the American Entomological Society* 117: 1–143.
- Ball SL, Hebert PDN, Burian SK, Webb JM (2005) Biological identifications of mayflies (Ephemeroptera) using DNA barcodes. *Journal of the North American Benthological Society* 24: 508–524. <https://doi.org/10.1899/04-142.1>
- Bauernfeind E, Soldán T (2012) *The Mayflies of Europe*. Apollo Books, Ollerup, 781 pp.
- Baumann R, Kondratieff BC (2000) A confirmed record of the Ephemeroptera genus *Baetisca* from west of the Continental Divide and an annotated list of the mayflies of the Humboldt River, Nevada. *Western North American Naturalist* 60: 459–461.
- Baumgardner DE (2009) *Tricorythus minutus* Traver, a new synonym of *Tricorythodes explicatus* Eaton (Ephemeroptera: Leptohyphidae). *Proceedings of the Entomological Society of Washington* 111: 57–67. <https://doi.org/10.4289/07-083.1>
- Baumgardner DE, Burian SK, Bass D (2003) Life stage descriptions, taxonomic notes, and new records for the mayfly family Leptohyphidae (Ephemeroptera). *Zootaxa* 332: 1–12.

- Berner L (1978) A review of the family Metretopodidae. *Transactions of the American Entomological Society* 104: 91–137.
- Bowman M, Spencer P, Dubé M, West D (2010) Regional reference variation provides ecologically meaningful protection criteria for northern world heritage site. *Integrated Environmental Assessment and Management* 6: 12–27.
- Burian SK (2000) A revision of the genus *Leptophlebia* Westwood in North America (Ephemeroptera: Leptophlebiidae: Leptophlebiinae). *Ohio Biological Survey, New Series* 13: 1–80.
- Burian SK (2017) The taxonomy of the North American species of *Parameletus* Bengtsson, 1908 (Ephemeroptera: Siphonuridae), with keys to nymphs and male imagos. *Canadian Journal of Arthropod Identification* 33. <https://doi.org/10.3752/cjai.2017.33>
- Burian SK, Myers LW (2011) A new species of *Acentrella* Bengtsson (Ephemeroptera: Baetidae) from New York and New England (USA), redescription of the nymph of *A. parvula* (McDunnough), and key to known adult males of Nearctic *Acentrella*. *Aquatic Insects* 33: 305–334. <https://doi.org/10.1080/01650424.2011.639305>
- Burian SK, Swartz BI, Wick PC (2008) Taxonomy of *Epeorus frisoni* (Burks) and a key to New England species of *Epeorus*. In: Hauer FR, Stanford JA, Newell RL (Eds) *International Advances in the Ecology, Zoogeography and Systematics of Mayflies and Stoneflies*. University of California Publications in Entomology 128, 277–294.
- Cordero RD, Sánchez-Ramírez S, Currie DC (2016) DNA barcoding of aquatic insects reveals unforeseen diversity and recurrent population divergence patterns through broad-scale sampling in northern Canada. *Polar Biology* 40: 1687–1695. <https://doi.org/10.1007/s00300-016-2062-3>
- Cruz PV, Usher AK, Jacobus LM (2017) Review of continental North and Central American *Paracloeodes* Day 1955 (Ephemeroptera: Baetidae), including description of a new species from Minnesota, U.S.A. *Zootaxa* 4263: 333–346.
- Danks HV (1979) Canada and its insect Fauna. *Memoirs of the Entomological Society of Canada* No. 108, 573 pp.
- Eaton AE (1883–1888) A revisional monograph of recent Ephemeridae or mayflies. *Transactions of the Linnean Society of London* 3(2): 1–352.
- Edmunds Jr GF (1951) New species of Utah mayflies. I. Oligoneuriidae (Ephemeroptera). *Proceedings of the Entomological Society of Washington* 53: 327–331.
- Edmunds Jr GF, Jensen SL, Berner L (1976) *The Mayflies of North and Central America*. University of Minnesota Press, Minneapolis, 330 pp.
- Engblom E, Lingdell P-E, Nilsson AN, Savolainen E (1993) The genus *Metretopus* (Ephemeroptera, Siphonuridae) in Fennoscandia — identification, faunistics and natural history. *Entomologica Fennica* 4: 213–222.
- Flannagan JF, Alba-Tercedor J, Lowen RG, Cobb DG (2001) Emergence of Ephemeroptera from the Assiniboine River, Canada. In: Dominguez E (Ed.) *Trends in Research in Ephemeroptera and Plecoptera*. Springer, New York, 97–106. https://doi.org/10.1007/978-1-4615-1257-8_13
- Giberson DJ, Burian SK (2017) How valid are old species lists? How archived samples can be used to update Ephemeroptera biodiversity information for northern Canada. *The Canadian Entomologist* 149: 755–773. <https://doi.org/10.4039/tce.2017.27>

- Giberson DJ, Burian SK, Shouldice M (2007) Life history of the northern mayfly *Baetis bundyae* in Rankin Inlet, Nunavut, Canada, with updates to the list of mayflies of Nunavut. *The Canadian Entomologist* 139: 628–642. <https://doi.org/10.4039/n06-089>
- Gorski MR, Fox ADE, McQueen JI, Jacobus LM (2015) *Plauditus cestus* (Provonsha & McCafferty, 1982) (Insecta: Ephemeroptera: Baetidae): New records from Virginia and the Northwest Territories, with notes on color variation. *Check List* 11(1) 1498: 1–3. <https://doi.org/10.15560/11.1.1498>
- Guenther JL, McCafferty WP (2008) Mayflies (Ephemeroptera) of the Great Plains. IV: South Dakota. *Transactions of the American Entomological Society* 134: 147–171. [https://doi.org/10.3157/0002-8320\(2008\)134\[147:MEOTGP\]2.0.CO;2](https://doi.org/10.3157/0002-8320(2008)134[147:MEOTGP]2.0.CO;2)
- Hill MA, Pfeiffer J, Jacobus LM (2010) A new genus and new species of Baetidae (Ephemeroptera) from lakes and reservoirs in eastern North America. *Zootaxa* 2481: 61–68.
- Ide FP (1930) The nymph of the mayfly genus *Cinygma* Eaton. *The Canadian Entomologist* 62: 42–45. <https://doi.org/10.4039/Ent6242-2>
- Jacobus LM (2010) Taxonomic review of the *Caudatella heterocaudata* (McDunnough) and *C. hystrix* (Traver) complexes (Insecta: Ephemeroptera: Ephemerellidae). *Psyche* 2010 (493935). <https://doi.org/10.1155/2010/493935>
- Jacobus LM (2013) A new junior synonym for the Holarctic species *Ametropus fragilis* Albarda, 1878 (Insecta: Ephemeroptera: Ametropodidae). *Proceedings of the Indiana Academy of Science* 122: 18–19.
- Jacobus LM (2015) Book review: *The Mayflies of Europe*. *American Entomologist* 61(3): 190–191. <https://doi.org/10.1093/ae/tmv056>
- Jacobus LM (2016) Addendum to Book Review of *The Mayflies of Europe*, by Bauernfeind & Soldan (2012), Published in *American Entomologist* 61(3), DOI: 10.1093/ae/tmv056. *The Mayfly Newsletter* 19(1): 8–9.
- Jacobus LM (2017) Identifying larvae of eastern North American mayfly species (Insecta: Ephemeroptera). Southeast Water Pollution Biologists Association, Grand Rivers, Kentucky, 29 pp.
- Jacobus LM (2018) Endemism among Canada's mayfly species (Insecta: Ephemeroptera). *Newsletter of the Biological Survey of Canada* 37(1): 13–19.
- Jacobus LM, McCafferty WP (2001) Additions to the Canadian Ephemeroptera. *Journal of the New York Entomological Society* 109: 367–371. [https://doi.org/10.1664/0028-7199\(2001\)109\[0367:ATTCE\]2.0.CO;2](https://doi.org/10.1664/0028-7199(2001)109[0367:ATTCE]2.0.CO;2)
- Jacobus LM, McCafferty WP (2005) *Apobaetis futilis*, a new combination in Nearctic Baetidae (Ephemeroptera). *Proceedings of the Entomological Society of Washington* 107: 979.
- Jacobus LM, McCafferty WP (2006) Reevaluation of the phylogeny of the Ephemeroptera Infraorder Pannota (Furcatergalia), with adjustments to higher classification. *Transactions of the American Entomological Society* 132: 81–90, 429–430. [https://doi.org/10.3157/0002-8320\(2006\)132\[81:ROTPOT\]2.0.CO;2](https://doi.org/10.3157/0002-8320(2006)132[81:ROTPOT]2.0.CO;2)
- Jacobus LM, McCafferty WP (2007) Reinstatement of *Serratella serratooides* (McDunnough) and status of *Ephemerella molita* McDunnough (Ephemeroptera: Ephemerellidae). *Proceedings of the Entomological Society of Washington* 109: 730–732.
- Jacobus LM, McCafferty WP (2008) Revision of Ephemerellidae genera (Ephemeroptera). *Transactions of the American Entomological Society* 134: 185–274. [https://doi.org/10.3157/0002-8320\(2008\)134\[185:ROEGE\]2.0.CO;2](https://doi.org/10.3157/0002-8320(2008)134[185:ROEGE]2.0.CO;2)

- Jacobus LM, Wiersema NA (2014) The genera *Anafroptilum* Kluge, 2011 and *Neocloeon* Traver, 1932, reinstated status, in North America, with remarks about the global composition of *Centroptilum* Eaton, 1869 (Ephemeroptera: Baetidae). *Zootaxa* 3814: 385–391. <https://doi.org/10.11646/zootaxa.3814.3.5>
- Jacobus LM, Wiersema NA, Webb JM (2014) Identification of Far Northern and Western North American mayfly larvae (Insecta: Ephemeroptera), north of Mexico; Version 2. Joint Aquatic Science meeting, Portland, Oregon, 176 pp. + suppl.
- Kjørstad G, Webb JM, Ekrem T (2012) A review of the Ephemeroptera of Finnmark – DNA barcodes identify Holarctic relations. *Norwegian Journal of Entomology* 59: 182–195.
- Klubertanz TH (2016a) Mayfly larvae of Wisconsin. Cooperative Extension Publishing, University of Wisconsin, Madison, Wisconsin, viii + 291 pp.
- Klubertanz TH (2016b) New records of mayflies (Ephemeroptera) from Kenora District and Northwestern Ontario, Canada. *Entomological News* 125: 341–350. <https://doi.org/10.3157/021.125.0505>
- Kluge NJ (1980) To the knowledge of mayflies (Ephemeroptera) of Taimyr National District. *Entomologicheskoe Obozrenie* 59(3): 561–579.
- Kluge NJ (1996) The Palaearctic Metretopodidae, with description of a new genus and species from Siberia (Ephemeroptera). *Zoosystematica Rossica* (1995) 4(1): 76–80.
- Kluge NJ, Novikova EA (2011) Systematics of the mayfly taxon *Acentrella* (Ephemeroptera, Baetidae), *Russian Entomological Journal* 20(1): 1–60.
- Kondratieff BC, Voshell Jr JR (1981) A new species of *Siphonurus* from Nova Scotia, Canada. *Annals of the Entomological Society of America* 74: 545–547. <https://doi.org/10.1093/aesa/74.6.545>
- Kondratieff BC, Voshell Jr JR (1984) The North and Central American species of *Isonychia* (Ephemeroptera: Oligoneuriidae). *Transactions of the American Entomological Society* 110: 129–244.
- Landa V, Soldán T (1985) Phylogeny and higher classification of the order Ephemeroptera: a discussion from the comparative anatomical point of view. *Studie CSAV, Československa Akademie Ved, Praha*, 121 pp.
- Langor DW (2019) The diversity of terrestrial arthropods in Canada. In: Langor DW, Sheffield CS (Eds) *The Biota of Canada – A Biodiversity Assessment. Part 1: The Terrestrial Arthropods*. *ZooKeys* 819: 9–40. <https://doi.org/10.3897/zookeys.819.31947>
- Lehmkuhl DM (1976) Additions to the taxonomy, zoogeography and biology of *Analetris eximia* (Acanthametropodinae: Siphonuridae: Ephemeroptera). *The Canadian Entomologist* 108: 199–207. <https://doi.org/10.4039/Ent108199-2>
- Lehmkuhl DM (1979) Ephemeroptera. In: Danks HV (Ed.) *Canada and its insect fauna*. *Memoirs of the Entomological Society of Canada* No. 108, 305–308. <https://doi.org/10.4039/entm111108305b-1>
- Lomond TM, Colbo MH (2000) Variations in lake-outlet Ephemeroptera, Plecoptera, and Trichoptera communities amongst regions of eastern Newfoundland, Canada. *Canadian Journal of Zoology* 78: 1536–1543. <https://doi.org/10.1139/z00-098>
- Lugo-Ortiz CR, McCafferty WP (1998) A new North American genus of Baetidae (Ephemeroptera) and key to *Baetis* complex genera. *Entomological News* 109: 345–353.
- McCafferty WP (1975) The burrowing mayflies (Ephemeroptera: Ephemeroidea) of the United States. *Transactions of the American Entomological Society* 101: 447–504.

- McCafferty WP (1991a) Comparison of Old and New World *Acanthametropus* (Ephemeroptera: Acanthametropodidae) and other psammophilous mayflies. *Entomological News* 102: 205–214.
- McCafferty WP (1991b) Toward a phylogenetic classification of the Ephemeroptera (Insecta): a commentary on systematics. *Annals of the Entomological Society of America* 84: 343–360. <https://doi.org/10.1093/aesa/84.4.343>
- McCafferty WP (1992) New larval descriptions and comparisons of North American *Choroterpes* (Ephemeroptera: Leptophlebiidae). *The Great Lakes Entomologist* 25: 71–78.
- McCafferty WP (1994) Distributional and classificatory supplement to the burrowing mayflies (Ephemeroptera: Ephemeroidea) of the United States. *Entomological News* 105: 1–13.
- McCafferty WP (1996) The mayflies of North America online. *Entomological News* 107: 61–63.
- McCafferty WP (2001) The gentle quest: 200 years in search of North American mayflies. In: Dominguez E (Ed.) *Trends in Research in Ephemeroptera and Plecoptera*. Springer, New York, 21–35. https://doi.org/10.1007/978-1-4615-1257-8_5
- McCafferty WP (2004) Higher classification of the burrowing mayflies (Ephemeroptera: Scaphodonta). *Entomological News* 115: 84–92.
- McCafferty WP (2009) New state and provincial North American records for 100 Ephemeroptera species. *Transactions of the American Entomological Society* 135: 353–368. <https://doi.org/10.3157/061.135.0302>
- McCafferty WP (2011a) A new genus and species of small minnow mayflies (Ephemeroptera: Baetidae) from far northern North America. *Transactions of the American Entomological Society* 137: 11–14.
- McCafferty WP (2011b) Notable new North and Central American records of Ephemeroptera species. *Transactions of the American Entomological Society* 137: 1–10.
- McCafferty WP, Bae YJ (1992) Taxonomic status of historically confused species of Potamanthidae and Heptageniidae (Ephemeroptera). *Proceedings of the Entomological Society of Washington* 94: 169–171.
- McCafferty WP, Edmunds Jr GF (1997) Critical commentary on the genus *Siphonisca* (Ephemeroptera: Siphonuridae). *Entomological News* 108: 141–147.
- McCafferty WP, Jacobus LM (2018) Mayfly central. <https://www.entm.purdue.edu/mayfly/> [Accessed 19.III.2018]
- McCafferty WP, Randolph RP (1998) Canada mayflies: a faunistic compendium. *Proceedings of the Entomological Society of Ontario* 129: 47–97.
- McCafferty WP, Waltz RD (1990) Revisionary synopsis of the Baetidae (Ephemeroptera) of North and Middle America. *Transactions of the American Entomological Society* 116: 769–799.
- McCafferty WP, Randolph RP, Jacobus LM (2012) Mayflies of the Intermountain West. *Memoirs of the American Entomological Institute* 85: i–iii + 1–317.
- McCafferty WP, Stark BP, Provonsha AV (1990) Ephemeroptera, Plecoptera, and Odonata. In: Koszarab M, Schaefer C (Eds) *Systematics of the North American insects and arachnids: status and needs*. Virginia Agricultural Experiment Station Information Series No. 90-1, Virginia Polytechnic Institute and State University, Blacksburg, Virginia, 43–58.

- McCafferty WP, Jacobus LM, Provonsha AV, Wiersema NA (2017) Ephemeroptera. In: Morse JC, McCafferty WP, Stark BP, Jacobus LM (Eds) Larvae of the Southeastern USA Mayfly, Stonefly, and Caddisfly Species (Ephemeroptera, Plecoptera, and Trichoptera). Biota of South Carolina, Vol. 9. Clemson University Public Service Publishing, Clemson University, Clemson, South Carolina, 15–160.
- McCafferty WP, Jacobus LM, Webb JM, Meyer MD (2008) Insecta, Ephemeroptera: Range extensions and new records for Ontario and Canada. Check List 4: 449–452. <https://doi.org/10.15560/4.4.445>
- McCafferty WP, Lenat DR, Jacobus LM, Meyer MD (2010) The mayflies (Ephemeroptera) of the southeastern United States. Transactions of the American Entomological Society 136: 221–233. <https://doi.org/10.3157/061.136.0303>
- McCafferty WP, Meyer MD, Webb JM, Jacobus LM (2004) New state and provincial records for North American small minnow mayflies (Ephemeroptera: Baetidae). Entomological News 115: 93–100.
- McCafferty WP, Waltz RD, Webb JM, Jacobus LM (2005) Revision of *Heterocloeon* McDunnough (Ephemeroptera: Baetidae). Journal of Insect Science 5(35): 1–11. <https://doi.org/10.1093/jis/5.1.35>
- McDunnough J (1925) New Canadian Ephemeridae with notes. III. The Canadian Entomologist 57: 168–176, 185–192. <https://doi.org/10.4039/Ent57185-8>
- Meyer MD, McCafferty WP (2007) Mayflies (Ephemeroptera) of the far western United States. Part I: Washington. Transactions of the American Entomological Society 133: 21–63. [https://doi.org/10.3157/0002-8320\(2007\)133\[21:MEOTFW\]2.0.CO;2](https://doi.org/10.3157/0002-8320(2007)133[21:MEOTFW]2.0.CO;2)
- Miller DB, Bartlett S, Sartori M, Breinholt JW, Ogden TH (2018) Anchored phylogenomics of burrowing mayflies (Ephemeroptera) and the evolution of tusks. Systematic Entomology 43(4): 692–701. <https://doi.org/10.1111/syen.12298>
- Miyazaki R, Lehmkuhl DM (2011) Insects of the Saskatchewan River system in Saskatchewan. Arthropods of Canadian Grasslands 2: 119–157.
- Molineri C (2010) A cladistic revision of *Tortopus* Needham & Murphy with description of the new genus *Tortopsis* (Ephemeroptera: Polymitarcyidae). Zootaxa 2481: 1–36.
- Newell RL, Anderson ML (2009) Note on the occurrence of *Siphonurus autumnalis* (Ephemeroptera: Siphonuridae) in a Montana spring brook. Western North American Naturalist 69: 551–555. <https://doi.org/10.3398/064.069.0415>
- Novikova EA, Kluge NJ (1987) Systematics of the genus *Baetis* (Ephemeroptera, Baetidae), with description of new species from Middle Asia. Vestnik Zoologii 4: 8–19.
- Ogden TH, Sartori M, Whiting MF (2008) Pisciforma, Setisura, and Furcatergalia (order: Ephemeroptera) are not monophyletic based on 18S rDNA sequences: a response to Sun et al. (2006) Annals of the Entomological Society of America 101: 1–6. [https://doi.org/10.1603/0013-8746\(2008\)101\[1:PSAFOE\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2008)101[1:PSAFOE]2.0.CO;2)
- Ogden TH, Gattolliat J-L, Sartori M, Staniczek AH, Soldán T, Whiting MF (2009) Towards a new paradigm in mayfly phylogeny (Ephemeroptera): combined analysis of morphological and molecular data. Systematic Entomology 34: 616–634. <https://doi.org/10.1111/j.1365-3113.2009.00488.x>

- Pescador ML (1985) Systematics of the Nearctic genus *Pseudiron* (Ephemeroptera: Heptageniidae: Pseudironinae). The Florida Entomologist 68: 432–444. <https://doi.org/10.2307/3495130>
- Pescador ML, Berner L (1981) The mayfly family Baetiscidae (Ephemeroptera). Part II. Biosystematics of the genus *Baetisca*. Transactions of the American Entomological Society 107: 163–228.
- Peters WL (1988) Origins of the North American Ephemeroptera fauna, especially the Leptophlebiidae. Memoirs of the Entomological Society of Canada No. 144, 13–24. <https://doi.org/10.4039/entm120144013-1>
- Peters WL, Peters JG (1993) Status changes in Leptohyphidae and Tricorythidae (Ephemeroptera). Aquatic Insects 15: 45–48. <https://doi.org/10.1080/01650429309361499>
- Prisecaru FS, Tabacaru I, Prisecaru M, Stoica I, Călin M (2014) Contributions to a revised species concept of the Ephemeroptera fauna from Romania (mayfliesyst). Studii si Cercetari, Biologie, Universitatea “Vasile Alecsandri” din Bacau 23(2): 20–30.
- Provonsha AV (1990) A revision of the genus *Caenis* in North America (Ephemeroptera: Caenidae). Transactions of the American Entomological Society 116: 801–884.
- Provonsha AV, McCafferty WP (1982) New species and previously undescribed larvae of North American Ephemeroptera. Journal of the Kansas Entomological Society 55: 23–33.
- Randolph RP, McCafferty WP (2001) Nunavut mayflies (Ephemeroptera) a supplement for far northern North America. Entomological News 112: 56–58.
- Randolph RP, McCafferty WP, Zaranko D, Jacobus LM, Webb JM (2002) New Canadian records of Baetidae (Ephemeroptera) and adjustments to North American *Cloeon*. Entomological News 113: 306–309.
- Ratnasingham S, Hebert PDN (2013) A DNA-based registry for all animal species: the Barcode Index Number (BIN) system. PLoS ONE 8(7): e66213. <https://doi.org/10.1371/journal.pone.0066213>
- Savolainen E, Drotz MK, Saura A, Ståhls G (2014) *Baetis bundyae* (Ephemeroptera: Baetidae), described from Arctic Canada is found in northernmost Europe. The Canadian Entomologist 146: 621–629. <https://doi.org/10.4039/tce.2014.21>
- Schmude KL, Tracey DL, Klubertanz TH, Dimick JJ (2012) First report of Neoephemeridae (Ephemeroptera) in Wisconsin. The Great Lakes Entomologist 45: 99–101.
- Sun L, McCafferty WP (2008) Cladistics, classification and identification of the brachycercine mayflies (Insecta: Ephemeroptera: Caenidae). Zootaxa 1801: 1–239.
- Sun L, Sabo A, Meyer MD, Randolph RP, Jacobus LM, McCafferty WP, Ferris VR (2006) Tests of current hypotheses of mayfly (Ephemeroptera) phylogeny using molecular (18s rDNA) data. Annals of the Entomological Society of America 99: 241–252. [https://doi.org/10.1603/0013-8746\(2006\)099\[0241:TOCHOM\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2006)099[0241:TOCHOM]2.0.CO;2)
- Tiunova TM, Kluge N (2016) Redescription of *Paraleptophlebia falcata* Traver 1934 with notes on status and composition of *Paraleptophlebia* Lestage 1917 and *Neoleptophlebia* Kluge 1997 (Ephemeroptera: Leptophlebiidae). Zootaxa 4098: 369–382. <https://doi.org/10.11646/zootaxa.4098.2.9>
- Traver JR (1935) Part II: Systematic. In: Needham JG, Traver JR, Hsu Y-C (Eds) The Biology of Mayflies, with a systematic account of North American species. Comstock, Ithaca, New York, 237–739.

- Waltz RD, Burian SK (2008) Ephemeroptera. In: Merritt RW, Cummins KW, Berg MB (Eds) An Introduction to the Aquatic Insects of North America. Kendall/Hunt, Dubuque, Iowa, 181–236.
- Wang T-Q, McCafferty WP (1995) Relationships of the Arthropleidae, Heptageniidae, and Pseudironidae (Ephemeroptera: Heptagenioidea). Entomological News 106: 251–256.
- Webb JM, McCafferty WP (2003) New records of mayflies (Ephemeroptera) from Alberta. Entomological News 114: 230–232.
- Webb JM, McCafferty WP (2008) Heptageniidae of the world. Part II: Key to the genera. Canadian Journal of Arthropod Identification 7. <https://doi.org/10.37551/cjai2008.7>
- Webb JM, McCafferty WP (2011) Contributions to the larvae of North American *Nixe* (Ephemeroptera: Heptageniidae), with description of *N. dorotheae* sp. nov. from southern Indiana. Zootaxa 3065: 27–37.
- Webb JM, Jacobus LM, Sullivan SP (2018) The state of systematics of North American *Baetis* Leach, 1815 (Ephemeroptera: Baetidae), with recommendations for identification of larvae. Zootaxa 4394: 105–127. <https://doi.org/10.11646/zootaxa.4394.1.6>
- Webb JM, Parker DW, Lehmkuhl DM, McCafferty WP (2004) Additions and emendations to the mayfly fauna of Saskatchewan, Canada. Entomological News 115: 213–218.
- Webb JM, Sun L, McCafferty WP, Ferris VR (2007) A new species and new synonym in *Heptagenia* Walsh (Ephemeroptera: Heptageniidae: Heptageniinae) based on molecular and morphological evidence. Journal of Insect Science 7(63): 1–16. <https://doi.org/10.1673/031.007.6301>
- Webb JM, Jacobus LM, Funk DH, Zhou X, Kondratieff B, Geraci CJ, DeWalt RE, Baird D, Richard B, Phillips I, Hebert PDN (2012) A DNA Barcode Library for North American Ephemeroptera: Progress and Prospects. PLoS ONE 7(5): e38063. <https://doi.org/10.1371/journal.pone.0038063>
- Wiersema NA, McCafferty WP (2000) Generic revision of the North and Central American Leptohiphidae (Ephemeroptera: Pannota). Transactions of the American Entomological Society 126: 337–371.
- Wiersema NA, McCafferty WP (2005) Contribution to the taxonomy of *Asioplax* (Ephemeroptera: Leptohiphidae: Tricorythodinae). Entomological News 116: 147–158.
- Zhou X, Adamowicz SJ, Jacobus LM, DeWalt RE, Hebert PDN (2009) Towards a comprehensive barcode library for Arctic life: Ephemeroptera, Plecoptera and Trichoptera of Churchill, Manitoba, Canada. Frontiers in Zoology 6(30). <https://doi.org/10.1186/1742-9994-6-30>
- Zhou X, Jacobus LM, DeWalt RE, Adamowicz SJ, Hebert PDN (2010) The Ephemeroptera, Plecoptera, and Trichoptera fauna of Churchill (Manitoba, Canada): insights into biodiversity patterns from DNA barcoding. Journal of the North American Benthological Society 29: 814–837. <https://doi.org/10.1899/09-121.1>
- Zloty J (1996) A revision of the Nearctic *Ameletus* mayflies based on adult males, with descriptions of seven new species (Ephemeroptera: Ameletidae). The Canadian Entomologist 128: 293–346. <https://doi.org/10.4039/Ent128293-2>
- Zloty J, Harper F (1999) Two new *Ameletus* mayflies (Ephemeroptera: Ameletidae) from western North America. The Canadian Entomologist 131: 1–9. <https://doi.org/10.4039/Ent1311-1>