

Checklist of acanthocephalan parasites of South Africa

Ali Halajian¹, Lesley R. Smales², Sareh Tavakol¹, Nico J. Smit³,
Wilmien J. Luus-Powell¹

1 DST-NRF SARChI Research Chair (Ecosystem Health), Department of Biodiversity, University of Limpopo, Sovenga, 0727, South Africa **2** Parasitology Section, South Australia Museum, North Terrace, Adelaide 5000 South Australia, Australia **3** Water Research Group, Unit for Environmental Sciences and Management, North-West University, Potchefstroom, South Africa

Corresponding author: Ali Halajian (ali_hal572002@yahoo.com)

Academic editor: Matthew Wayland | Received 23 June 2018 | Accepted 11 August 2018 | Published 10 October 2018

<http://zoobank.org/9C50AE22-CD06-478C-B81A-D74E5F0E384D>

Citation: Halajian A, Smales LR, Tavakol S, Smit NJ, Luus-Powell WJ (2018) Checklist of acanthocephalan parasites of South Africa. ZooKeys 789: 1–18. <https://doi.org/10.3897/zookeys.789.27710>

Abstract

Twenty-one species of acanthocephalans, representative of thirteen genera from ten families of seven orders and three classes, are included in this updated checklist of acanthocephalans in South Africa. Although South Africa appears to have a less diverse acanthocephalan fauna compared to some other countries such as Iran in Asia, or Brazil in South America, this is probably an artefact of fewer parasitological surveys.

Keywords

Acanthocephala, helminths, diversity, wildlife, Africa

Introduction

South Africa's landscape is the third most biologically diverse in the world with 6% of the world's mammal species, 8% of bird species and 5% of reptile species of which many are endemic (da Silva and Willows-Munro 2016). In regard to invertebrates, only 36,803 species are listed for Africa, and 12,098 for South Africa (Hamer 2010). It is said that in South African habitats, there are many undiscovered and undescribed animal species, especially invertebrates. It is estimated that as many as 80,000 South

African animal species remain to be discovered or described, and most of these animals are invertebrates (Hamer 2013).

The most comprehensive checklist of helminth parasites of Africa was compiled for freshwater fishes by Khalil (1971) and updated by Khalil and Polling (1997). It included very few records of acanthocephalans considering the total number of freshwater fish species present in the continent. The updated list (Khalil and Polling 1997) comprised 568 adult helminth parasites of which only 21 species are acanthocephalans. These records were compiled from 359 species of African freshwater fishes (assigned to 89 genera belonging to 32 families) of an estimated 3000 existing inland fish species (Khalil and Polling 1997). These examples illustrate the lack of knowledge of the helminth fauna of the wildlife of the African continent in general and in South Africa in particular.

This is the first checklist of acanthocephalans of South Africa and the aim is to provide a comprehensive record of all the previously reported species of Acanthocephala occurring in South African hosts as well as new records from our on-going research on parasites of wildlife, while simultaneously demonstrating the need for more extensive parasitological surveys.

Materials and methods

Data were obtained from two sources, published records and our own ongoing studies on the Acanthocephala of South African wildlife. These data are presented in two parts. In the first part parasites are listed systematically, with families, genera, and species in alphabetical order. The scientific name, including any synonyms, followed by the scientific and common name of the host, the locality in which the parasite was reported and museum (location) of type specimens where known. In the second part, the hosts are listed systematically by their scientific names and parasite records from each host are given together with locality and reference. The records without references are those of our ongoing study that are being reported here for the first time.

Classification of the Acanthocephala follows Amin (2013). For the hosts, fish taxonomy is based on Skelton (2001, 2016) and Fishbase (Froese and Pauly 2016), bird taxonomy is based on Clements et al. (2016) and mammal taxonomy on Wilson and Reader (2005) and Apps (2012).

Abbreviations for museums are:

BMNH	Natural History Museum London, London, UK;
GNM	Gothenburg Natural History Museum, Gothenburg, Sweden;
SAMCTA	South African Museum at Cape Town, South Africa;
SAM	South Australian Museum, Adelaide, Australia;
USNM Helm. Coll.	United States National Museum Helminthological Collection
USNPC	United States National Parasite Collection now held in the Invertebrate Zoology collection of the Smithsonian Museum, Washington, USA.

Acanthocephalan specimens from our ongoing wildlife parasitology projects were mostly collected from roadkill animals, museum collections, hunting/culling surveys and other research permits received for a limited number of specimens through the Limpopo Department of Economic Development, Environment and Tourism (LEDET) (permit number CPM004961 and ZA/LP/HO/3370 for freshwater fish research, 001-CPM403-00012 and ZA/LP/HO/3448 for frogs, ZA/LP/HO/3432 for rodents and ZA/LP/87586 for roadkills).

Acanthocephalans from freshly dead animals were placed in tap water and refrigerated for a few hours to one day until the proboscis was everted and then fixed and stored in 70% ethanol until studied. Acanthocephalans from frozen hosts were fixed and stored in 70% ethanol. Some specimens were prepared for examination by staining in Mayer's acid carmine, destained in HCl in 70% ethanol, dehydrated through increasing concentrations of ethanol, cleared in xylene, and mounted as whole worms in Canada balsam. Other worms were examined as temporary mounts following clearing in lactophenol or beechwood creosote.

A total of 102 species of birds (151 individuals), 72 of mammals (420 individuals), 9 of reptiles (18 individuals) and 42 (1050 individuals) of fishes were examined for this study (details in Table 1).

Table 1. Total number of host taxa examined and those infected with acanthocephalans (i.e. number of taxa that harboured acanthocephalans, in parenthesis) during our ongoing study on wildlife parasites in South Africa.

Taxon Group	Order	Family	Genus	Species
Amphibians	1 (0)	8 (0)	13 (0)	19 (0)
Birds	21 (5)	50 (5)	87 (5)	102 (5)
Fishes (freshwater)	8 (1)	13 (1)	24 (1)	42 (1)
Mammals	10 (4)	26 (4)	59 (6)	72 (6)
Reptiles	3 (1)	6 (1)	7 (1)	9 (1)
Totals	43 (11)	103 (11)	190 (13)	244 (13)

Parasite-Host List

Acanthocephala

Class: Archiacanthocephala Meyer, 1931

Order: Gigantorhynchida Southwell & Macfie, 1925

Family: Gigantorhynchidae Hamann, 1892

Genus: *Mediorhynchus* Van Cleave, 1916

Mediorhynchus africanus Amin, Evans, Heckmann & El-Naggar, 2013

Empodius segmentatus (de Marval, 1902) Southwell & Mac-Fie, 1925

Mediorhynchus selengensis Harris, 1973

M. gallinarum (Bhalerao, 1937) Van Cleave, 1947 *sensu* Junker & Boomker, 2006

Notes. *M. gallinarum* is found only in Asia and the species in Africa is actually *M. africanus* (Amin et al. 2013; Amin 2013) and not *M. gallinarum* previously reported in South Africa (Junker and Boomker 2006).

Host. *Numida meleagris* (L. 1758) (Helmeted Guineafowl) (Numididae) (type host).

Localities. Kruger National Park, Mpumalanga Province, South Africa (type locality) (Junker and Boomker 2006); Vicinity of Petrus Steyn, Free State Province, South Africa (Davies et al. 2008); Musina, Limpopo Province, South Africa (Junker and Boomker 2007; Junker et al. 2008).

***Mediorhynchus mokgalongi* Smales & Halajian, 2018**

Host. *Turdus smithi* Bonaparte, 1850 (Karoo Thrush) (Turdidae) (type host).

Locality. Polokwane, Limpopo Province, South Africa (type locality) (Smales et al. 2018).

Type specimens. Holotype male SAM AHC 48068, allotype female SAM AHC 48069, paratype SAM AHC 48070.

***Mediorhynchus numidae* (Baer, 1925) Meyer, 1932**

Heteroplus numidae Baer, 1925; *Empodisma numidae* (Baer, 1925) Yamaguti, 1963

Host. *Numida meleagris* (L. 1758) (Helmeted Guineafowl) (Numididae)

Locality. Pretoria, Gauteng Province, South Africa (Oosthuizen and Markus 1967).

***Mediorhynchus taeniatus* (von Linstow, 1901) Dollfus, 1936**

Echinorhynchus taeniatus von Linstow, 1901; *E. segmentatus* de Marval, 1902

Host. *Numida meleagris* (L. 1758) (Helmeted Guineafowl) (Numididae).

Locality. Rooipoort farm, Kimberley, Northern Cape Province, South Africa (Crowe 1977).

Host. *Tockus leucomelas* (Lichtenstein, 1842) (Southern Yellow-billed Hornbill) (Bucerotidae)

Locality. Limpopo Province, South Africa.

Order: Moniliformida Schmidt, 1972
Family: Moniliformidae Van Cleave, 1924
Genus: *Moniliformis* Travassos, 1915

***Moniliformis kalahariensis* Meyer, 1931**

Host. *Atelerix frontalis* (Smith, 1831) (Southern African Hedgehog) (Erinaceidae)

Locality. Mhlonong village and University of Limpopo, Limpopo Province, South Africa (Amin et al. 2014).

***Moniliformis moniliformis* (Bremser, 1811) Travassos, 1915 (type species)**

Echinorhynchus moniliformis Bremser, 1811

E. grassi Railliet, 1893

E. canis Porter, 1914

E. belgicus Railliet, 1919

Moniliformis moniliformis aegypticus Meyer in Petrochenko, 1958

M. dubius Meyer, 1932

M. travassosi Meyer, 1932 (*vide* Machado Filho 1946, Van Cleave 1952)

Host. *Atelerix frontalis* (Smith, 1831) (Southern African Hedgehog) (Erinaceidae).

Locality. Hammanskraal, Gauteng Province, South Africa (Le Roux 1930).

Notes. Host recorded as *Aethechinus frontalis* in Le Roux (1930).

***Moniliformis acomysi* Ward & Nelson, 1967**

Host. *Gerbilliscus leucogaster* (Peters, 1852) (Bushveld Gerbil), *Mastomys natalensis* (Smith, 1834) (Natal Mastomys), *Mus minutoides* (Pygmy mouse) (Muridae).

Localities. Bloemhof, Free State Province; Vyeboom village, Limpopo Province; Hoopstad, Free State Province; South Africa.

***Moniliformis* sp.**

Host. *Otolemur crassicaudatus* (É. Geoffroy Saint-Hilaire, 1812) (Thick-tailed Bush-baby) (Galagidae).

Locality. Venda, Limpopo Province, South Africa.

Remarks. One male and one female worm were found in the small intestine of an adult bushbaby.

Order: Oligacanthorhynchida Petrochenko, 1956
Family: Oligacanthorhynchidae Southwell & Macfie, 1925
Genus: *Heptamegacanthus* Spencer Jones, 1990

Heptamegacanthus niekerki Spencer Jones, 1990 (type species)

Host. *Chrysofalax trevelyani* (Günther, 1875) (Giant Golden Mole) (Chrysochloridae) (type host).

Locality. Nqadu Forest, Transkei, Eastern Cape Province, South Africa (type locality) (Spencer Jones 1990).

Type specimens. Holotype male BMNH 1988.2480; allotype female BMNH 1988.2481; paratypes BMNH 1988.2482–2491.

Oligacanthorhynchidae sp.

Host. *Varanus albigularis* Daudin, 1802 (Rock Monitor) (Varanidae).

Localities. Tzaneen; Tolwe, Limpopo Province, South Africa.

Class: Eoacanthocephala Van Cleave, 1936
Order: Gyracanthocephala Van Cleave, 1936
Family: Quadrigyridae Van Cleave, 1920
Genus: *Acanthogyryrus* Thapar, 1927
Subgenus: *Acanthosentis* Verma & Datta, 1929

Acanthogyryrus (Acanthosentis) phillipi (Mashego, 1988) Amin, 2005

Acanthosentis phillipi Mashego, 1988

Host. *Enteromius neefi* (Greenwood, 1962) (syn. *Barbus neefi*) (Sidespot Barb) (Cyprinidae) (type host).

Locality. Lingwe River, Venda, Limpopo Province, South Africa (type locality) (Mashego 1988).

Type specimens. Holotype in Transvaal Museum no. TM14659; Paratypes TM5 at University of Limpopo, Zoology, 5.

***Acanthogyryrus* sp.**

Host. *Oreochromis mossambicus* (Peters, 1852) (Mozambique Tilapia) (Cichlidae).

Locality. Molepo Dam, Limpopo Province, South Africa (Kunutu et al. 2013).

Order: Neoechinorhynchida Southwell & Macfie, 1925

Family: Neoechinorhynchidae (Ward, 1917) Van Cleave, 1928

Genus: *Neoechinorhynchus* Stiles & Hassall, 1905

Subgenus: *Neoechinorhynchus* Hamann, 1892

***Neoechinorhynchus (Neoechinorhynchus) dorsovaginatus* Amin & Christison, 2005**

Host. *Argyrosomus japonicus* (Temminck & Schlegel, 1843) (Japanese Meagre, Dusky Kob) Sciaenidae) (type host).

Locality. Breede River Estuary, Western Cape Province, South Africa (type locality) (Amin and Christison 2005).

Type specimens. No. SAMCTA29536 (holotype male and allotype female; same slide), nos SAMCTA29537-29545 (paratypes), USNPC no. 94918 (paratypes).

Class: Palaeacanthocephala Meyer, 1931

Order: Echinorhynchida Southwell & Macfie, 1925

Family: Pomphorhynchidae Yamaguti, 1939

Genus: *Longicollum* Yamaguti, 1935

***Longicollum chabanaudi* Dollfus & Golvan, 1963**

Host. *Barnardichthys fulvomarginata* (Gilchrist, 1904) (syn. *B. fulvomarginatus*) (Soleidae) (Sole).

Locality. False Bay, Western Cape Province, South Africa (Dollfus and Golvan 1963).

***Longicollum* sp. innom.**

Host. *Pegusa nasuta* (Pallas, 1814) (syn. *Solea bleekeri*) (Blackhand Sole) (Soleidae).

Locality. Klein River estuary, Hermanus, Western Cape Province, South Africa (Bray 1974).

Remarks. Three contracted and immature specimens were found in one fish and one male in another fish. Thus it could not be identified to species level (Bray 1974).

Family: Rhadinorhynchidae Lühe, 1912

Genus: *Rhadinorhynchus* Lühe, 1911

***Rhadinorhynchus cadenati* (Golvan & Houin, 1964) Golvan, 1969**

Nipporhynchus cadenati Golvan & Houin, 1964

Host. *Thyrsites atun* (Euphrasen, 1791) (Snoek) (Gempylidae).

Locality. South Africa's West and South coasts (Nunkoo et al. 2016).

***Rhadinorbynchus capensis* Bray, 1974**

Host. *Pegusa nasuta* (Pallas, 1814) (syn. *Solea bleekeri*) (Blackhand Sole) (Soleidae) (type host)

Locality. Heuninges River estuary, near Cape Agulhas, Western Cape Province, South Africa (type locality) (Bray 1974).

Type specimens. British Museum, Registration number 1974.521-550.

***Rhadinorbynchus* sp.**

Host. *Ruvettus pretiosus* Cocco, 1833 (Oilfish) (Gempylidae).

Locality. South Africa's West coast, Atlantic Ocean (Nunkoo et al. 2017).

Order: Polymorphida Petrochenko, 1956

Family: Centrorhynchidae Van Cleave, 1916 (Golvan, 1960)

Genus: *Centrorhynchus* Lühe, 1911

***Centrorhynchus sarebae* Smales & Halajian, 2017**

Host. *Kaupifalco monogrammicus* (Temminck, 1824) (Lizard Buzzard) (Accipitridae) (type host).

Locality. Makhado (Louis Trichardt), Limpopo Province, South Africa (type locality) (Smales et al. 2017).

Type Specimens. Holotype male SAM AHC 47858; allotype female SAM AHC 47859; paratypes SAM AHC 47860.

***Centrorhynchus clitorideus* (Meyer, 1931) Golvan, 1958**

Gordiorhynchus clitorideus Meyer, 1931 (*nec clitorideum*)

Host. *Bubo africanus* (Temminck, 1821) (Spotted Eagle Owl) (Strigidae).

Locality. Zandriverspoort Farm, Polokwane, Limpopo Province, South Africa.

***Centrorhynchus* sp.**

Host. *Felis catus* L., 1758 (Domestic Cat) (Felidae).

Locality. Pretoria, Gauteng Province, South Africa (Baker et al. 1989).

***Centrorhynchus* sp.**

Host. *Mungos mungo* (Gmelin, 1788) (Banded Mongoose) (Herpestidae).

Locality. Polokwane, Limpopo Province, South Africa.

Family: Plagiorhynchidae Golvan, 1960

Genus: *Plagiorhynchus* Lühe, 1911

Subgenus: *Prosthorhynchus* Kostylew, 1915

***Plagiorhynchus (Prosthorhynchus) cylindraceus* (Goeze, 1782) Schmidt & Kuntz, 1966**

Echinorhynchus cylindraceus Goeze, 1782

E. pici Gmelin, 1791 *fide* Florescu and Ienistea 1984

E. merulae Gmelin, 1791 *fide* Florescu and Ienistea 1984

E. transversus (Rudolphi, 1819) Travassos 1926

E. obliquus Dujardin, 1845 *fide* Florescu and Ienistea 1984

Centrorhynchus cylindraceus (Goeze 1782) Kostylew, 1914

C. fasciatus (Westrumb, 1821) Travassos, 1926 *fide* de Marval 1905

C. rostratus de Marval, 1902 *fide* Florescu and Ienistea 1984

Prosthorhynchus rosai (Porta, 1910) Meyer, 1932

Prosthorhynchus rostratus (de Marval, 1902) Meyer, 1932

Plagiorhynchus formosus Van Cleave, 1918 *fide* Amin et al. 1999

Plagiorhynchus taiwanensis Schmidt et Kuntz, 1966 *fide* Amin et al. 1999.

Host. *Calidris ferruginea* (Pontoppidan, 1763) (Curlew Sandpiper); *Charadrius pecuarius* Temminck, 1823 (Kittlitz's Plover); *Charadrius tricollaris* Vieillot, 1818 (Three-banded Plover) (Charadriidae).

Locality. Berg River, Western Cape Province, South Africa (Amin et al. 1999).

Host. *Crexopsis egregia* (Peters, 1854) (syn. *Crex egregia*) (African Crane) (Rallidae).

Locality. Blouberg, Limpopo Province, South Africa.

Host. *Vanellus armatus* (Burchell, 1822) (Blacksmith Lapwing) (Charadriidae).

Locality. Berg River, Western Cape Province, South Africa (Amin et al. 1999).

Unidentified plagiorhynchid

Host. *Charadrius marginatus* Vieillot, 1818 (White-fronted Plover), *Charadrius pallidus* Strickland, 1852 (Chestnut-banded Plover); *Charadrius pecuarius* Temminck, 1823 (Kittlitz's Plover); *Himantopus himantopus* (L., 1758) (Black-winged Stilt); *Vanelus armatus* (Burchell, 1822) (Blacksmith Lapwing)

Locality. Berg River, Western Cape Province, South Africa (Amin et al. 1999).

Family: Polymorphidae Meyer, 1931

Genus: *Andracantha* Schmidt, 1975

Andracantha tunitae (Weiss, 1914) Zdzitowiecki, 1989

Corynosoma tunitae Weiss, 1914

Host. *Microcarbo africanus* (Gmelin, 1789) (Long-tailed Cormorant) (Phalacrocoracidae).

Locality. Dyer Island, South Africa (Van Cleave 1937).

Notes. Van Cleave (1937) is mentioning that he looked at a large number of immature worms and he tentatively assigning them to *C. tunitae*.

Genus: *Arhythmorhynchus* Lühe, 1911

Skrjabinorhynchus Petrochenko, 1956

Arhythmorhynchus turbidus (Van Cleave, 1937) Golvan, 1994

Corynosoma turbidum Van Cleave, 1937

Host. *Phalacrocorax neglectus* (Wahlberg, 1855) (Bank Cormorant) (Phalacrocoracidae) (type host).

Locality. Dyer Island, South Africa (type locality) (Van Cleave 1937).

Type specimens. Holotype female (1737, 3) and one paratype female (1737, 1) in GNM. One paratype female (1737, 2) in the collection of H.J. Van Cleave, Urbana, Illinois, U.S.A.

Genus: *Bolbosoma* Porta, 1908***Bolbosoma capitatum* (von Linstow, 1880) Porta, 1908***Echinorhynchus capitatum* von Linstow, 1880*Bolbosoma physeteris* Gubanov, 1952 (*fide* Amin & Margolis, 1998)**Host.** *Ruvettus pretiosus* Cocco, 1833 (Oilfish) (Gempylidae).**Locality.** South Africa's West coast, Atlantic Ocean (Nunkoo et al. 2017).***Bolbosoma vasculosum* (Rudolphi, 1819) Porta, 1908***Echinorhynchus vasculosum* Rudolphi, 1819*Bolbosoma annulatus* Molin, 1858*B. aurantiacus* Risso, 1826*B. pellucidus* Leukart, 1828*B. serrani* Linton, 1888*B. thunni* Harada, 1935 (*fide* Petrochenko 1958)**Host.** *Thyrsites atun* (Euphrasen, 1791) (Snoek) (Gempylidae).**Locality.** South Africa's West and South coasts (Nunkoo et al. 2016).**Genus: *Corynosoma* Lühe, 1904 (*fide* Van Cleave, 1945)***Chentrosoma* Monticelli, 1905*Centrosoma* Lühe, 1912*Coryusoma* Railliet & Henry, 1907 (misprint)*Echinosoma* Porta, 1907***Corynosoma australe* Johnston, 1937***Corynosoma otariae* Morini & Boero, 1961**Host.** *Thyrsites atun* (Euphrasen, 1791) (Snoek) (Gempylidae).**Locality.** South Africa's West and South coasts (Nunkoo et al. 2016).

Host-Parasite List

Host	Parasite	Locality Country/ Reference
Class Aves		
Order Accipitriformes		
Family Accipitridae		
<i>Kaupifalco monogrammicus</i> (type host)	<i>Centrorhynchus sarebae</i> (Centrorhynchidae)	Makhado (Louis Trichardt), Limpopo Province, South Africa
Order Bucerotiformes		
Family Bucerotidae		
<i>Tockus leucomelas</i>	<i>Mediorhynchus taeniatus</i> (Gigantorhynchidae)	Limpopo Province, South Africa
Order Charadriiformes		
Family Charadriidae		
<i>Calidris ferruginea</i>	<i>Plagiorhynchus (Prosthorhynchus) cylindraceus</i> (Plagiorhynchidae)	Berg River, Western Cape Province, South Africa (Amin et al. 1999)
<i>Charadrius marginatus</i>	Unidentified plagiorhynchid acanthocephalan (Plagiorhynchidae)	Berg River, Western Cape Province, South Africa (Amin et al. 1999)
<i>Charadrius pallidus</i>	Unidentified plagiorhynchid acanthocephalan (Plagiorhynchidae)	Berg River, Western Cape Province, South Africa (Amin et al. 1999)
<i>Charadrius pecuarius</i>	<i>Plagiorhynchus (Prosthorhynchus) cylindraceus</i> (Plagiorhynchidae)	Berg River, Western Cape Province, South Africa (Amin et al. 1999)
	Unidentified plagiorhynchid acanthocephalan (Plagiorhynchidae)	Berg River, Western Cape Province, South Africa (Amin et al. 1999)
<i>Charadrius tricollaris</i>	<i>Plagiorhynchus (Prosthorhynchus) cylindraceus</i> (Plagiorhynchidae)	Berg River, Western Cape Province, South Africa (Amin et al. 1999)
<i>Vanellus armatus</i>	<i>Plagiorhynchus (Prosthorhynchus) cylindraceus</i> (Plagiorhynchidae)	Berg River, Western Cape Province, South Africa (Amin et al. 1999)
	Unidentified plagiorhynchid acanthocephalan (Plagiorhynchidae)	Berg River, Western Cape Province, South Africa (Amin et al. 1999)
Family Recurvirostridae		
<i>Himantopus himantopus</i>	Unidentified plagiorhynchid Acanthocephalan (Plagiorhynchidae)	Berg River, Western Cape Province, South Africa (Amin et al. 1999)
Order Galliformes		
Family Numididae		
<i>Numida meleagris</i> (type host)	<i>Mediorhynchus africanus</i> (previously identified as <i>Mediorhynchus gallinarum</i>) (Gigantorhynchidae)	Kruger National Park, Mpumalanga Province, South Africa (Junker and Boomker 2006); Vicinity of Petrus Steyn, Free State Province, South Africa (Davies et al. 2008); Musina, Limpopo Province, South Africa (Junker and Boomker 2007; Junker et al. 2008)
	<i>Mediorhynchus numidae</i> (Gigantorhynchidae)	Pretoria, Gauteng Province, South Africa (Oosthuizen and Markus 1967)
	<i>Mediorhynchus taeniatus</i> (Gigantorhynchidae)	Rooipoort farm, Kimberley, Northern Cape Province, South Africa (Crowe 1977)
Order Gruiformes		
Family Rallidae		
<i>Crecopsis egregia</i>	<i>Plagiorhynchus (Prosthorhynchus) cylindraceus</i> (Plagiorhynchidae)	Blouberg, Limpopo Province, South Africa
Order Passeriformes		
Family Turdidae		
<i>Turdus smithi</i> (type host)	<i>Mediorhynchus mokgalongi</i> (Gigantorhynchidae)	Polokwane, Limpopo Province, South Africa (type locality) (Smales et al. 2018)

Host	Parasite	Locality Country/ Reference
Order Strigiformes		
Family Strigidae		
<i>Bubo africanus</i>	<i>Centrorhynchus clitorideus</i> (Centrorhynchidae)	Zandriviervoort Farm, Polokwane, Limpopo Province, South Africa
Order Suliformes		
Family Phalacrocoracidae		
<i>Microcarbo africanus</i>	<i>Andracantha tunitae</i> (Polymorphidae)	Dyer Island, South Africa (Van Cleave 1937)
<i>Phalacrocorax neglectus</i> (type host)	<i>Arhythmorhynchus turbidus</i> (Polymorphidae)	Dyer Island, South Africa (type locality) (Van Cleave 1937)
Class Actinopterygii		
Order Cypriniformes		
Family Cyprinidae		
<i>Enteromius neefi</i>	<i>Acanthogyrus (Acanthosentis) phillipi</i> (Quadrigyridae)	Lingwe River, Venda, Limpopo Province, South Africa (Mashego 1988)
Order Perciformes		
Family Cichlidae		
<i>Oreochromis mossambicus</i>	<i>Acanthogyrus</i> sp. (Quadrigyridae)	Molepo dam, Limpopo Province, South Africa (Kunutu et al. 2013)
Family Gempylidae		
<i>Ruvettus pretiosus</i>	<i>Bolbosoma capitatum</i> (Polymorphidae)	South Africa's West coast, Atlantic Ocean (Nunkoo et al. 2017)
	<i>Rhadinorhynchus</i> sp. (Rhadinorhynchidae)	South Africa's West coast, Atlantic Ocean (Nunkoo et al. 2017)
<i>Thyrsites atun</i>	<i>Bolbosoma vasculosum</i> (Polymorphidae)	South Africa's West and South coasts (Nunkoo et al. 2016)
	<i>Corynosoma australe</i> (Polymorphidae)	South Africa's West and South coasts (Nunkoo et al. 2016)
	<i>Rhadinorhynchus cadenati</i> (Rhadinorhynchidae)	South Africa's West and South coasts (Nunkoo et al. 2016)
Family Sciaenidae		
<i>Argyrosomus japonicus</i> (type host)	<i>Neoechinorhynchus (Neoechinorhynchus)</i> <i>dorsovaginatus</i> (Neoechinorhynchidae)	Breed River Estuary, Western Cape Province, South Africa (type locality) (Amin and Christison 2005)
Order Pleuronectiformes		
Family Soleidae		
<i>Pegusa nasuta (Solea)</i> <i>bleekeri</i> (type host)	<i>Rhadinorhynchus capensis</i> (Rhadinorhynchidae)	Heuninges River estuary (Bray 1974)
	<i>Longicollum</i> sp. innom. (Pomphorhynchidae)	Klein River estuary (Bray 1974)
<i>Barnardichthys</i> <i>fulvomarginata</i> (type host)	<i>Longicollum chabanaudi</i> (Pomphorhynchidae)	False Bay, Western Cape Province (Dollfus and Golvan 1963)
Class Mammalia		
Order Afrosoricida		
Family Chrysochloridae		
<i>Chrysospalax trevelyani</i> (type host)	<i>Heptamegacanthus niekerki</i> (Oligacanthorhynchidae)	Nqadu Forest, Transkai, Eastern Cape Province, South Africa (Spencer Jones 1990)
Order Carnivora		
Family Felidae		
<i>Felis catus</i>	<i>Centrorhynchus</i> sp. (Centrorhynchidae)	Pretoria, Gauteng Province, South Africa (Baker et al. 1989)

Host	Parasite	Locality Country/ Reference
Family Herpestidae		
<i>Mungos mungo</i>	<i>Centrorhynchus</i> sp. (Centrorhynchidae)	Polokwane, Limpopo Province, South Africa.
Order Eulipotyphla		
Family Erinaceidae		
<i>Atelerix frontalis</i> (<i>Aethechinus frontalis</i>)	<i>Moniliformis kalahariensis</i> (Moniliformidae)	Mohlonong village and University of Limpopo, Limpopo Province, South Africa (Amin et al. 2014)
	<i>Moniliformis moniliformis</i> (Moniliformidae)	Hammanskraal, Gauteng Province, South Africa (Le Roux 1930)
Order Primates		
Family Galagidae		
<i>Otolemur crassicaudatus</i>	<i>Moniliformis</i> sp. (Moniliformidae)	Venda, Limpopo Province, South Africa
Order Rodentia		
Family Muridae		
<i>Gerbilliscus leucogaster</i>	<i>Moniliformis acomysi</i> (Moniliformidae)	Vyeboom village, Limpopo Province, South Africa
<i>Mastomys natalensis</i>	<i>Moniliformis acomysi</i> (Moniliformidae)	Bloemhof, Free State Province
<i>Mus minutoides</i>	<i>Moniliformis acomysi</i> (Moniliformidae)	Hoopstad, Free State Province, South Africa
Class Reptilia		
Order Squamata		
Family Varanidae		
<i>Varanus albigularis</i>	Oligacanthorhynchidae sp. (Oligacanthorhynchidae)	Tzaneen; Tolwe, Limpopo Province, South Africa

Results and discussion

A total of twenty-one species of acanthocephalans, from thirteen genera from ten families of seven orders, comprise this updated checklist of acanthocephalans in South Africa. Representatives of three of the four classes of acanthocephalans (Amin 2013) have been reported in South Africa, with only the Polyacanthocephala Amin, 1987 not having been recorded yet. The composition of reported acanthocephalan fauna shows that the Polymorphidae is the most represented family with five named species parasitic in marine fish and wild birds.

In South Africa, birds have the highest species richness of acanthocephalans to this date with nine named species (from five genera) and five records only identified to group level, followed by fish with eight named species (from six genera) and two species only identified to genus level, mammals with four named species (from two genera) and three species only identified to genus level and finally reptiles with a single species only identified to group level. No acanthocephalans have been reported in amphibians to date. During the current study 110 frog specimens belonging to 19 species were examined but none harboured any acanthocephalans. However, this forms a small part of the entire amphibian fauna of the country which includes 128 described frog species (Frost 2018).

Only a small fraction of the vertebrate fauna of South Africa has been surveyed for acanthocephalans and we expect that in future additional acanthocephalan species will be discovered and described. For example it is estimated that many of South Africa's marine fish parasites have yet to be discovered (Smit and Hadfield 2015). South Africa has an extremely rich biodiversity (Huntley et al. 2005), with nearly 8% of the world's known species of birds, 6% of mammal species and 5% of reptile species (Driver et al. 2012). Therefore we might expect a more diverse acanthocephalan fauna compared to that of Brazil which has 23 genera and 34 species (from only 119 fish species) (Santos et al. 2008) or Iran with 30 described species (Tavakol et al. 2015). The lower species richness reported for South Africa probably reflects sampling effort rather than the true diversity of the acanthocephalan fauna. Until more data are available it will not be possible to determine the true species richness of the South African acanthocephalan assemblage.

Acknowledgements

We would like to thank all the scientists who assisted us to access literature, particularly Geoffrey Allan Boxshall and David Gibson, and all those who helped AH with roadkill animals, especially Kgethedi Michael Rampedi and K David Kunutu. Special thanks are due to Ms Lorna Modiba (Librarian, University of Limpopo). Thanks also to Paul Skelton for his advice on fish species name changes. This work is based on the research supported by the South African Research Chairs Initiative of the Department of Science and Technology and National Research Foundation (NRF) of South Africa (Grant No 101054). Any opinion, finding and conclusion or recommendation expressed in this material is that of the author(s) and the NRF does not accept any liability in this regard.

References

- Amin OM (2013) Classification of the Acanthocephala. *Folia Parasitologica* 60(4): 273–305. <https://doi.org/10.14411/fp.2013.031>
- Amin OM, Canaris AG, Kinsella JM (1999) A taxonomic reconsideration of the genus *Plagiorhynchus* s. lat. (Acanthocephala: Plagiorhynchidae), with descriptions of South African *Plagiorhynchus* (*Prosthorhynchus*) *cylindraceus* from shore birds and *P. (P.) malayensis*, and a key to the species of the subgenus *Prosthorhynchus*. *Journal of Helminthological Society of Washington* 66: 123–132.
- Amin OM, Christison KW (2005) *Neoechinorhynchus* (*Neoechinorhynchus*) *dorsovaginatus* n. sp. (Acanthocephala: Neoechinorhynchidae) from the dusky kob *Argyrosomus japonicus* (Scienidae) on the southern coast of South Africa. *Systematic Parasitology* 61: 173–179. <https://doi.org/10.1007/s11230-005-3130-1>

- Amin OM, Evans P, Heckmann RA, El-Naggar AM (2013) The description of *Mediorhynchus africanus* n. sp. (Acanthocephala: Gigantorhynchidae) from galliform birds in Africa. *Parasitology Research* 112: 2897–2906. <https://doi.org/10.1007/s00436-013-3461-9>
- Amin OM, Heckmann RA, Halajian A, El-Naggar AM, Tavakol S (2014) Description of *Moniliformis kalahariensis* (Acanthocephala: Moniliformidae) from the South African Hedgehog, *Atelerix frontalis* (Erinaceidae) in South Africa. *Comparative Parasitology* 81(1): 33–43. <https://doi.org/10.1654/4664.1>
- Apps P (2012) *Smither's mammals of southern Africa: a field guide*. Penguin Random House, South Africa, 392 pp.
- Baker MK, Lange L, Verster A, Van der Plaat S (1989) A survey of helminths in domestic cats in the Pretoria area of Transvaal, Republic of South Africa. Part 1: The prevalence and comparison of burdens of helminths in adult and juvenile cats. *Journal of The South African Veterinary Association* 60(3): 139–142.
- Bray R (1974) Acanthocephala in the flatfish *Solea bleekeri* (Soleidae) from Cape Province, South Africa. *Journal of Helminthology* 48: 179–185. <https://doi.org/10.1017/S0022149X00022811>
- Clements JF, Schulenberg TS, Iliff MJ, Roberson D, Fredericks TA, Sullivan BL, Wood CL (2016). The eBird/Clements checklist of birds of the world: v2016. Downloaded from <http://www.birds.cornell.edu/clementschecklist/download/>
- Crowe TM (1977) Variation in intestinal helminth infestation of the helmeted guineafowl. *South African Journal of Wildlife Research* 7(1): 1–3.
- da Silva JM, Willows-Munro S (2016) A review of over a decade of DNA barcoding in South Africa: a faunal perspective. *African Zoology* 51(1): 1–12. <https://doi.org/10.1080/15627020.2016.1151377>
- Davies OR, Junker K, Jansen R, Crowe TM, Boomker J (2008) Age- and sex-based variation in helminth infection of helmeted guineafowl (*Numida meleagris*) with comments on Swainson's spurfowl (*Pternistis swainsonii*) and Orange River francolin (*Scleroptila levaillantoides*). *South African Journal of Wildlife Research* 38(2): 163–170. <https://doi.org/10.3957/0379-4369-38.2.163>
- Dollfus RP, Golvan YJ (1963) Extension a l'Afrique du sud de la distribution géographique du genre *Longicollum* S. Yamaguti 1935; *L. chabanaudi* n. sp. (Palaeacanthocephala, Pomphorhynchidae) parasite d'un *Barnardichthys*. (Soleidae). *Bulletin de la Société Zoologique de France* 88: 65–70.
- Driver A, Sink KJ, Nel JN, Holness S, Van Niekerk L, Daniels F, Jonas Z, Majiedt PA, Harris L, Maze K (2012) *National Biodiversity Assessment 2011: An assessment of South Africa's biodiversity and ecosystems. Synthesis Report*. South African National Biodiversity Institute and Department of Environmental Affairs, Pretoria.
- Froese R, Pauly D (Eds) (2017) *FishBase*. World Wide Web electronic publication. www.fishbase.org, version 10/2017.
- Frost DR (2018) *Amphibian Species of the World: an Online Reference*. Version 6.0. Electronic Database accessible at <http://research.amnh.org/herpetology/amphibia/index.html>. American Museum of Natural History, New York, USA.

- Hamer M (2013) A national strategy for zoological taxonomy (2013–2020). South African National Biodiversity Institute, Pretoria, 53 pp.
- Huntley PM, Van Noort S, Hamer M (2005) Giving increased value to invertebrates through ecotourism, *South African Journal of Wildlife Research* 35(1): 53–62.
- Junker K, Boomker J (2006) *Mediorhynchus gallinarum* (Acanthocephala: Gigantorhynchidae) in Helmeted guineafowls, *Numida meleagris*, in the Kruger National Park, South Africa. *Onderstepoort Journal of Veterinary Research* 73: 283–292.
- Junker K, Boomker J (2007) Helminths of guineafowls in Limpopo Province, South Africa, *Onderstepoort Journal of Veterinary Research* 74: 265–280.
- Junker K, Debusho L, Boomker J (2008) The helminth community of Helmeted guineafowls, *Numida meleagris* (Linnaeus, 1758), in the north of Limpopo Province, South Africa. *Onderstepoort Journal of Veterinary Research* 75: 225–235. <https://doi.org/10.4102/ojvr.v75i3.98>
- Khalil LF (1971) Check list of the helminth parasites of African freshwater fishes. Commonwealth Agricultural Bureaux, Farnham Royal, Slough, England, 80 pp.
- Khalil LF, Polling L (1997) Check list of the helminth parasites of African freshwater fishes. Department of Zoology/Biology, University of the North, Sovenga, South Africa, 185 pp.
- Kunutu KD, Luus-Powell WJ, Tavakol S, Halajian A, Hattingh HE, Geldenhuys G, Sara JR (2013) Fish health and parasites from Molepo dam, Limpopo province: new geographical and host records. 42nd PARSA Conference, Parys (South Africa), September 22–24, 45.
- Le Roux PL (1930) A new nematode (*Rictularia aethechini*, sp. nov.), a *Physaloptera* and an acanthocephalan from the hedgehog (*Aethechinus frontalis*), 16th Report of the Director of Veterinary Services and Animal Industry, Union of South Africa, 217–227.
- Mashego SN (1988) A new species of *Acanthosentis* Verma & Datta, 1929 (Acanthocephala: Quadrigyridae) from *Barbus neefi* in South Africa. *Annals of the Transvaal Museum* 34: 545–549.
- Nunkoo MAI, Reed CC, Kerwath SE (2016) Community ecology of the metazoan parasites of snoek *Thyrsites atun* (Euphrasen, 1791) (Perciformes: Gempylidae) off South Africa. *African Journal of Marine Science* 38(3): 363–371. <https://doi.org/10.2989/1814232X.2016.1216892>
- Nunkoo I, Weston MJ, Reed CC, van der Lingen CD, Kerwath S (2017) First account of the metazoan parasite fauna of oilfish *Ruvettus pretiosus* Cocco, 1829 (Perciformes: Gempylidae) in South African waters. *African Zoology* 52(4): 237–241. <https://doi.org/10.1080/15627020.2017.1411831>
- Oosthuizen JH, Markus, MB (1967) The haematozoa of South African birds. I: Blood and other parasites of two species of game birds. *Ibis* 109: 115–117. <https://doi.org/10.1111/j.1474-919X.1967.tb00009.x>
- Santos CP, Gibson DI, Tavares LE, Luque JL (2008) Checklist of acanthocephalan associated with the fishes of Brazil. *Zootaxa* 1938: 1–22.
- Skelton PH (2001) A complete guide to the freshwater fishes of Southern Africa, 2nd ed. Southern Book Publishers, Cape Town, 395 pp.
- Skelton PH (2016) Name changes and additions to the southern African freshwater fish fauna. *African Journal of Aquatic Science* 41(3): 1–7. <https://doi.org/10.2989/16085914.2016.1186004>

- Smit NJ, Hadfield KA (2015) Marine fish parasitology in South Africa: history of discovery and future direction. *African Zoology* 50(2): 79–92. <https://doi.org/10.1080/15627020.2015.1043644>
- Spencer Jones ME (1990) *Heptamegacanthus niekerki* n.g., n. sp. (Acanthocephala: Oligacanthorhynchidae) from the south-east African insectivore *Chrysospalax trevelyani* (Günther, 1857). *Systematic Parasitology* 15: 133–140. <https://doi.org/10.1007/BF00009991>
- Smales LR, Halajian A, Mokgawa MP, Luus-Powell WJ (2017) A new species of *Centrorhynchus* Lühe, 1911 (Acanthocephala: Centrorhynchidae) from the lizard buzzard *Kaupifalco monogrammicus* (Temminck) (Aves: Acciptridae) in South Africa. *Systematic Parasitology* 94: 423–430. <https://doi.org/10.1007/s11230-017-9710-z>
- Smales LR, Halajian A, Luus-Powell WJ, Tavakol S (2018) Acanthocephalans, including the description of a new species of *Mediorhynchus* (Giganthorhynchidae) and a redescription of *Centrorhynchus clitorideus* (Centrorhynchidae) from vertebrate hosts from South Africa. *Comparative Parasitology* 85(1): 95–106. <https://doi.org/10.1654/1525-2647-85.1.95>
- Tavakol S, Amin OM, Luus-Powell WJ, Halajian A (2015) The acanthocephalan fauna of Iran, a check list. *Zootaxa* 4033(2): 237–258. <https://doi.org/10.11646/zootaxa.4033.2.3>
- Van Cleave HJ (1937) Acanthocephala of the genus *Corynosoma* from birds of Dyer Island, South Africa. *Göteborgs Kungl. Vetenskaps och Vitterhets-Samhälles Handlingar. Ser. B.* 5(2): 1–6.
- Wilson DE, Reader DM (2005) *Mammal Species of the World* (3rd edn). Johns Hopkins University Press, Baltimore, 2142 pp.