

# *Ceratophysella* species from mushrooms in China (Collembola, Hypogastruridae)

Wanda Maria Weiner<sup>1</sup>, Zhijing Xie<sup>2,3</sup>, Yu Li<sup>4</sup>, Xin Sun<sup>2,4,5</sup>

**1** Institute of Systematics and Evolution of Animals, Polish Academy of Sciences, Slawkowska 17, 31-016 Kraków, Poland **2** Key laboratory of Wetland Ecology and Environment, Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences, Changchun 130102, China **3** University of Chinese Academy of Sciences, Beijing 100049, China **4** Engineering Research Center of Chinese Ministry of Education for Edible and Medicinal Fungi, Jilin Agricultural University, Changchun 130118, China **5** J.F. Blumenbach Institute of Zoology and Anthropology, University of Göttingen, 37073 Göttingen, Germany

Corresponding author: Xin Sun ([sunxin@iga.ac.cn](mailto:sunxin@iga.ac.cn))

---

Academic editor: L. Deharveng | Received 27 October 2018 | Accepted 21 January 2019 | Published 5 February 2019

<http://zoobank.org/DDE9A97D-0715-45E8-879F-8FCE235F8335>

---

**Citation:** Weiner WM, Xie Z, Li Y, Sun X (2019) *Ceratophysella* species from mushrooms in China (Collembola, Hypogastruridae). ZooKeys 822: 67–77. <https://doi.org/10.3897/zookeys.822.30880>

---

## Abstract

Four species of the genus *Ceratophysella* living on mushrooms are reported from China, including a new species, *Ceratophysella skarzynskii* Weiner & Sun, **sp. n.**, which is described from alpine mushrooms. The new species belongs to the *Ceratophysella* group of species with a dorsal chaetotaxy of type B and differs from the other species in a combination of characters. *Ceratophysella skarzynskii* **sp. n.** is distinguished by its small body size (maximum length 1.09 mm), number of peg-like s-chaetae (30–32) in the ventral sensory file, the trilobed apical vesicle of antennal segment IV, five modified chaetae on dens, and serrated dorsal chaetae. A key to the Chinese species of the genus has been provided.

## Keywords

*Ceratophysella skarzynskii* sp. n., key, new species, taxonomy

## Introduction

The genus *Ceratophysella* Börner, 1932 is distributed worldwide, having more than 130 species (Bellinger et al. 1996–2018). The main diagnostic characters for the genus are the pigmented body, 8 + 8 ocelli, body chaetae mostly differentiated into micro- and macrochaetae, an eversible integumental sac usually present between antennal segments III and IV, the ventral side of antennal segment IV with a sensory file often well-developed of short, erect, curved, and flattened at tips s-chaetae, unguiculus with broad basal lamella, furca well developed, mucro usually boat-like with a spoon-like apex and distinct lamella, and anal spines usually long and curved.

Until now, fourteen species of the genus *Ceratophysella* have been reported from China (Zhao 1992, Shen 1993, Liu et al. 1998, Zhao et al. 1997, Jia et al. 2010). As a common group of Collembola living on mushrooms, species have often caused significant economic damage in China (Wei 2002; Zhu 2012). Within a large collection of the mushroom Collembola in China, three known species, *C. communis* (Folsom, 1898), *C. denticulata* (Bagnall, 1941), *C. liguladorsi* (Lee, 1974), and the new species described here, *C. skarzynskii* sp. n., are reported in the present paper.

## Materials and methods

Specimens were collected by hand using a brush and stored in ethanol; they were then cleared in lactic acid and KOH, and mounted in Marc André II medium. Drawings and measurements were made using a phase contrast microscope LEICA DM2500 equipped with a camera lucida.

Mushroom species were determined by the third author, Yu Li.

## Abbreviations used in the descriptions:

<b>Abd.</b>	abdominal segments,	<b>s-chaetae</b>	sensorial chaetae on Th. and Abd.
<b>Ant.</b>	antennal segments,	<b>Th.</b>	thoracic segments,
<b>av</b>	apical vesicle	<b>VT</b>	ventral tube,
<b>AIIO</b>	sensory organ of Ant. III,	<b>IGA-CAS</b>	Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences;
<b>l.p.</b>	lateral processus on labial palp,	<b>ISEA-PAS</b>	Institute of Systematics and Evolution of Animals, Polish Academy of Sciences.
<b>ms</b>	s-microsetae (ms) (microsensillum),		
<b>or</b>	subapical organite		
<b>PAO</b>	postantennal organ,		
<b>S</b>	sensillum,		

Terminology for the descriptions follows that given in Fjellberg (1984, 1999), Babenko et al. (1994), and Thibaud et al. (2004).

## Taxonomy

### *Ceratophysella skarzynskii* Weiner & Sun, sp. n.

<http://zoobank.org/BA3F1CD5-D62C-4AFA-8450-26FB928994CA>

Figs 1–3, Table 1

**Type material.** *Holotype*: preadult male, China: Jilin: Changbai Mountains, alt. 2000 m, on *Russula* sp., leg. Xin Sun, 29 July 2015. *Paratypes*: 10 females and one juvenile, the same data as holotype. Type material: the holotype and 8 paratypes are housed in IGA-CAS, China, two paratypes in ISEA-PAS, Poland.

**Diagnosis.** Dorsal chaetotaxy of type B with serrated chaetae. Maximal length 1.09 mm. Antennal segment IV with bi- or trilobed apical vesicle and ventral sensory file with 30–32 peg-like s-chaetae. Dens with seven chaetae, five of them modified.


**Description.** Body length 0.9–1.09 mm (holotype: 1.07 mm). Body colour violet or blue in alive specimens, grey or grey-black in alcohol, ventrally pale. Granulation rather coarse, 10–14 granules between chaetae  $p_1$  on Abd. V (Yosii's parameter).

*Antennae.* Ant. IV with bilobed or trilobed apical vesicle (av), subapical organite (or), dorso-lateral microsensillum (ms), seven cylindrical, subequal sensilla (dorsal S0, S1–4, dorsolateral S7–8), ca. 30 small, peg-like sensilla and one subcylindrical sensillum in ventral sensory file (sensory rasp) (Fig. 2A, B). Ant. III-organ with two long (external) and two short (internal) curved sensilla (Fig. 2A). Microsensillum on ant. III present. Eversible sac between Ant. III–IV present (Fig. 2B). Ant. I with seven chaetae, Ant. II with 13 chaetae.

*Head.* Ocelli 8 + 8. Postantennal organ 1.5 times as large as single ocellus with four lobes of which the anterior pair is larger than the posterior pair (Fig. 2C). Accessory boss present (Fig. 2C).

*Labrum* with 5, 5, 4 chaetae, four prelabrals present. Head of maxilla of the *C. armata* type. Maxillary outer lobe with two sublobal hairs. Labium of the *C. armata* type, with five papillae (A–E) and six proximal chaetae. Guard chaetae  $a_1$ ,  $b_{1-2}$ ,  $d_2$ ,  $e_2$  and lateral processus (l.p.) as accessory papillae with short terminal sensillum. Guards  $b_{3-4}$ ,  $d_{3-4}$ , and  $e_{1-6}$  with long sensilla. Dorsal guards  $b_{3-4}$ ,  $d_{3-4}$ , and  $e_3$  distally expanded and flattened.

*Chaetotaxy.* Differentiation of dorsal chaetae into micro-/meso- and macrochaetae quite distinct (Figs 1A, 1B, 2D, 3C). Arrangement of chaetae on head typical for the genus, spine-like chaetae absent. Cephalic chaetae  $d_{2,4}$ ,  $v_2$ ,  $p_{3,4}$ ,  $g_{1,5}$ ,  $l_{01}$ ,  $l_{11}$  as macrochaetae. Dorsal chaetotaxy of B type (sensu Gisin 1947, Bourgeois and Cassagnau 1972, and Babenko et al. 1994) (Fig. 1B). Chaetae of medium length, pointed and serrated. Th. I with macrochaetae  $p_4$ , without  $p_2$ . Th. II–III with macrochaetae  $p_2$  (shifted forward),  $p_{5,6}$ ,  $m_5$ , chaetae  $m_4$  and  $m_5$  (Th. II with  $m_4$  and microsensillum ms), chaetae  $p_4$ ,  $m_6$  as sensorial chaetae s, chaetae  $a_2$  as long as  $a_3$ . Abd. I–III with macrochaetae  $p_{2,6}$ , sensorial chaetae s =  $p_5$ . Abd. IV with macrochaetae  $p_{1,3,6}$ , s-chaetae as  $p_4$ . Abd. V with macrochaetae  $p_{1,5}$ , 4 + 4 a-chaetae inside two macrochaetae p5 ( $a_{2,2}$  absent, chaeta  $a_5$



**Figure 1.** *Ceratophysella skarzynskii* sp. n. **A** Chaetotaxy of head and Th. I–III **B** Chaetotaxy of Abd. I–VI. Scale bars: 0.1 mm.

straight above  $p_3$ ) (Figs 1B, 3C). Body s-chaetae relatively long, but shorter than macrochaetae, only on Abd. V as long as macrochaetae  $p_{1,5}$  (Figs 1A, B, 3C).

*Tibiotarsi* I, II, III with 19, 19, 18 chaetae respectively, including one acuminate tenent hair  $A_1$  each, femora with 13, 13, 12 chaetae, trochanters with 7, 7, 7 chaetae, coxae with 3, 7, 8 chaetae, subcoxae II with 0, 3, 3 chaetae, subcoxae I with 1, 2, 3 chaetae. Claws with inner tooth and two pairs of lateral teeth. Empodial appendage with broad basal lamella and apical filament reaching slightly below inner tooth (ratio empodial filament: inner edge of claw = 0.47) (Fig. 3A). Ventral tube with 4 + 4 chaetae.

*Retinaculum* with 4 + 4 teeth. Furca well developed. Ratio dens + mucro: inner edge of claw III = 2.11: 1, ratio dens: mucro = 1.79: 1. Cuticular skeleton of furca well visible. Dens with uniform granulation and seven dorsal chaetae of which five are modified, two strongly thickened and three moderately so; basal macrochaeta longer than others chaetae, shorter than dens (3/5 of its length). Mucro boat-like with clear outer lamella (Fig. 3B, D).

*Anal spines* as long as inner edge of claw III slightly curved, situated on basal papillae, colourless (Figs 1B, 3C).

**Table 1.** Morphological characters for *C. skarzynski* sp. nov. and similar species: *C. denisana* Yosii, 1954 (Yosii 1956), *C. empodialis* Babenko, 1994, *C. longispina* (Tullberg, 1876) and *C. scotica* (Carpenter & Evans, 1899) after authors and Babenko et al. (1994).

Species/characters	<i>C. denisana</i>	<i>C. skarzynskii</i>	<i>C. empodialis</i>	<i>C. longispina</i>	<i>C. scotica</i>
Maximal body size (mm)	1.20	1.09	1.80	2.00	2.00
Yosii' parameter	14–16 (20)	10–14	8–13	12–16	13–15
Ant.IV: apical vesicle	trilobed	trilobed	simple/slightly bilobed	simple/slightly bilobed	simple
Ant.IV: number of peg-like chaetae in ventral file	>50	30–32	20	max. 15	15–20
Maxillar palp: number of sublobal hairs	2	2	2	2	1
Dorsal chaetae	smooth	serrated	rather smooth	serrated	rather smooth
Th.II–III: length of chaetae $s = p_4/p_3$	$p_4 > p_3$	$p_4 > p_3$	$p_4 > p_3$	$p_4 \approx p_3$	$p_4 \approx p_3$
Abd. V: chaetae $s = p_3/p_1$	$p_3 \approx p_1$	$p_3 \approx p_1$	$p_3 < p_1$	$p_3 \approx p_1$	$p_3 < p_1$
Tibiotarsial tenent hair	?	pointed	pointed	?	pointed
Empodial appendage : inner edge of claw	$\pm 1/2$	$\pm 1/2$	$\pm 1 1/4$	$\pm 1/2$	$\pm 1$
Empodial basal lamella : inner edge of claw	1/4	1/5	1/5	?	1/5
Lateral teeth of claw	basal 3 pairs	2 pairs	2 pairs	2 pairs	2 pairs
Chaetae on dens: total number/number of modified chaetae	7/5	7/5	7/2	7(8)/2	7/2

**Etymology.** The species is cordially dedicated to our colleague and friend Dr Dariusz Skarżyński, a prominent Polish specialist in Hypogastruridae, Collembola.


**Remarks.** The new species belongs to the *armata*-group of species, group B (Abd. tergum IV with  $p_1$  as macrochaeta) and subgroup B2 (Abd. tergum IV without chaeta  $p_3$ ) (Bourgeois and Cassagnau 1972). Among the species which could belong to this subgroup, *C. skarzynskii* is most similar to four species: *C. denisana* (Yosii, 1956), *C. empodialis* Babenko, 1994 (in Babenko et al. 1994), *C. longispina* (Tullberg, 1876), and *C. scotica* (Carpenter & Evans, 1899), due to the absence of transformed into spines, spine-like chaetae or spine-like integumentary protuberance on the head or Abd. V. They differ in the shape of the apical vesicle, the number of modified chaetae on the dens, the number of peg-like chaetae in the ventral sensory file on Ant. IV, length of empodial appendage, in the type of dorsal chaetae (serrated or smooth), and in the number of sublobal hairs on maxillary palp (one or two). A comparison of these species is presented in Table 1.

### *Ceratophysella communis* (Folsom, 1898)

*Acorutes communis* Folsom, 1898: 52.

Syn: *Hypogastrura yuasai* Yosii, 1954: Yosii 1960.

**Studied material.** China: Henan: Zhengzhou, on *Pleurotus ostreatus*, 7 specimens on slides and 20 in alcohol, leg. Xin Sun, 05 Dec 2014; China: Henan: Zhumadian, on




**Figure 2.** *Ceratophysella skarzynskii* sp. n. **A** ant. III and IV dorsal **B** ant. III and IV dorsal **C** PAO and eyes **D** macro-, microchaetae and s-chaeta. Scale bars: 0.01 mm.

*Pleurotus ostreatus*, 9 specimens on slides and 50 in alcohol, leg. ZhijingXie, 10 Nov 2017; China: Zhejiang: Pinghu, on *Pleurotus ostreatus*, 13 specimens on slides and 100 in alcohol, leg. Xin Sun, 07 Dec 2014; China: Sichuan: Qingchuan, on *Morchella esculenta*, 4 specimens on slides and 30 in alcohol, leg. Zhijing Xie, 19 Mar 2017.

### *Ceratophysella denticulata* (Bagnall, 1941)

*Achorutes denticulatus* Bagnall, 1941: 218.

Syn: *Achorutes armata* var. nov. *cuspidate* Axelson, 1905: Thibaud et al. 2004.



**Figure 3.** *Ceratophysella skarzynskii* sp. n. **A** tibiotarsus and claw III **B** dens and mucro **C** abd. V and VI **D** set of dens chaetae. Scale bars: 0.01 mm.

*Achorutes distinguendus* Bagnall, 1941: Thibaud et al. 2004.

*Hypogastrura (Ceratophysella) exilis* Yosii, 1956: Yosii 1965.

*Hypogastrura (Ceratophysella) afghanistanensis* Stach, 1963: Thibaud et al. 2004.

**Studied material.** China: Tibet: Lasa, on *Ganoderma* sp., 9 specimens on slides, leg. Weiping Xiong, 27 Mar 2015.

***Ceratophysella liguladorsi* (Lee, 1974)**

*Hypogastrura liguladorsi* Lee, 1974: 95.

**Studied material.** China: Zhejiang: Wuyi, on *Lentinus* sp., 11 specimens on slides and 80 in alcohol, leg. Xin Sun, 09 Dec 2014.

## Key to the Chinese species of *Ceratophysella*

In 2007 Wu and Yin proposed a key to the six species of the genus *Ceratophysella* known from China. Jia et al. (2010) proposed a list of 14 Chinese species of *Ceratophysella*. The present key includes 15 species. Some of the species are not sufficiently described, but the available characters given in the descriptions are sufficient to include them in the key. The characters for *C. adexilis* have been verified on type material, and for *C. communis* on fresh material from the type locality (Tokyo).

- 1 Abd. IV with p1 chaeta shorter than p2 chaeta (A-type) ..... 2
- Abd. IV with p1 chaeta longer than p2 chaeta (B-type)..... 9
- 2 Dens with 6 chaetae..... 3
- Dens with 7 chaetae..... 5
- 3 Unguiculus as long as 1/2–1 of internal edge of claw ..... 4
- Unguiculus very short, as long as 1/3, claw without teeth.... *C. zhangi* (Zhao, 1998)\*
- 4 Labial palp with 4 papillae (papilla C absent), macrochaetae rather short, Th. II with p4 = s longer than macrochaeta p5..... *C. succinea* (Gisin, 1949)
- Labial palp with 5 papillae (papillae A–E present), macrochaetae long, Th. II with p4 = shorter than macrochaeta p5..... *C. taiguensis* Jia, Skarżyński & Li, 2010
- 5 Body chaetae smooth ..... 6
- Body chaetae serrated..... 8
- 6 Dens with thickened chaetae..... 7
- Dens without thickened chaetae ..... *C. yinae* (Yue & Fu, 2000)
- 7 Four internal chaetae on dens thickened, Ant. IV with 8 dorsal sensilla ..... *C. baichengensis* Wu & Yin, 2007
- Two internal chaetae on dens thickened, Ant. IV with 7 dorsal sensilla..... *C. adexilis* Stach, 1964
- 8 Abd. V tergum with chaeta a2' present, Ant. IV with simple apical vesicle, Abd. IV tergum with p5=s equal to 1/2 macrochaetae..... *C. denticulata* (Bagnall, 1941)
- Abd. V tergum without chaeta a2, Ant. IV with bi- or trilobed apical vesicle, Abd. IV with chaeta p5=s equal to 3/4 macrochaetae..... *C. communis* (Folsom, 1898)
- 9 Abd. V tergum with cuticular projection or medial spines..... 10
- Abd. V tergum without such projection or spines ..... 11
- 10 Abd. V tergum with medial cuticular projection, dens with 7 normal chaetae, ventral sensory file (sensory rasp) with ca. 40 peg-like sensilla ..... *C. liguladorsi* (Lee, 1974)
- Abd. V tergum with chaetae p1 modified in spines, dens with 7 chaetae among which two modified, central sensory file (sensory rasp) with ca. 25–35 peg-like sensilla ..... *C. duplicispinosa* (Yosii, 1954)
- 11 Head with chaetae d5 and sd5 modified into spines..... 12
- Head without chaetae modified into spines..... 13

\* following the description in Tamura and Zhao 2008

- 12 Length of unguiculus as 1/3 of inner edge of claw, tibiotarsi with A1 (tenent hair) short (= 1/2 of inner edge of claw) ..... *C. xiaoi* (Tamura, 1998) (in: Tamura and Zhao 1998)
- Length of unguiculus as 1/2 of inner edge of claw, tibiotarsi with A1 (tenent hair) long (= length of inner edge of claw) .... *C. anshanensis* (Wu & Xie, 2007)
- 13 Head without a pair of cornea-like convexity ..... 14
- Head with a pair of cornea-like convexity ..... *C. sinensis* Stach, 1964
- 14 Dens with thickened chaetae, tibiotarsi with prolonged chaeta A1 as pointed tenent hair, claws with one internal tooth and two pairs of lateral teeth ..... *C. skarzynskii* sp. n.
- Dens without thickened chaetae, tibiotarsi without prolonged chaeta A1, claws without internal and lateral teeth ..... *C. flectochaeta* Lin & Xia, 1983

## Acknowledgements

We would like to express our thanks to Dariusz Skarżyński, Anatoly Babenko, and Louis Deharveng for their valuable and helpful advice. We are also grateful to Taizo Nakamori for collecting *C. communis* specimens in the type locality. The present study was mainly supported by funding from the National Basic Research Program of China (No. 2016YFA0602303-1), the National Natural Science Foundation of China (No.: 41571052, 41430857, 41811530086), the Science and Technology Development Plan Project of Jilin Province (No. 20160520051JH), the Postdoctoral Science Foundation of China (No. 2015M570281), the funding provided by the Alexander von Humboldt Foundation, the Youth Innovation Promotion Association, CAS, and the academic exchange program between Chinese and Polish Academies of Sciences.

## References

- Axelson WM (1905) Einigene Collembolen aus Finnland. Zoologischer Anzeiger 28: 788–794.
- Babenko AB, Chernova N, Potapov MB, Stebaeva SK (1994) Collembola of Russia and adjacent countries: Family Hypogastruridae. Moscow, Nauka, 336 pp.
- Bagnall RS (1941) Notes on British Collembola. VIII. The Entomology Monthly Magazine 77: 217–226.
- Bellinger P, Christiansen KA, Janssens F (1996–2018) Checklist of the Collembola of the World. <http://www.collembola.org> [Date of access: 22.08.2018]
- Börner C (1932) Apterygota. In: Brohmer P (Ed.) Fauna von Deutschland, Auflage 4. Leipzig, 136–143.
- Bourgeois A, Cassagnau P (1972) La différenciation du type ceratophysellien chez les Collemboles Hypogastruridae. Nouvelle Revue d'Entomologie 2: 271–291.
- Carpenter GH, Evans W (1899) Collembola and Thysanura of the Edinburgh District. Proceedings of the Royal Physical Society, Edinburgh 14: 221–266.

- Fjellberg A (1984) Maxillary structures in Hypogastruridae (Collembola). *Annales de la Société Royale Zoologique de Belgique* 114: 89–99.
- Fjellberg A (1999) The Labial Palp in Collembola. *Zoologischer Anzeiger* 237: 309–330.
- Folsom JW (1898) Japanese Collembola, Part I. *Bulletin of the Essex Institute* 29: 51–58. <https://doi.org/10.5962/bhl.part.14789>
- Gisin H (1947) Notes taxonomiques sur quelques espèces des genres *Hypogastrurab* et *Xenylla* (Collembola). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 20: 341–344.
- Gisin H (1949) Notes sur les Collemboles avec description de quatorze espèces et d'un genre nouveaux. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 22: 385–410.
- Jia J, Skarżyński D, Li Y (2010) *Ceratophysella taiguensis* sp. nov. (Collembola, Hypogastruridae) from China, with an annotated checklist of Chinese *Ceratophysella* Börner, 1932. *Zootaxa* 2644: 57–63. <https://doi.org/10.11646/zootaxa.2644.1.4>
- Lee BH (1974) Étude de la faune Coréenne des Insectes Collemboles II. Description de quatre espèces nouvelles de la famille Hypogastruridae. *Nouvelle Revue d'Entomologie* 4(2): 89–102.
- Lin S, Xia F (1983) A new species of the genus *Ceratophysella* (Collembola: Hypogastruridae). *Acta Entomologica Sinica* 26: 426–427.
- Liu Y, Hou D, Li Z (1998) A checklist of Collembola species from China. *Journal of Southwest Agricultural University* 20: 125–131.
- Shen XC (1993) A checklist of insects from Henan. Chinese Agriculture Science and Technological Press, 353 pp.
- Stach J (1963) Materials to the knowledge of Chinese Collembolan Fauna. *Acta zoologica cracoviensia* 10: 345–372.
- Stach J (1964) Materials to the knowledge of Chinese Collembolan fauna. *Acta zoologica cracoviensia* 9: 1–26.
- Tamura H, Zhao L (1998) Three species of Collembola from Yunnan, Southwest China (Apterygota). *Zoological Research* 19(2): 153–159.
- Thibaud JM, Schulz HJ, Gama Assalino MM da (2004) Hypogastruridae. In: Dunger W (Ed.) *Synopses on Palearctic Collembola*, Vol. 4. *Abhandlungen und Berichte des Naturkundemuseums, Görlitz* 75(2): 1–287.
- Tullberg T (1876) *Collembola borealia* (Nordiska Collembola). *Öfversigt af Kongliga Vetenskaps-Akademiens Förhandlingar* 33: 23–42.
- Wei W (2002) The harm and prevention of Collembola on Stropharia. *Guangxi Tropical Agriculture* 2: 17. [In Chinese]
- Wu D, Xie R (2007) New record of the genus *Mitchellania* Wray from China (Collembola, Hypogastruridae) with description of a new species. *Acta Zootaxonomica Sinica* 32: 287–289.
- Wu D, Yin W (2007) A new species of *Ceratophysella* Börner, 1932 (Collembola: Hypogastruridae), and description of Chinese specimens of *Ceratophysella succinea* Gisin, 1949. *The Pan-Pacific Entomologist* 83: 255–256. <https://doi.org/10.3956/0031-0603-83.3.255>
- Yosii R (1954) Springschwänze des Ozé-Naturschutzgebietes. *Scientific Researches of the Ozegahara Moor*, Tokyo, 777–830.
- Yosii R (1956) Monografie zur höhlen collembolen Japans. *Contributions from the Biological Laboratory Kyoto University* 3: 1–109.

- Yosii R (1960) Studies on the Collembolan genus *Hypogastrura*. American Midland Naturalist 64: 257–281. <https://doi.org/10.2307/2422661>
- Yosii R (1965) On some Collembola of Japan and adjacent countries. Contributions from the Biological Laboratory Kyoto University 19: 1–71.
- Yue Q, Fu R (2000) New records and new species of fresh water springtails from China (Collembola). Acta Entomologica Sinica 43: 394–402.
- Zhao L (1992) Collembola. In: Yin W (Ed) Subtropical soil animals of China. Science Press, Beijing, 414–457.
- Zhao L, Tamura H, Ke X (1997) Tentative checklist of collembolan species from China (Insecta). Publications of Itako Hydrobiological Station 9: 15–40.
- Zhu F (2012) Occurrence and comprehensive control measures of Collembola on edible fungi. Edible and medicinal mushrooms 20(5): 310–311.