

New Chinese record of the genus *Spinonychiurus* (Collembola, Onychiuridae), with the description of a new species

Xin Sun^{1,2}, Yu Li¹

1 Engineering Research Center of Chinese Ministry of Education for Edible and Medicinal Fungi, Jilin Agricultural University, Changchun 130118, China **2** Key laboratory of Wetland Ecology and Environment, Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences, Changchun 130102, China

Corresponding author: Yu Li (yuli966@126.com)

Academic editor: L. Deharveng | Received 25 April 2014 | Accepted 20 August 2014 | Published 8 September 2014

<http://zoobank.org/30AC763E-9C31-46E2-8CAA-8D6CD8461F60>

Citation: Sun X, Li Y (2014) New Chinese record of the genus *Spinonychiurus* (Collembola, Onychiuridae), with the description of a new species. ZooKeys 439: 19–26. doi: 10.3897/zookeys.439.7789

Abstract

A new collembolan species is described, *Spinonychiurus sinensis* sp. n., which has seven chaetae in the distal row of the tibiotarsi. It is placed in the genus *Spinonychiurus* due to two important characters: the two sub-segments on Abd. III sternum and the absence of d0 on the head. This is the first report of the genus *Spinonychiurus* in China. The diagnosis of *Spinonychiurus* is broadened and the key to the world species is provided.

Keywords

Taxonomy, Thalassaphorurini, tibiotarsi, key

Introduction

The genus *Spinonychiurus* was established by Weiner (1996) for the Scottish species *Onychiurus edinensis* Bagnall, 1935, and it was assigned to the tribe Onychiurini Börner, 1901 as having compound vesicles on the postantennal organ. Kaprus' and Tsalan (2009) revised the diagnosis characters of the genus and moved the genus into the tribe Thalassaphorurini Pomorski, 1998 by the structure of the furcal area and the distinct S-chaeta on the antennae. So far, only four species of the genus *Spinonychiurus* are

reported throughout the world: *S. edinensis*, *S. epaphius* Kaprus' & Tsalan, 2009 from Ukraine, *S. subedinensis* (Arbea & Jordana, 1985) from Spain and *S. vandeli* (Cassagnau, 1960) from France.

During our study on specimens collected from Changbai Mountain Range, we found a new species closest to the genus *Spinonychiurus* Weiner, 1996 but having 7 chaetae in the distal row of tibiotarsi (there are 11 in *Spinonychiurus* according to Kaprus' and Tsalan (2009)). In the present paper, we assign the new species to the genus *Spinonychiurus* with two important characters: two subsegments on Abd. III sternum and the absence of d0 on the head. The description of the new species and the broadened diagnosis of the genus are given below. An updated key to the species of the genus *Spinonychiurus* is provided.

Material and methods

Specimens were mounted in Marc André II solution, after clearing in lactic acid, and were studied using a Nikon Eclipse 80i microscope. Material is deposited in the Key Laboratory of Wetland Ecology and Environment, Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences, Changchun (NEIGAE).

Labial types are identified after Fjellberg (1999). Labium areas and chaetal nomenclature follow Massoud (1967) and D'Haese (2003). Chaetae on anal valves are recognised after Yoshii (1996). Furcal area is classified after Weiner (1996). Tibiotarsus chaetotaxy formula follows Deharveng (1983), and is expressed as: total number of chaetae (number of chaetae in row C, number of chaetae in row B, number of basal chaetae in rows A+T).

Abbreviations used in descriptions and figures:

Ant. – antennal segments, PAO – postantennal organ, Th. – thoracic segments, Abd. – abdominal segments, pso – pseudocellus, psp – pseudopore, psx – parapseudocelli, psx^m – unpaired parapseudocelli, ms – microsensillum, p-chaeta – chaeta of row p, S – S-chaeta, Sp – posterior S-chaeta on Abd. V, AIII O – sensory organ of antennal segment III.

The pseudocelli and pseudopores formulae are the number of pseudocelli or pseudopores per half-tergum or half-sternum. The S-chaetae formula is the number of S-chaetae per half-tergum or half-sternum from head to Abd. VI.

Systematics

Family Onychiuridae Börner, 1913

Tribe Thalassaphorurini Pomorski, 1998

Genus *Spinonychiurus* Weiner, 1996

Type species. *Onychiurus edinensis* Bagnall, 1935: 117.

Diagnosis. Postantennal organ oval, with numerous compound vesicles perpendicular to the long axis; clubs of AIIIIO smooth, ribbed or granulated; Ant. IV with differentiated S-chaetae; posterior pso on head present; chaeta d0 on head absent; S-chaetae on the body well marked; Abd. V tergum with or without spines; Abd. VI with one or two axial chaetae; anal spines present or absent; distal whorl of tibiotarsal chaetae as 7 or 11; Abd. III sternum divided into two subsegment; furcal rudiment as a finely granulated area with 4 small dental chaetae in two rows posteriorly, one manubrial row of chaetae present posteriorly to dental chaetae.

***Spinonychiurus sinensis* sp. n.**

<http://zoobank.org/BC4B8586-C7AE-4DB2-A1FB-E02B17174A8A>

Figs 1–2

Type material. Holotype: female; paratypes: 3 females and 1 male on slides - China, Jilin, Changbai Mountain Range (alt. 689 m, 43.0376°N, 128.9965°E), 3.Oct.2011, litter and soil, Berlese extraction, leg. Tang Xuguang.

Diagnosis. Pso formula as 32/133/33343 dorsally, absent ventrally, subcoxa 1 of legs I–III with 1, 1 and 1 pso respectively; psx formula as 11/000/122211^m ventrally, absent dorsally, subcoxa 1 of legs I–III with 1, 1 and 1 psx respectively; S-chaetae formula as 1/011/111021 dorsally, 11/000/000110 ventrally; sterna of Th. I, II, and III with 0+0, 2+2, 2+2 chaetae respectively; Abd. IV tergum with axial chaeta p0, Abd. V tergum with a0 and m0, Abd. VI tergum with a0; the distal row of tibiotarsi with 7 chaetae; male ventral organ absent; anal spines present, 0.8 times as long as inner edge of hind unguis.

Description. Body white in alcohol. Size 970–1200 µm in females, 900 µm in male; holotype: 1200 µm. Body subcylindrical, body sides parallel.

Pseudocellar formula: 32/133/33343 dorsally, absent ventrally (Figs 1A, G, 2A), subcoxa 1 of legs I–III with 1, 1 and 1 pso respectively. Parapseudocellar formula: 11/000/122211^m ventrally, absent dorsally (Figs 1A, 1G, 2A), subcoxa 1 of legs I–III with 1, 1 and 1 psx respectively. Pseudopore formula: 0/011/11110 dorsally, -/111/- ventrally (Figs 1A, G, 2A).

Head. Antennae short and distinctly segmented, as long as head. Length ratio of Ant. I: II: III: IV as about 1: 2: 2: 2. Ant. IV with two distinct thickened S-chaetae, subapical organite with globular apex; basolateral ms at approx. half length from base, above the second proximal row of chaetae (Fig. 1E). AIIIIO composed of 5 papillae, 5 guard chaetae, 2 sensory rods and 2 smooth sensory clubs, the inner bigger than the outer, and a lateral ms (Fig. 1E). Ant. II with 15 chaetae. Ant. I with 9 chaetae. Antennal base not marked (Fig. 1D). PAO with 12–13 compound vesicles (Fig. 1D). Dorsal cephalic chaeta d0 absent (Fig. 1A). 3+3 p-chaetae present between two inner posterior pso, p1 in line with others (Fig. 1E). Mandible with strong molar plate and 4 apical teeth. Maxilla bearing 3 teeth and 6 lamellae. Maxillary palp simple with 1 basal chaeta, without sublobal hair (Fig. 1B). Labral formula 4/1,4,2; labium with 6 proxi-




Figure 1. *Spinonychiurus sinensis* sp. n. **A** dorsal chaetotaxy of body **B** maxillary palp **C** labium **D** postantennal organ **E** antenna **F** distal part of leg III **G** chaetotaxy of Abd. II–VI sterna. Scale bars: 0.1 mm (A, E, G), 0.01 mm (B–D, F).




Figure 2. *Spinonychiurus sinensis* sp. n. **A** ventral chaetotaxy of head **B–C** central part of abdominal sternum IV **D** dorsal chaetotaxy of Abd. IV–VI. Scale bars: 0.1 mm (**A, D**), 0.01 mm (**B–C**).

mal, 4 basomedian (E, F, G, f) and 6 basolateral (a, b, c, d, e, e') chaetae (Fig. 2A); labial type A, papillae A–E respectively with 1, 4, 0, 3 and 3 guard chaetae (Fig. 1C). Head ventrally with 4+4 postlabial chaetae along ventral groove (Fig. 2A).

Body chaetotaxy. S-chaetae subcylindrical, apically rounded, 1/011/111021 dorsally, 11/000/000110 ventrally (Fig. 1A, G); subcoxae 2 of legs I, II and III with 0, 0, 1 S-chaeta respectively. Tiny and blunt ms, present on Th. II–III (Fig. 1A). Ordinary chaetae differentiated into meso- and macrochaetae, ratio Sp: m1: p1 on Abd. V tergum = 1: 1: 2.5. Th. I tergum with 7+7 dorsal chaetae. Th. II–Abd. III terga with 3+3 chaetae along axis respectively (Fig. 1A). Abd. IV tergum with one axial chaeta (p0), Abd. V tergum with two axial chaetae (a0 and m0), Abd. VI tergum with one axial chaeta (a0) (Figs 1A, 2D). Sterna of Th. I, II, and III with 0+0, 2+2, 2+2 chaetae respectively.

Table 1. Main diagnostic characters of world species of *Spinonychiurus*.

	<i>S. sinensis</i> sp. n.	<i>S. edinensis</i> *	<i>S. epaphius</i>	<i>S. subedinensis</i> **	<i>S. vandeli</i>
Dorsal pso formula	32/133/33343	32/223/1(?)1(?)3(?)43	5-6,5/4-5,8,8-10/9-13,9-12,9-14,9-14,7-10	34/233/44454	32/233/33343
Ventral pso formula	absent	absent	1/000/00000	1/000/01110	1/000/00000
Number of vesicles on PAO	12–13	14–16	13–16	18–22	22–25
Sensory clubs on AIIIIO	smooth	smooth	smooth	smooth	granulated
Chaetae in distal row of tibiotarsi	7	11***	11	11	11***
Male ventral organ	absent	?	present on ventral tube	present on ventral tube	absent
Spines on Abd. V	absent	present	absent	absent	absent
Anal spines	present	present	absent	present	present

* The details of *S. edinensis* follow Bagnall (1935) and Weiner (1996); ** the ventral pso formula has been verified by Javier Arbea (based on the type materials collected in University of Navarra); *** the number of chaetae in distal row of tibiotarsi in *S. edinensis* and *S. vandeli* has been verified by Louis Deharveng (based on the collections in Muséum National d'Histoire Naturelle).

Appendages. Subcoxa 1 of legs I–III with 4, 5 and 5 chaetae, subcoxa 2 with 1, 4 and 4 chaetae respectively. Tibiotarsi of legs I, II and III with 16 (1, 8, 7), 16 (1, 8, 7) and 15 (1, 7, 7) chaetae each (Fig. 1F). Unguis without teeth. Unguiculus approx. 0.6 times as long as inner edge of unguis, with inner basal lamella (Fig. 1F). Ventral tube with 4+4 basal and 7+7 distal chaetae. Furca reduced to a field of fine granulation with 4 small dental chaetae arranged in 2 rows posteriorly; only one manubrial row of chaetae posterior to dental chaetae (Figs 1G, 2B, C).

Genital plate with 9–11 chaetae in females, 32 chaetae in male. Male ventral organ absent. Anal valves with numerous acuminate chaetae; each lateral valve with a0, 2a1, 2a2; upper valves with chaetae a0, 2b1, 2b2, c0, 2c1, 2c2. Anal spines set on distinct papillae, 0.8 times as long as inner edge of hind unguis.

Derivatio nominis. Named for the first record of the genus *Spinonychiurus* in China.

Ecology. Found in the coniferous forest.

Discussion. The new species is closest to the genus *Spinonychiurus* in two important characters: two subsegments on Abd. III sternum and the absence of d0 on the head. However, it does not fit the current definition of the genus as proposed by Kaprus' and Tsalan (2009) in having 7 chaetae in the distal row of tibiotarsi instead of 11. The distal tibiotarsal chaetae have been verified to be an unstable character in the generic level (Sun et al. 2010; Sun et al. 2011; Sun and Zhang 2012; Sun et al. 2013), so we propose the placement of the new species in the genus *Spinonychiurus* and broaden its diagnosis accordingly. The main diagnostic characters of all known species of the genus are given in Table 1 and a key to these species is provided below.

Key to the known species of the genus *Spinonychiurus*

- 1 Spines on Abd. V tergum present ***S. edinensis* (Bagnall, 1935)**
 – Spines on Abd. V tergum absent..... **2**
 2 Chaetae in distal row of tibiotarsi as 7..... ***S. sinensis* sp. n.**
 – Chaetae in distal row of tibiotarsi as 11..... **3**
 3 Anal spines absent..... ***S. epaphius* Kaprus' & Tsalan, 2009**
 – Anal spines present **4**
 4 Dorsal pso formula as 34/233/44454.....
***S. subedinensis* (Arbea & Jordana, 1985)**
 – Dorsal pso formula as 32/233/33343..... ***S. vandeli* (Cassagnau, 1960)**

Acknowledgements

Thanks should be given to Louis Deharveng (France) and Javier Arbea (Spain) for their kind help in the examination of the species *S. edinensis*, *S. vandeli* and *S. subedinensis*. The present study was supported by the Fauna of Jilin Province, China, the National Natural Sciences Foundation of China (NO. 31301862, 31311130106, 31370532) and the Fund for Excellent Young Scholars of Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences (DLSYQ13003).

References

- Arbea JI, Jordana R (1985) Estudio ecológico de la colembofauna de los suelos del Macizo de Quinto Real (Pirineos Occidentales) y descripción de dos especies nuevas: *Anurida flaoelata* sp. n. y *Onychiurus subedinensis* sp. n. (Insecta, Collembola). Boletín de la Estación Central de Ecología 14(28): 57–80.
- Bagnall RS (1935) Contributions towards a knowledge of the Scottish Onychiuridae (Collembola), I. Scottish Naturalist 214: 111–117.
- Cassagnau P (1960) Faune française des Collemboles (XI): deux espèces nouvelles des Pyrénées centrales. Bulletin de la Société d'Histoire Naturelle de Toulouse 95: 3–4.
- Deharveng L (1983) Morphologie évolutive des Collemboles Neanurinae en particulier de la lignée Neanurienne. Travaux du Laboratoire d'Ecobiologie des Arthropodes Edaphiques, Toulouse 4(2): 1–63.
- D'Haese C (2003) Homology and morphology in Poduromorpha (Hexapoda, Collembola). European Journal of Entomology 101: 385–407. doi: 10.14411/eje.2003.060
- Fjellberg A (1999) The labial palp in Collembola. Zoologischer Anzeiger 237: 309–330.
- Kaprus' IJ, Tsalan JV (2009) New Collembola species from the floodplain forests of the Transcarpathian Lowland (Ukraine). Vestnik Zoologii 43(2): 173–178. doi: 10.2478/v10058-009-0008-8

- Massoud Z (1967) Monographie des Neanuridae, Collemboles Poduromorphes à pièces buccales modifiées. In: Delamare Deboutteville C, Rapoport EH (Eds) Biologie de l'Amérique Australe, III. CNRS, Paris, 7–399.
- Pomorski RJ (1998) Onychiurinae of Poland (Collembola: Onychiuridae). Genus 9: 1–201.
- Sun X, Chen JX, Deharveng L (2010) Six new species of *Thalassaphorura* (Collembola, Onychiuridae) from southern China, with a key to world species of the genus. Zootaxa 2627: 20–38.
- Sun X, Chen JX, Deharveng L (2011) Redefinition of the genus *Allonychiurus* Yoshii, 1995 (Collembola, Onychiuridae) with description of a new species from China. Zookeys 78: 27–41. doi: 10.3897/zookeys.78.977
- Sun X, Zhang F (2012) Two new species of *Onychiurus* (Collembola: Onychiuridae) from Eastern China. Journal of Natural History 46(31/32): 1895–1904. doi: 10.1080/00222933.2012.707236
- Sun X, Deharveng L, Wu DH (2013) Broadening the definition of the genus *Thalassaphorura* Bagnall, 1949 (Collembola, Onychiuridae) with a new aberrant species from China. ZooKeys 364: 1–9. doi: 10.3897/zookeys.364.6332
- Weiner WM (1996) Generic revision of Onychiurinae (Collembola: Onychiuridae) with a cladistic analysis. Annals of the Entomological Society of France (NS) 32: 163–200.
- Yoshii R (1996) Identity of some Japanese Collembola IV. “*Deuteraphorura*” Group of *Onychiurus*. Annals of the speleological research institute of Japan (Iwaizumi) 14: 1–15.