

Taxonomic interpretation of chromosomal and mitochondrial DNA variability in the species complex close to *Polyommatus (Agrodiaetus) dama* (Lepidoptera, Lycaenidae)

Nazar A. Shapoval^{1,2}, Vladimir A. Lukhtanov^{1,2}

1 Department of Karyosystematics, Zoological Institute of Russian Academy of Sciences, Universitetskaya nab. 1, St. Petersburg 199034, Russia **2** Department of Entomology, Faculty of Biology, St. Petersburg State University, Universitetskaya nab. 7/9, St. Petersburg 199034, Russia

Corresponding author: Nazar A. Shapoval (nazaret@bk.ru)

Academic editor: V. Kuznetsova | Received 13 September 2015 | Accepted 26 October 2015 | Published 19 November 2015

<http://zoobank.org/8BC2F1EA-6DFF-49DD-9EA4-CE7CC23AFF54>

Citation: Shapoval NA, Lukhtanov VA (2015) Taxonomic interpretation of chromosomal and mitochondrial DNA variability in the species complex close to *Polyommatus (Agrodiaetus) dama* (Lepidoptera, Lycaenidae). In: Lukhtanov VA, Kuznetsova VG, Grozeva S, Golub NV (Eds) Genetic and cytogenetic structure of biological diversity in insects. ZooKeys 538: 1–20. doi: 10.3897/zookeys.538.6559

Abstract

In this paper, by using combination of molecular and chromosomal markers, populations of *Polyommatus (Agrodiaetus) karindus* (Riley, 1921) from north-west and central Iran are analyzed. It has been found that taxon usually identified as *P. (A.) karindus* is represented in Iran by two geographically separated groups of individuals, strongly differentiated by their karyotypes and mitochondrial haplotypes. It is demonstrated that populations from NW Iran have the haploid chromosome number $n = 68$, while the haploid chromosome number of *P. (A.) karindus* from central Iran is found to be $n = 73$. Phylogenetic analysis revealed that these groups also differ by at least eight nucleotide substitutions in a 690 bp fragment of the mitochondrial *COI* gene and form separated groups of clusters in Bayesian inference tree. Thus, population entities from central Iran are described here as a new subspecies *Polyommatus (Agrodiaetus) karindus saravandi* ssp. n. Strong chromosomal and molecular differentiation are confirmed between *P. (A.) karindus* and its sister species, *P. (A.) dama* (Staudinger, 1892).

Keywords

COI, Iran, karyotype, molecular marker, chromosome number

Introduction

Agrodiaetus Hübner, 1822 is the most species-rich subgenus within the genus *Polyommatus* Latreille, 1804 (Talavera et al. 2013a, Lukhtanov et al. 2015a). It consists of approximately 130 species distributed in the western Palearctic (Vila et al. 2010, Lukhtanov et al. 2008, 2014, Vershinina and Lukhtanov 2010, Przybyłowicz et al. 2014, Lukhtanov and Tikhonov 2015). Today *Agrodiaetus* has become a model group in studies of speciation (Lukhtanov et al. 2005, 2015b), intraspecific differentiation (Dincă et al. 2013, Przybyłowicz et al. 2014, Lukhtanov et al. 2015a), and rapid karyotype evolution (Lukhtanov et al., 2005, Kandul et al. 2007). From the point of view of taxonomy, *Agrodiaetus* is a very complicated group. Many *Agrodiaetus* taxa display extremely similar phenotype (Hesselbarth et al. 1995) and, in contrast to other Lepidoptera taxa, genitalia offer only few distinctive features. Furthermore, many taxa represent allopatric populations which differ only slightly in morphology, and a conclusion on their status as distinct species or subspecies is controversial and can be misleading (Wiemers 2003, Lukhtanov et al. 2015a). This resulted in description of numerous polytypic species based on geographic distribution and classic morphological characters (Forster 1956, 1960a, b, 1961).

In particular, *Polyommatus (Agrodiaetus) dama* (Staudinger, 1892) was traditionally regarded as a polytypic species that included two subspecies: *Polyommatus (Agrodiaetus) dama dama* (Staudinger, 1892) (orig. comb. *Lycaena Dama*) and *Polyommatus (Agrodiaetus) dama karindus* (Riley, 1921) (orig. comb. *Lycaena dama* subsp. *karinda*). *P. (A.) dama dama* has only been found in South Anatolia (a few localities in Malatya, Maraş, and Mardin provinces (Turkey), while *P. (A.) dama karindus* distribution range is restricted to Zagros Mountains in Iran.

The karyotype studies of de Lesse (1957, 1959a, b, c, d, 1960a, b, 1961, 1962a, b, 1963a, b, 1964, 1966, 1968) revealed that *Agrodiaetus* species exhibit a wide diversity of karyotypes. Karyotyping may provide necessary diagnostic character for many *Agrodiaetus* species, and therefore become an important requirement for describing new taxa (de Lesse 1960a, b, Lukhtanov and Dantchenko 2002, 2003, Lukhtanov et al. 2008). Karyological investigations showed strong chromosomal differentiation between Turkish and Iranian populations of *P. (A.) dama* s. l.. De Lesse (1959a) described karyotype of *P. (A.) dama dama* from Kahramanmaraş and Olivier et al. (1999) confirmed his results from the type locality Malatya. It has an asymmetric karyotype with $n = 41$ chromosomes, about eleven of them are large, gradually decreasing in size, the others medium-sized; whereas the karyotype of Iranian taxon was determined as $n = 68$ (Wiemers 2003). Thus, on the basis of karyotype studies, *P. (A.) dama* s. l. was split into two species, *P. (A.) dama* and *P. (A.) karindus*, that can be characterized by species-specific haploid chromosome numbers.

However, the chromosome number of *P. (A.) karindus* was determined only for one population from NW Iran (Saqqez, Kordestan Province) (Wiemers 2003). Further investigations showed that Iranian species *P. (A.) karindus* has complicated genetic and

phylogeographic structure (Lukhtanov et al. 2015b). Here a combination of molecular mitochondrial (*COI*) and nuclear chromosomal (karyotype) markers are used to analyze different Iranian populations of *P. (A.) karindus*. Our study demonstrates that butterflies from central Iran strongly differentiated by their karyotypes and mitochondrial haplotypes from NW Iranian populations. Thus, population entities from central Iran are described here as a separate subspecies *Polyommatus (Agrodiaetus) karindus saravandi* ssp. n.

Material and methods

Specimens sampling

The butterflies were collected in the period of 2007–2014 in Iran (list of collected specimens is given in Table 1). In north-west Iran we collected material in two localities: 1) in the mountain range between Saqqez and Baneh (30–40 km SW of Saqqez), and 2) in the vicinity of Dare Dozdan (30–40 km W of Divandarreh). In central Iran we collected butterflies in the vicinity of Vennai (18 km W of Borujerd), in the vicinity of Saravand (15 km SE of Dorud), in the vicinity of Nahavand and in the vicinity of Darreh Takht (35 km NE of Dorud) (information about sampling localities is given in Figure 1 and Table 1).




Figure 1. Distribution ranges of *P. (A.) dama* (green circles), *P. (A.) karindus karindus* (red circles) and *P. (A.) karindus saravandi* (blue circles). The asterisk indicates the type locality of *P. (A.) karindus karindus*.

Table I. List of studied material (129 specimens) with information on karyotype (48 specimens) and *COI* sequences (54 specimens). Collectors: V. Lukhtanov (VL), N. Shapoval (NS) and A. Barabanov (AB).

Species	Sex	Sample ID	Chromosome number (n)	COI	GenBank number	Province	Locality and coordinates		Altitude	Date	Collectors
<i>karindus</i>	M	E391	70			Kordesstan	ca. 40 km SW Saqez	36°06'.18'N; 046°00'.27'E	1725 m	29 July 2004	VL
<i>karindus</i>	M	E399	68			Kordesstan	ca. 40 km SW Saqez	36°06'.18'N; 046°00'.27'E	1725 m	29 July 2004	VL
<i>karindus</i>	M	E400	68			Kordesstan	ca. 40 km SW Saqez	36°06'.18'N; 046°00'.27'E	1725 m	29 July 2004	VL
<i>karindus</i>	M	E402	68			Kordesstan	ca. 40 km SW Saqez	36°06'.18'N; 046°00'.27'E	1725 m	29 July 2004	VL
<i>karindus</i>	M	Z726	68	h02/GH5	KT582701	Kordesstan	ca. 40 km SW Saqez	36°05'.97'N; 045°59'.63'E	1720m	30 July 2007	VL & NS
<i>karindus</i>	M	Z727	68	h03/GH5	KT582702	Kordesstan	ca. 40 km SW Saqez	36°05'.97'N; 045°59'.63'E	1720m	30 July 2007	VL & NS
<i>karindus</i>	M	Z729	68			Kordesstan	ca. 40 km SW Saqez	36°05'.97'N; 045°59'.63'E	1720m	30 July 2007	VL & NS
<i>karindus</i>	M	Z749		h01/GH5	KT582703	Kordesstan	ca. 40 km SW Saqez	36°04'.82'N; 045°58'.88'E	1880m	31 July 2007	VL & NS
<i>karindus</i>	M	Z750		h01/GH5	KT582704	Kordesstan	ca. 40 km SW Saqez	36°04'.82'N; 045°58'.88'E	1880m	31 July 2007	VL & NS
<i>karindus</i>	M	Z753	68	h01/GH5	KT582705	Kordesstan	ca. 40 km SW Saqez	36°04'.82'N; 045°58'.88'E	1880m	31 July 2007	VL & NS
<i>karindus</i>	M	Z800	68	h01/GH5	KT582706	Kordesstan	ca. 40 km SW Saqez	36°04'.09'N; 045°58'.82'E	2050m	31 July 2007	VL & NS
<i>karindus</i>	M	Z809	68	h01/GH5	KT582707	Kordesstan	ca. 40 km SW Saqez	36°04'.09'N; 045°58'.82'E	2050m	31 July 2007	VL & NS
<i>karindus</i>	M	Z820	68	h01/GH5	KT582708	Kordesstan	ca. 40 km SW Saqez	36°04'.09'N; 045°58'.82'E	2050m	31 July 2007	VL & NS
<i>karindus</i>	M	Z843	68	h01/GH5	KT582709	Kordesstan	ca. 40 km SW Saqez	36°04'.64'N; 045°59'.16'E	1920-1950m	1 August 2007	VL & NS
<i>karindus</i>	M	Z845	69	h01/GH5	KT582710	Kordesstan	ca. 40 km SW Saqez	36°04'.64'N; 045°59'.16'E	1920-1950m	1 August 2007	VL & NS
<i>karindus</i>	M	W233	68			Kordesstan	ca. 40 km SW Saqez	36°03'.00'N; 045°58'.54'E	2027m	29 July 2009	VL & NS
<i>karindus</i>	M	W254	68			Kordesstan	ca. 40 km SW Saqez	36°03'.00'N; 045°58'.54'E	2027m	29 July 2009	VL & NS
<i>karindus</i>	M	W259	68			Kordesstan	ca. 40 km SW Saqez	36°03'.00'N; 045°58'.54'E	2027m	29 July 2009	VL & NS
<i>karindus</i>	M	W271	68			Kordesstan	ca. 40 km SW Saqez	36°04'.39'N; 045°59'.06'E	1869m	29 July 2009	VL & NS
<i>karindus</i>	M	W272	68			Kordesstan	ca. 40 km SW Saqez	36°04'.39'N; 045°59'.06'E	1869m	29 July 2009	VL & NS
<i>karindus</i>	M	W273	68			Kordesstan	ca. 40 km SW Saqez	36°04'.39'N; 045°59'.06'E	1869m	29 July 2009	VL & NS
<i>karindus</i>	M	W274	68			Kordesstan	ca. 40 km SW Saqez	36°04'.39'N; 045°59'.06'E	1869m	29 July 2009	VL & NS
<i>karindus</i>	M	W275	68			Kordesstan	ca. 40 km SW Saqez	36°04'.39'N; 045°59'.06'E	1869m	29 July 2009	VL & NS
<i>karindus</i>	M	W276	68			Kordesstan	ca. 40 km SW Saqez	36°04'.39'N; 045°59'.06'E	1869m	29 July 2009	VL & NS
<i>karindus</i>	M	W277	68			Kordesstan	ca. 40 km SW Saqez	36°04'.39'N; 045°59'.06'E	1869m	29 July 2009	VL & NS
<i>karindus</i>	M	W278	68			Kordesstan	ca. 40 km SW Saqez	36°04'.39'N; 045°59'.06'E	1869m	29 July 2009	VL & NS
<i>karindus</i>	M	W279	68			Kordesstan	ca. 40 km SW Saqez	36°04'.39'N; 045°59'.06'E	1869m	29 July 2009	VL & NS

Species	Sex	Sample ID	Chromosome number (n)	COI	GenBank number	Province	Locality and coordinates	Altitude	Date	Collectors
<i>karindus</i>	M	W280	68			Kordesstan	ca. 40 km SW Saqqez 36°04'39"N; 045°59'06"E	1869m	29 July 2009	VL & NS
<i>karindus</i>	M	W281	68			Kordesstan	ca. 40 km SW Saqqez 36°04'39"N; 045°59'06"E	1869m	29 July 2009	VL & NS
<i>karindus</i>	M	W282	68			Kordesstan	ca. 40 km SW Saqqez 36°04'39"N; 045°59'06"E	1869m	29 July 2009	VL & NS
<i>karindus</i>	M	W283	68			Kordesstan	ca. 40 km SW Saqqez 36°04'39"N; 045°59'06"E	1869m	29 July 2009	VL & NS
<i>karindus</i>	M	W340	h01/GH5	KT582732	Kordesstan	Dare Dozdan	35°52.05N; 046°33.03'E	2066m	30 July 2009	VL & NS
<i>karindus</i>	M	W341	h01/GH5	KT582733	Kordesstan	Dare Dozdan	35°52.05N; 046°33.03'E	2066m	30 July 2009	VL & NS
<i>karindus</i>	M	W342	h01/GH5	KT582734	Kordesstan	Dare Dozdan	35°52.05N; 046°33.03'E	2066m	30 July 2009	VL & NS
<i>karindus</i>	M	W354	h04/GH5	KT582737	Kordesstan	Dare Dozdan	35°52.05N; 046°33.03'E	2277m	31 July 2009	VL & NS
<i>karindus</i>	M	W355	h01/GH5	KT582735	Kordesstan	Dare Dozdan	35°52.05N; 046°33.03'E	2277m	31 July 2009	VL & NS
<i>karindus</i>	M	W361	h01/GH5	KT582736	Kordesstan	Dare Dozdan	35°52.05N; 046°33.03'E	2066m	31 July 2009	VL & NS
<i>karindus</i>	M	W366	h01/GH5	KT582738	Kordesstan	Dare Dozdan	35°52.05N; 046°33.03'E	2066m	31 July 2009	VL & NS
<i>karindus</i>	M	V069	h01/GH5	KT582739	Kordesstan	Dare Dozdan	35°51.30N; 046°42.60"E	2200m	27 July 2014	NS & AB
<i>karindus</i>	M	V070	h01/GH5	KT582740	Kordesstan	Dare Dozdan	35°51.30N; 046°42.60"E	2200m	27 July 2014	NS & AB
<i>karindus</i>	M	W370	73	h05/GH4	KT582722	Lorestan	Nahavand 34°02.57N; 048°20.22'E	2173m	2 August 2009	VL & NS
<i>karindus</i>	M	W371	73	h05/GH4	KT582723	Lorestan	Nahavand 34°02.57N; 048°20.22'E	2173m	2 August 2009	VL & NS
<i>karindus</i>	M	W372	73	h09/GH2	KT582724	Lorestan	Nahavand 34°02.57N; 048°20.22"E	2173m	2 August 2009	VL & NS
<i>karindus</i>	M	W373	73	h05/GH4	KT582725	Lorestan	Nahavand 34°02.57N; 048°20.22"E	2173m	2 August 2009	VL & NS
<i>karindus</i>	M	W374	h09/GH2	KT582726	Lorestan	Nahavand 34°02.57N; 048°20.22"E	2173m	2 August 2009	VL & NS	
<i>karindus</i>	M	W375	h09/GH2	KT582727	Lorestan	Nahavand 34°02.57N; 048°20.22"E	2173m	2 August 2009	VL & NS	
<i>karindus</i>	M	W376	h09/GH2	KT582728	Lorestan	Nahavand 34°02.57N; 048°20.22"E	2173m	2 August 2009	VL & NS	
<i>karindus</i>	M	W388	h08/GH2	KT582731	Lorestan	Nahavand 34°02.57N; 048°20.22"E	1950-2173m	3 August 2009	VL & NS	
<i>karindus</i>	M	W389	h05/GH4	KT582729	Lorestan	Nahavand 34°02.57N; 048°20.22"E	1950-2173m	3 August 2009	VL & NS	
<i>karindus</i>	M	W390	h09/GH2	KT582730	Lorestan	Nahavand 34°02.57N; 048°20.22"E	1950-2173m	3 August 2009	VL & NS	
<i>karindus</i>	M	W391			Lorestan	Nahavand 34°02.57N; 048°20.22"E	1950-2173m	3 August 2009	VL & NS	
<i>karindus</i>	M	W392			Lorestan	Nahavand 34°02.57N; 048°20.22"E	1950-2173m	3 August 2009	VL & NS	
<i>karindus</i>	M	U217			Lorestan	Nahavand, 34°02.92N; 48°20.40"E	2161 m	19 July 2011	VL & NS	
<i>karindus</i>	M	U218			Lorestan	Nahavand, 34°02.92N; 48°20.40"E	2161 m	19 July 2011	VL & NS	
<i>karindus</i>	M	U219			Lorestan	Nahavand, 34°02.92N; 48°20.40"E	2161 m	19 July 2011	VL & NS	
<i>karindus</i>	M	U220			Lorestan	Nahavand, 34°02.92N; 48°20.40"E	2161 m	19 July 2011	VL & NS	
<i>karindus</i>	M	U223			Lorestan	Nahavand, 34°02.92N; 48°20.40"E	2161 m	19 July 2011	VL & NS	

Species	Sex	Sample ID	Chromosome number (n)	COI	GenBank number	Province	Locality and coordinates	Altitude	Date	Collectors
<i>karindus</i>	M	U228				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U229				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U230				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U231				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U232				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U233				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U234				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U235				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U236				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U237				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U238				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U239				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U240				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U256				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U257				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U262				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U263				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U264				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U265				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U266				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U267				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U278				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U279				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U280				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	U281				Lorestān	Nahavand, 34°02.91'N; 48°21.08'E	2020 m	20 July 2011	VL & NS
<i>karindus</i>	M	Z381	73	h09/GH2	KT582691	Lorestān	W of Borujerd, Kuh-Garin mount. Vennnai, 33°53.89'N; 48°34.03'E	2150m	21 July 2007	VL & NS
<i>karindus</i>	M	Z382	73	h09/GH2	KT582692	Lorestān	W of Borujerd, Kuh-Garin mount. Vennnai, 33°53.89'N; 48°34.03'E	2150m	21 July 2007	VL & NS
<i>karindus</i>	M	Z396	73	h09/GH2	KT582693	Lorestān	W of Borujerd, Kuh-Garin mount. Vennnai, 33°53.89'N; 48°34.03'E	2150m	21 July 2007	VL & NS

Species	Sex	Sample ID	Chromosome number (n)	COI	GenBank number	Province	Locality and coordinates		Altitude	Date	Collectors
<i>karindus</i>	M	Z397	73	h10/GH2	KT582694	Lorestan	W of Borujerd, Kuh-e Garin mount.Vennnai, 33°53.89'N; 48°34.03'E	2150m	21 July 2007	VL & NS	
<i>karindus</i>	M	Z398	73	h09/GH2	KT582695	Lorestan	W of Borujerd, Kuh-e Garin mount.Vennnai, 33°53.89'N; 48°34.03'E	2150m	21 July 2007	VL & NS	
<i>karindus</i>	M	Z399	73	h11/GH1	KT582696	Lorestan	W of Borujerd, Kuh-e Garin mount.Vennnai, 33°53.89'N; 48°34.03'E	2150m	21 July 2007	VL & NS	
<i>karindus</i>	M	Z400	73	h10/GH2	KT582697	Lorestan	W of Borujerd, Kuh-e Garin mount.Vennnai, 33°53.89'N; 48°34.03'E	2150m	21 July 2007	VL & NS	
<i>karindus</i>	M	Z408	73	h10/GH2	KT582698	Lorestan	W of Borujerd, Kuh-e Garin mount.Vennnai, 33°53.89'N; 48°34.03'E	2150m	21 July 2007	VL & NS	
<i>karindus</i>	M	Z412	73	h11/GH1	KT582700	Lorestan	W of Borujerd, Kuh-e Garin mount.Vennnai, 33°53.89'N; 48°34.03'E	2150m	22 July 2007	VL & NS	
<i>karindus</i>	M	Z413				Lorestan	W of Borujerd, Kuh-e Garin mount.Vennnai, 33°53.89'N; 48°34.03'E	2150m	22 July 2007	VL & NS	
<i>karindus</i>	M	Z416		h09/GH2	KT582699	Lorestan	W of Borujerd, Kuh-e Garin mount.Vennnai, 33°53.89'N; 48°34.03'E	2150m	22 July 2007	VL & NS	
<i>karindus</i>	M	V331				Lorestan	W of Borujerd, Kuh-e Garin mount.Vennnai, 33°53.89'N; 48°34.03'E	2150m	2 August 2014	NS & AB	
<i>karindus</i>	M	V335				Lorestan	W of Borujerd, Kuh-e Garin mount.Vennnai, 33°53.89'N; 48°34.03'E	2150m	2 August 2014	NS & AB	
<i>karindus</i>	M	V336				Lorestan	W of Borujerd, Kuh-e Garin mount.Vennnai, 33°53.89'N; 48°34.03'E	2150m	2 August 2014	NS & AB	
<i>karindus</i>	M	W061	73,74,75	h12/GH1	KT582711	Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	21 July 2009	VL & NS	
<i>karindus</i>	M	W062	ca73			Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	22 July 2009	VL & NS	
<i>karindus</i>	M	W063	71	h12/GH1	KT582712	Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	22 July 2009	VL & NS	
<i>karindus</i>	M	W064	73	h12/GH1	KT582713	Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	22 July 2009	VL & NS	
<i>karindus</i>	M	W065	ca73	h12/GH1	KT582714	Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	22 July 2009	VL & NS	
<i>karindus</i>	M	W072	ca73	h12/GH1	KT582715	Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	22 July 2009	VL & NS	
<i>karindus</i>	M	W073				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	22 July 2009	VL & NS	
<i>karindus</i>	M	W074				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	22 July 2009	VL & NS	
<i>karindus</i>	M	W075				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	22 July 2009	VL & NS	

Species	Sex	Sample ID	Chromosome number (n)	COI	GenBank number	Province	Locality and coordinates	Altitude	Date	Collectors
<i>karindus</i>	M	W081				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	22 July 2009	VL & NS
<i>karindus</i>	M	W082				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	22 July 2009	VL & NS
<i>karindus</i>	M	W083				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	22 July 2009	VL & NS
<i>karindus</i>	M	W084				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	22 July 2009	VL & NS
<i>karindus</i>	M	W085				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	22 July 2009	VL & NS
<i>karindus</i>	M	W086				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	22 July 2009	VL & NS
<i>karindus</i>	M	W087				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	22 July 2009	VL & NS
<i>karindus</i>	M	W093				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	23 July 2009	VL & NS
<i>karindus</i>	M	W094				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	23 July 2009	VL & NS
<i>karindus</i>	M	W095				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	23 July 2009	VL & NS
<i>karindus</i>	M	W096				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2070m	23 July 2009	VL & NS
<i>karindus</i>	M	W377	h12/GH1	KT582716		Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2100–2250m	3 August 2009	VL & NS
<i>karindus</i>	M	W378	h12/GH1	KT582717		Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2100–2250m	3 August 2009	VL & NS
<i>karindus</i>	M	W379	h06/GH3	KT582718		Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2100–2250m	3 August 2009	VL & NS
<i>karindus</i>	M	W380	h12/GH1	KT582719		Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2100–2250m	3 August 2009	VL & NS
<i>karindus</i>	M	W381	h12/GH1	KT582720		Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2100–2250m	3 August 2009	VL & NS
<i>karindus</i>	M	W382	h07/GH3	KT582721		Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2100–2250m	3 August 2009	VL & NS
<i>karindus</i>	M	W383				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2100–2250m	3 August 2009	VL & NS
<i>karindus</i>	M	W386				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2100–2250m	3 August 2009	VL & NS
<i>karindus</i>	M	W387				Lorestan	Saravand, 33°22.39'N; 49°10.25'E	2100–2250m	3 August 2009	VL & NS
<i>karindus</i>	M	U168	h12/GH1	KT582741		Lorestan	Dareh Takht, 33°21.19'N; 49°22.34'E	2000–2100 m	18 July 2011	VL & NS
<i>karindus</i>	M	U178	h12/GH1	KT582742		Lorestan	Dareh Takht, 33°21.19'N; 49°22.34'E	2000–2100 m	18 July 2011	VL & NS
<i>karindus</i>	M	U179	h12/GH1	KT582743		Lorestan	Dareh Takht, 33°21.19'N; 49°22.34'E	2000–2100 m	18 July 2011	VL & NS
<i>karindus</i>	F	U169	h06/GH3	KT582744		Lorestan	Dareh Takht, 33°21.19'N; 49°22.34'E	2000–2100 m	18 July 2011	VL & NS

In addition, we used the following sequences from GenBank:

COI: AY557145 *P. (A.) karindus* (h01/GH5); AY557007 *P. (A.) damii*; AY56887 *P. (A.) birunii*.

Fresh (not worn) adult males were used to investigate the karyotypes. After capturing a butterfly in the field, it was placed in a glassine envelope for 1–2 hours to keep it alive until processed. Butterflies were killed by pressing the thorax. Testes for karyotype analysis were removed from the abdomen and placed into a 0.5 mL vial with a freshly prepared fixative (ethanol and glacial acetic acid 3:1). Then each wing was carefully removed from the body using forceps and placed into glassine envelope. The wingless body was placed into a plastic, 2 mL vial with pure 100% ethanol (for DNA analysis). Each vial with ethanol has already been numbered. This ID number was also used to label a vial with the fixative and a glassine envelope, in which the wings are preserved. Thus, each specimen was individually fixed. All collected specimens are kept in the Zoological Institute of the Russian Academy of Science (St. Petersburg) (ZIN RAS). All the testes are kept in the Department of Karyosystematics (ZIN RAS).

Chromosome preparation and karyotyping

Testes were stored in the fixative for 1–12 months at 4 °C. Then the gonads were stained in 2% acetic orcein for 30–60 days at 18–20 °C. Chromosome preparations were obtained as previously described (Talavera et al. 2013b). Different stages of male meiosis were examined by using a light microscope (Amplival, Carl Zeiss). An original two-phase method of chromosome analysis was used (Lukhtanov et al. 2006).

DNA Extraction and Sequencing

A fragment of the mitochondrial cytochrome *c* oxidase subunit I gene (first 690 positions) served as a mitochondrial molecular marker. Thoracic muscles and first abdominal segments were used for DNA extraction. The segments were homogenized in CTAB buffer and digested with proteinase K (10 mg/mL) for three hours at 60 °C. DNA was purified through successive ethanol precipitations and stored in dd H₂O at -20 °C.

For DNA amplification of *COI* we used primers K698 and Nancy (Caterino and Sperling 1999). PCR reactions (50 µl) contained 10 pmol each of forward and reverse primer, 1 mM dNTPs, 10x PCR Buffer (0.01 mM Tris-HCl, 0.05 M KCl, 0.1% Triton X-100: pH 9.0), 1 unit Taq DNA Polymerase (Fermentas), 5 mM MgCl₂ and were conducted using the following profile: initial 4 min denaturation at 94 °C and 30 cycles of 30 sec denaturation at 94 °C, 1 min annealing at 55 °C, 1 min extension at 72 °C and 5 min final elongation at 72 °C. PCR products were analyzed on 1.5% agarose gel, and purified using GeneJET PCR purification kit (Fermentas). Sequencing of double-stranded product was carried out at the Research Resource Center for Molecular and Cell Technologies (St. Petersburg State University).

Sequence alignments and phylogeny inference

The sequences were edited and aligned using CHROMAS 2.4.3 (<http://www.techne-lysimium.com.au/>), Geneious 8.1.6 (Kearse et al. 2012), and BioEdit 7.0.3 (Hall 2011) software. The alignment was unambiguous, as all the sequences were of equal length and included no insertions/deletions. Primer sequences were cropped. This resulted in final alignment of 690 bp *COI* fragments. The analysis involved *COI* sequences inferred from 54 *P. (A.) karindus* specimens. Additional sequences of the *P. (A.) dama* (accession number AY557007) and *P. (A.) karindus* (accession number AY557145) were found in GenBank (Wiemers 2003) and were included into analysis, since these sequences completely overlapped with our fragment. We used sequence of *P. (A.) birunii* (Eckweiler & ten Hagen, 1998) (accession number AY556558) as an outgroup to root the phylogeny (according to available data, this species does not belong to the group closely related to *P. (A.) dama*). Thus, the final analysis included in total 57 *COI* sequences. A Bayesian approach for estimating phylogeny was used. Bayesian analyses were performed using the program MrBayes 3.2 (Ronquist et al. 2012), with the nucleotide substitution model GTR+G+I as suggested by jModelTest (Posada 2008). TRACER, v. 1.4 was used for summarizing the results of Bayesian phylogenetic analyses (<http://beast.bio.ed.ac.uk/Tracer>). A maximum-parsimony haplotype network was built using TCS v. 1.21, with a 99% parsimony connection limit (Clement et al. 2000).

Results

Analysis of karyotypes

Meiotic karyotypes were studied in 48 specimens of *P. (A.) karindus* from different Iranian localities. Depending on karyotypes and localities, 2 groups of individuals can be distinguished (Table 1 and see below).

Group I (*P. (A.) karindus* from NW Iran)

The haploid chromosome number $n = 68$ was found in meiotic metaphase I (MI) and meiotic metaphase II (MII) cells. The MI karyotype displayed 5 large bivalents in the center of metaphase plate and 63 smaller bivalents in the periphery (Fig. 2A).

Group II (will be described below as *P. (A.) karindus saravandi* from central Iran)

The haploid chromosome number $n = 73$ was found in meiotic MI and MII cells of studied individuals (Fig. 2B). The MI karyotype was strongly asymmetric with 5–6 larger bivalents in the center of the MI plate and 67–68 smaller bivalents in the periphery.




Figure 2. Male meiosis I karyotypes of: **A** *P. (A.) karindus karindus*, sample E399, Iran, Kordestan, 40 km SW Saqpez, 1800–1900 m, 2004.VII.29, V. Lukhtanov leg., n = 68 **B** *P. (A.) karindus saravandi*, sample W372, Iran, Nahavand 34°02.57'N; 048°20.22'E, 2173m, 2009.VIII.02, V. Lukhtanov & N. Shapoval leg., n = 73. Scale bar = 10 μ m.

Phylogenetic analysis of molecular data

A Bayesian inference recovered *P. (A.) karindus* as a strongly supported monophyletic clade characterized by a specific set of fixed nucleotide substitutions (Fig. 3). Specimens of *P. (A.) karindus* were divided into several clusters: one cluster united specimens of *P. (A.) karindus* collected in north–west Iran (Fig. 3, GH5, highlighted in pink) and the others (Fig. 3, GH1–GH4, highlighted in blue) included specimens of central Iran populations (described here as a novel subspecies *P. (A.) karindus saravandi*). Most parsimonious *COI* haplotype network demonstrated similar pattern (Fig. 4). *P. (A.) dama* differs from *P. (A.) karindus* by at least 20 fixed nucleotide substitutions. Specimens of *P. (A.) karindus* form several haplotypes clustered in five different haplogroups. In general, composition of each haplogroup reflects geographical distribution of butterflies. Thus, majority of the specimens from easternmost (Saravand and Darreh Takht) and central west (Vennai, Nahavand) localities form two distinct haplogroups: GH1 and GH2. Nevertheless, two specimens from Vennai (approx. 80 km NW from Saravand) were found to have mitochondrial haplotype similar to that in easternmost populations, which has led to the suggestion that there is no complete isolation (reproductive or/and geographical) between population from Vennai and easternmost populations. The third haplogroup (GH3) consists of only three specimens, which were collected in Saravand and Darreh Takht. Interestingly, the third haplogroup differs drastically (by 10–12 fixed nucleotide substitutions) from the haplotypes, which comprise all other specimens from Saravand and Darreh Takht (group GH1). The fourth haplogroup (GH4) unites four specimens from Nahavand. Finally, all the haplotypes found in NW Iran constituted a subset of the distinct haplogroup (GH5). Thus, most parsimonious *COI* haplotype network reflects complex phylogeographic pattern of *P. (A.) karindus*.



Figure 3. The Bayesian tree of *Polyommatus (Agrodiætus) dama* and *Polyommatus (Agrodiætus) karindus* based on analysis of the cytochrome c oxidase subunit I gene from 57 specimens. Numbers at nodes indicate Bayesian posterior probability. *Agrodiætus karindus karindus* and *Agrodiætus karindus saravandi* clusters highlighted in pink and blue respectively.




Figure 4. *COI* Haplotype analysis. **A** geographical distribution of haplogroups. Number of studied individuals sharing the same haplogroup is given in parentheses **B** most parsimonious *COI* haplotype network; h01–h12 are *COI* haplotypes; GH1–GH5 are *COI* haplogroups. Number of studied individuals sharing the same haplotype is given in parentheses.

Discussion

We have found that a taxon usually identified as *P. (A.) karindus* is represented in Iran by two geographically separated groups of individuals. The first group unites specimens collected in NW Iran, while the second group comprises specimens from central Iran. The representatives of these groups have different chromosome numbers, $n = 68$ and $n = 73$ respectively. They also have at least eight fixed nucleotide differences in 690 bp fragment of mitochondrial *COI* gene. The first group is monophyletic with respect to

both *COI* gene and karyotype ($n = 68$). The second group has complicated genetic structure, comprises several differentiated populations and is paraphyletic with respect to the *COI* gene. Despite this gene paraphyly, it appears as a clearly monophyletic group with respect to its karyotype ($n = 73$). Thus, the NW and central Iranian groups are differentiated by at least five fixed chromosome fusions/fissions. Fixed chromosome differences are often considered as characters associated with reproductive isolation (King 1993). From this point of view, the NW and central Iranian groups could be theoretically treated as a different species. However, our recent studies on *Agrodiaetus* demonstrated that multiple chromosome fusions and fissions did not block fertility in chromosomal hybrids (Lukhtanov et al. 2015b). In other words, differentiation by five fixed chromosome rearrangements would not guarantee impossibility of blending populations together when they occur in sympatry. Thus, NW and central Iranian groups of populations should be considered as a subspecies rather than separate species.

Since *Polyommatus (Agrodiaetus) karindus* (Riley, 1921) (orig. comb. *Lycaena dama* subsp. *karinda*) was described from NW Iran (type locality is “Harir, Karind, and Karind Gorge, N.W. Persia” according to original description, and “N.W. Persia, Karind Gorge, 6000 ft” according to lectotype designation made by Bálint (1999) (not from central Iran), the name *P. karindus karindus* should be attributed to the NW Iranian group of populations. The formal description and naming of the central Iranian group is provided below.

Description of the novel taxon

Polyommatus (Agrodiaetus) karindus saravandi ssp. n.

<http://zoobank.org/ADC4F3C8-B804-4869-955A-9F7C7164B0C5>

Fig. 1 – map, Fig. 2B karyotype, Figs 3–4 phylogeny, Fig. 5 – Holotype of *P. (A.) karindus saravandi*, Fig. 6 A, B – Underside and upperside of the male and female wings

Holotype. ♂. Forewing length 34.0 mm. Iran, Lorestan province, Zagros Mt., vicinity of Saravand village, 33°22.39'N; 49°10.25'E, 2070 m, 22.07.2009. N. Shapoval and V. Lukhtanov leg. In the Zoological Institute of the Russian Academy of Sciences (St. Petersburg). Specimen field code W064, GenBank code for mitochondrial cytochrome *c* oxidase subunit I (*COI*) gene (partial cds) is KT582713.

Paratypes. 87 ♂♂, field codes W061, W062, W063, 21.07.2009; W065, W072, W073, W074, W075, W081, W082, W083, W084, W085, W086, W087, the same locality, date and collectors as the holotype. Field codes W093, W094, W095, W096 23.07.2009, the same locality and collectors as the holotype. Field codes W377, W378, W379, W380, W381, W382, W383, W386, W387 03.08.2009, the same locality and collectors as the holotype. Field codes W370, W371, W372, W373, W374, W375, W376, Iran, Lorestan province, Zagros Mt., vicinity of Nahavand village, 34°02.57'N; 048°20.22'E, 2170 m, 02.08.2009, the same collectors as the holotype. Field codes W388, W389, W390, W391, W392, Iran, Lorestan province, Zagros Mt., vicinity



Figure 5. Holotype of *P. (A.) karindus saravandi*, sample W064. Upperside (left) and underside (right) of the male wings.




Figure 6. Underside and upperside of the *P. (A.) karindus saravandi* ssp. n. wings. **A** upperside (left) and underside (right) of the male wings **B** upperside (left) and underside (right) of the female wings.

of Nahavand village, 34°02.57'N; 048°20.22'E, 2170 m, 02.08.2009, the same collectors as the holotype. Field codes U217, U218, U219, U220, U223, Iran, Lorestan province, Zagros Mt., vicinity of Nahavand village, 34°02.57'N; 048°20.22'E, 2170 m, 19.07.2011, the same collectors as the holotype. Field codes U228, U229, U230, U231, U232, U233, U234, U235, U236, U237, U238, U239, U240, U256, U257, U262, U263, U264, U265, U266, U267, U278, U279, U280, U281, Iran, Lorestan province, Zagros Mt., vicinity of Nahavand village, 34° 02.92'N; 48° 20.40'E, 2160 m, 20.07.2011 the same collectors as the holotype. Field codes Z381, Z382, Z396, Z397, Z398, Z399, Z400, Z408, Iran, Lorestan province, Zagros Mt., W of Borujerd, Kuh-e Garin mount mount., Vennnai, 33°53.89'N; 48°34.03'E, 2150 m, 21.07.2007, the same collectors as the holotype. Field codes Z412, Z413, Z416, Iran, Lorestan province, Zagros Mt., W of Borujerd, Kuh-e Garin mount., Vennnai, 33°53.89'N; 48°34.03'E, 2150 m, 22.07.2007, the same collectors as the holotype. Field codes V331, 335, V336, Iran, Lorestan province, Zagros Mt., W of Borujerd, Kuh-e Garin mount., Vennnai, 33°53.89'N; 48°34.03'E, 2150 m, 02.08.2014, N. Shapoval and A. Barabanov leg. Field codes U169, U178, U179, Iran, Lorestan province, Zagros Mt., Darreh Takht, 33° 21.19'N; 49° 22.34'E, 2000–2100 m, 18.07.2011, the same collectors as the holotype. 1 ♀, field code U169 Iran, Lorestan province, Zagros Mt., Darreh Takht, 33° 21.19'N; 49° 22.34'E, 2000–2100 m, 18.07.2011, same collectors as the holotype. All paratypes are kept in the Zoological Institute of the Russian Academy of Sciences (St. Petersburg). GenBank accession numbers of the paratypes are presented in the Table 1.

Derivatio nominis. The new taxon is named after the village Saravand, one of the places where it was found.

Description. *Male upperside.* Forewing length 30–36 mm, ground colour bright blue with azure tint. Discoidal, submarginal and antemarginal marking absent on both fore- and hindwings. Black outer marginal line on forewings and hindwings very narrow; forewing hind margin with long white pubescence. Fringes of both wings dark grey; tips of hindwings veins indicated with fine black.

Male underside. Ground colour light grey, white streak on the hindwings absent. Basal black spots present only on hindwings. Discoidal series of spots present on fore- and hindwings, although the black spots composing it are minute. Postdiscal black marking very narrow, longitudinal, present only on forewings. Submarginal and marginal lunules only faintly indicated.

Female upperside. Ground colour brown with vastly darker veins. Discoidal black spots present on forewings. Submarginal markings dark brown with orange submarginal lunules well developed on forewing and hindwing. Fringe greyish-brown.

Female underside. General design as in males, but ground colour slightly darker.

Genitalia. The male genitalia have a structure typical for other species of the subgenus *Agrodiaetus* (Coutsis 1986). No specific characters in genitalia are found.

Diagnosis. Genetically *P. (A.) karindus saravandi* differs from all other taxa of *Agrodiaetus* by fixed substitutions in mitochondrial gene *COI*. Phenotypically the new taxon is extremely similar to *P. (A.) karindus karindus* from north-west Iran, but they have different chromosome numbers, n=73 and n = 68 respectively.

Distribution. Central part of Zagros Mountains, Iran.

Flight period. From July to August.

Ecology. Dry slopes, gorges and plateaus with xerophyte or steppe vegetation, sometimes wooded areas from 1800 up to 2800 m. Butterflies fly together with *P. (A.) alcestis* (Zerny, 1932), *P. (A.) cyaneus* (Staudinger, 1899), *P. (A.) hamadanensis* (de Lesse, 1959), *P. (A.) lorestanus* (Eckweiler, 1997) and *P. (A.) zarathustra* (Eckweiler, 1997).

Acknowledgements

The complete financial support for this study was provided by the grant N 14–14–00541 from the Russian Science Foundation to the Zoological Institute of the Russian Academy of Sciences. Postdoctoral fellowship (N 1.50.1617.2013) was provided to N. Shapoval from St. Petersburg State University. We thank A. Barabanov for help in obtaining material during the field expedition to Iran in 2014. The work was partially performed using equipment of the ‘Chromas’ Core Facility and Centre for Molecular and Cell Technologies of St. Petersburg State University.

References

- Bálint Z (1999) Annotated list of type specimens of *Polyommatus* sensu Eliot of the Natural History Museum, London (Lepidoptera, Lycaenidae). Neue Entomologische Nachrichten 46: 1–89.
- Caterino MS, Sperling FAH (1999) *Papilio* phylogeny based on mitochondrial *cytochrome Oxidase I* and *II* genes. Molecular Phylogenetics and Evolution 11(1): 122–137. doi: 10.1006/mpev.1998.0549
- Clement M, Posada D, Crandall K (2000) TCS: a computer program to estimate gene genealogies. Molecular Ecology 9: 1657–1660. doi: 10.1046/j.1365-294x.2000.01020.x
- Coutsis JG (1986) The blue butterflies of the genus *Agrodiaetus* Hübner (Lep., Lycaenidae): symptoms of taxonomic confusion. Nota Lepidopterologica 9: 159–169.
- Coyne JA, Orr AH (2004) Speciation. Sinauer Associates, Sunderland, 480 pp.
- de Lesse H (1957) Description de deux nouvelles espèces d'*Agrodiaetus* (Lep. Lycaenidae). Lambillionea 57(9–10): 65–71.
- de Lesse H (1959a) Description d'une nouvelle sous-espèce d'*Agrodiaetus dama* Stgr. (Lep. Lycaenidae). Bulletin de la Société Entomologique de Mulhouse, 13–15.
- de Lesse H (1959b) Note sur deux espèce d'*Agrodiaetus* (Lep. Lycaenidae) récemment séparées d'après leurs formules chromosomiques. Lambillionea 59(1–2): 5–10.
- de Lesse H (1959c) Nouvelle étude chorologique et cytologique conduisant à changer l'application du nom d'*Agrodiaetus iphigenia* H. S. (Lep. Lycaenidae). Lambillionea 59(7–8): 56–61.
- de Lesse H (1959d) Sur la valeur spécifique de deux sous-espèce d'*Agrodiaetus* (Lep. Lycaenidae) récemment décrites. Bulletin Mensuel de la Société Linnéenne de Lyon 28(10): 312–315.

- de Lesse H (1960a) Les nombres de chromosomes dans la classification du groupe d'*Agrodiaetus ripartii* Freyer (Lepidoptera, Lycaenidae). Revue Française d'Entomologie 27: 240–264.
- de Lesse H (1960b) *Agrodiaetus iphigenia* H.S. et son espèce jumelle *A. tankeri* n. sp. Séparées d'après sa formule chromosomique. Bulletin de la Société Entomologique de Mulhouse, 75–78.
- de Lesse H (1961a) Les nombres de chromosomes chez *Agrodiaetus dolus* Hübner et les espèces voisines (Lycaenidae). Alexanor 2(2): 57–63.
- de Lesse H (1962a) Variation chromosomique chez *Agrodiaetus actis* H.S. et *A. altivagans* Forst. (Lep. Lycaenidae). Revue Française d'Entomologie 29: 66–67.
- de Lesse H (1962b) Variation chromosomique chez *Agrodiaetus dolus* Hübner (Lep. Lycaenidae). Alexanor 2: 283–286.
- de Lesse H (1963a) Variation chromosomique chez *Agrodiaetus carmon* H.S. et *A. cyanea* Stgr. (Lep. Lycaenidae). Revue Française d'Entomologie 30(3): 177–181.
- de Lesse H (1963b) Variation chromosomique chez les *Agrodiaetus* (Lep. Lycaenidae). Revue Française d'Entomologie 30(3): 182–189.
- de Lesse H (1964) Une nouvelle sous-espèce d'*Agrodiaetus altivagans* Forster (Lep. Lycaenidae). Alexanor 4: 167–168.
- de Lesse H (1966) Variation chromosomique chez *Agrodiaetus dolus* Hübner (Lep. Lycaenidae). Annales de la Société Entomologique de France 2(1): 209–214.
- de Lesse H (1968) *Agrodiaetus ripartii* Frey dans la région de Barcelone (Lycaenidae). Alexanor 5(5): 203–205.
- Dincă V, Runquist M, Nilsson M, Vila R (2013) Dispersal, fragmentation and isolation shape the phylogeography of the European lineages of *Polyommatus (Agrodiaetus) ripartii* (Lepidoptera: Lycaenidae). Biological Journal of the Linnean Society 109: 817–829. doi: 10.1111/bij.12096
- Forster W (1956) Bausteine zur Kenntnis der Gattung *Agrodiaetus* Scudd. (Lep. Lycaen) I. Zeitschrift der Wiener Entomologischen Gesellschaft 41: 42–61, 70–89, 118–127.
- Forster W (1960a) Bausteine zur Kenntnis der Gattung *Agrodiaetus* Scudd. (Lep. Lycaen) II. Zeitschrift der Wiener Entomologischen Gesellschaft 45: 105–142.
- Forster W (1960b) Einige neue Formen der Gattung *Agrodiaetus* Scudd. (Lepidoptera, Lycaenidae). Entomologische Zeitschrift 70(3): 17–22.
- Forster W (1960c) Bausteine zur Kenntnis der Gattung *Agrodiaetus* Scudd. (Lep. Lycaen) II. Zeitschrift der Wiener Entomologischen Gesellschaft 45: 105–142.
- Forster W (1961) Bausteine zur Kenntnis der Gattung *Agrodiaetus* Scudd. (Lep. Lycaen) II. Zeitschrift der Wiener Entomologischen Gesellschaft 46: 8–13, 38–47, 74–79, 88–94, 110–116.
- Hall T (2011) BioEdit: An important software for molecular biology. GERF Bulletin of Biosciences 2(1): 60–61.
- Hesselbarth G, van Oorschot H, Wagener S (1995) Die Tagfalter der Türkei unter Berücksichtigung der angrenzenden Länder. Bocholt, 2200 pp.
- Kandul NP, Lukhtanov VA, Pierce NE (2007) Karyotypic diversity and speciation in *Agrodiaetus* butterflies. Evolution 61(3): 546–559. doi: 10.1111/j.1558-5646.2007.00046.x

- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thirer T, Ashton B, Mentjies P, Drummond A (2012) Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28(12): 1647–1649. doi: 10.1093/bioinformatics/bts199
- King M (1993) Species evolution: the role of chromosomal change. Cambridge University Press, Cambridge, 358 pp.
- Lukhtanov VA, Dantchenko AV (2002) Descriptions of new taxa of the genus *Agrodiaetus* Hubner, [1822] based on karyotype investigation (Lepidoptera, Lycaenidae). *Atalanta* 33(1/2): 81–107, 224–225.
- Lukhtanov VA, Dantchenko AV (2003) Two new chromosome races of the *Agrodiaetus altivagans-wagneri* complex (Lepidoptera, Lycaenidae). *Atalanta* 34(3–4): 421–428, 480–481.
- Lukhtanov VA, Tikhonov AV (2015) Chromosomal and molecular evidence for presence of *Polyommatus* (*Agrodiaetus*) *poseidon* (Lepidoptera, Lycaenidae) in Caucasus region. *Comparative Cytogenetics* 9(2): 249–255. doi: 10.3897/CompCytogen.v9i2.5020
- Lukhtanov VA, Shapoval NA, Dantchenko AV (2008) *Agrodiaetus shahkuhensis* sp. n. (Lepidoptera, Lycaenidae), a cryptic species from Iran discovered by using molecular and chromosomal markers. *Comparative Cytogenetics* 2(2): 99–114.
- Lukhtanov VA, Shapoval NA, Dantchenko AV (2014) Taxonomic position of several enigmatic *Polyommatus* (*Agrodiaetus*) species (Lepidoptera, Lycaenidae) from Central and Eastern Iran: insights from molecular and chromosomal data. *Comparative Cytogenetics* 8 (4): 313–322. doi: 10.3897/CompCytogen.v8i4.8939
- Lukhtanov VA, Vila R, Kandul NP (2006) Rearrangement of the *Agrodiaetus dolus* species group (Lepidoptera, Lycaenidae) using a new cytological approach and molecular data. *Insect Systematics and Evolution* 37(3): 325–334. doi: 10.1163/187631206788838563
- Lukhtanov VA, Dantchenko AV, Vishnevskaya MS, Saifitdinova AF (2015a) Detecting cryptic species in sympatry and allopatry: analysis of hidden diversity in *Polyommatus* (*Agrodiaetus*) butterflies (Lepidoptera: Lycaenidae). *Biological Journal of the Linnean Society* 116(2): 468–485. doi: 10.1111/bij.12596
- Lukhtanov VA, Shapoval NA, Anokhin BA, Saifitdinova AF, Kuznetsova VG (2015b) Homoploid hybrid speciation and genome evolution via chromosome sorting. *Proceedings of the Royal Society B* 282: 20150157. doi: 10.1098/rspb.2015.0157
- Lukhtanov VA, Kandul NP, Plotkin JB, Dantchenko AV, Haig D, Pierce NE (2005) Reinforcement of pre-zygotic isolation and karyotype evolution in *Agrodiaetus* butterflies. *Nature* 436(7049): 385–389. doi: 10.1038/nature03704
- Oliver A, de Prins W, Van Der Poorten D, Puplesiene J (1999) On the identity of *Polyommatus* (*Agrodiaetus*) *dama*, with lectotype designation and redescription of its karyotype (Lycaenidae). *Nota Lepidopterologica* 22(3): 197–211.
- Posada D (2008) jModel Test: phylogenetic model averaging. *Molecular Biology and Evolution* 25: 1253–1256. doi: 10.1093/molbev/msn083
- Przybyłowicz Ł, Lukhtanov V, Lachowska-Cierlik D (2014) Towards the understanding of the origin of the Polish remote population of *Polyommatus* (*Agrodiaetus*) *ripartii* (Lepidoptera:

- Lycaenidae) based on karyology and molecular phylogeny. *Journal of Zoological Systematics and Evolutionary Research* 52(1): 44–51. doi: 10.1111/jzs.12040
- Riley ND (1921) Some undescribed Rhopalocera from Mesopotamia and N. W. Persia; and other notes. *The Annals and Magazine of Natural History* 8: 59–600. doi: 10.1080/00222-932108632622
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3): 539–542. doi: 10.1093/sysbio/sys029
- Talavera G, Lukhtanov VA, Pierce NE, Vila R (2013a) Establishing criteria for higher-level classification using molecular data: the systematics of *Polyommatus* blue butterflies (Lepidoptera, Lycaenidae). *Cladistics* 29: 166–192. doi: 10.1111/j.1096-0031.2012.00421.x
- Talavera G, Lukhtanov VA, Rieppel L, Pierce NE, Vila R (2013b) In the shadow of phylogenetic uncertainty: the recent diversification of *Lysandra* butterflies through chromosomal change. *Molecular Phylogenetics and Evolution* 69: 469–478.
- Vershinina AO, Lukhtanov VA (2010) Geographical distribution of the cryptic species *Agrodiaetus alcestis alcestis*, *A. alcestis karacetinae* and *A. demavendi* (Lepidoptera, Lycaenidae) revealed by cytogenetic analysis. *Comparative Cytogenetics* 4(1): 1–11. doi: 10.3897/compcytogen.v4i1.21
- Vila R, Lukhtanov VA, Talavera G, Gil-T F, Pierce NE (2010) How common are dot-like distribution ranges? Taxonomical oversplitting in Western European *Agrodiaetus* (Lepidoptera, Lycaenidae) revealed by chromosomal and molecular markers. *Biological Journal of the Linnean Society* 101: 130–154. doi: 10.1111/j.1095-8312.2010.01481.x
- Wiemers M (2003) Chromosome differentiation and the radiation of the butterfly subgenus *Agrodiaetus* (Lepidoptera: Lycaenidae: *Polyommatus*) a molecular phylogenetic approach. PhD thesis, University of Bonn, Bonn, Germany, 203 pp. <http://hss.ulb.uni-bonn.de/2003/0278/0278.htm>