

# Three new species of the genus *Leptonetela* from Greece (Araneae, Leptonetidae)

Yi Wu<sup>1</sup>, Chunxia Wang<sup>2</sup>, Guo Zheng<sup>1</sup>, Shuqiang Li<sup>2</sup>

**1** College of Life Science, Shenyang Normal University, Shenyang 110034, China **2** Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China

Corresponding authors: Guo Zheng ([zhengguo@synu.edu.cn](mailto:zhengguo@synu.edu.cn)); Shuqiang Li ([lisq@ioz.ac.cn](mailto:lisq@ioz.ac.cn))

Academic editor: Y. Marusik | Received 21 October 2015 | Accepted 28 January 2016 | Published 24 February 2016

<http://zoobank.org/9EFB75EF-B126-441A-93DD-9FB14765F125>

**Citation:** Wu Y, Wang C, Zheng G, Li S (2016) Three new species of the genus *Leptonetela* from Greece (Araneae, Leptonetidae). ZooKeys 569: 23–35. doi: 10.3897/zookeys.569.6921

## Abstract

Three new species of the spider genus *Leptonetela* collected from caves in Greece are described: *L. arvanitidis* **sp. n.** (male & female), *L. paragamiani* **sp. n.** (male & female) and *L. penevi* **sp. n.** (male & female). Detailed illustrations of the new species are provided. DNA barcodes were obtained for future use.

## Keywords

Haplogynae, taxonomy, DNA-barcoding, Balkan Peninsula, spider

## Introduction

The leptonetids are minute (1.0–3.0 mm) spiders that can be easily distinguished from other families by a distinctive 6-eyed pattern, with the posterior median eyes situated behind the posterior lateral eyes; however, in some cave species, the eyes are reduced to vestiges or may be completely absent (Gertsch 1974).

A total of 23 genera and 276 species of the spider family Leptonetidae are known worldwide (World Spider Catalog 2016). The genus *Leptonetela* was established by Kratochvíl (1978), using *Leptonetela kanellisi* (Deeleman-Reinhold, 1971) from a cave in Greece as the type species. *Leptonetela* can be distinguished from other genera of the family by the palpal femur without spines, the retrolateral surface of the palpal tibia with a longitudinal row of strong spines and the male palpal tarsus without appendices.

A total of 50 *Leptonetela* species are known from Europe and Asia. Two species of *Protoleptoneta* were transferred to *Leptonetela* by Brignoli in 1979: *L. strinatii* (Brignoli, 1976) from Greece and *L. deltshevi* (Brignoli, 1979) from Turkey. Deltshev described *L. andreevi* from Greece in 1985. Dunin (1990) reported *L. caucasica* from Georgia and Azerbaijan. *L. thracia* was described by Gasparo in 2005 from Greece. Subsequently, Lin and Li (2010) described 24 species occurring in the Yunnan-Guizhou Plateau, China, including *L. quinquespinata* (Chen & Zhu, 2008) which was transferred from *Qianleptoneta* Chen & Zhu, 2008. Wang and Li (2011) reported 17 *Leptonetela* species from South China, 2 species from Greece and 1 species from Vietnam.

Other than *L. deltshevi* (Brignoli, 1979) from Turkey and *L. pungitia* Wang & Li, 2011 from Vietnam which have been collected in epigeal habitats, all species are found in caves. Some of them have characters typical to true troglobites, such as lacking eyes and pigmentation and elongated legs.

In this paper, three *Leptonetela* species collected from caves in Greece are described as new to science. The total number of *Leptonetela* species from Europe reaches 9 species.

## Material and methods

Specimens were examined with a LEICA M205C stereomicroscope. Images were captured with an Olympus C7070 wide zoom digital camera (7.1 megapixels) mounted on an Olympus SZX12 dissecting microscope. Epigynes and male palps were examined after dissection from the spiders' bodies.

Terminology and abbreviations in this paper generally follow Wang and Li (2011) and Ledford (2011). The unit of measurement in this paper is millimetres (mm). Leg metric data were recorded as total length (femur, patella, tibia, metatarsus, tarsus). Leg segments were measured on their dorsal side.

DNA barcodes were obtained for future use. A partial fragment of the mitochondrial gene cytochrome oxidase subunit I (COI) was amplified and sequenced for *Leptonetela arvanitidisi* sp. n., *L. paragamiani* sp. n. and *L. penevi* sp. n. following the protocol in Miller et al. (2010). Primers used in this study are: LCO1490 (5'-CWACAAAY-CATARRGATATTGG-3') and HCO-N-2198 (5'-TAAACTTCAGGGTGAC-CAAAAAATCA-3') (Folmer et al. 1994). Voucher information and GenBank accession number for all samples are listed in Table 1.

The specimens studied in the current paper are deposited in the Institute of Zoology, Chinese Academy of Sciences (IZCAS) in Beijing, China.

**Table 1.** Voucher specimen information.

Species	Sequence length	Collecting localities	GenBank accession number
<i>L. arvanitidisi</i> sp. n.	620 bp	Greece Athens Attica: Leondari Cave	KU318407
<i>L. paragamiani</i> sp. n.	620 bp	Greece Athens: Pan Cave	KU318410
<i>L. penevi</i> sp. n.	620 bp	Greece Thiva: Skoteini Cave	KU318411

## Taxonomy

### Family Leptonetidae Simon, 1890

### Genus *Leptonetela* Kratochvíl, 1978

*Leptonetela*: Kratochvíl 1978: 11, f. 1G. Type species *Sulcia kanellisi* Deeleman-Reinhold, 1971 from Greece.

### *Leptonetela arvanitidisi* Wang & Li, sp. n.

<http://zoobank.org/24A56F21-B537-4A53-B19B-53A86AA86C96>

Figs 1–2, 7

**Types.** Holotype ♂ (IZCAS): GREECE, Athens, Attica, Leondari Cave, 37°59'14.61"N, 23°49'47.03"E, elevation 553 m, 28 March 2013, S. Li leg. Paratypes 2 ♀ (IZCAS), same data as holotype.


**Etymology.** The specific name is dedicated to Dr. Christos Arvanitidis of the Hellenic Centre for Marine Research in Crete, a leading taxonomist on Polychaeta; noun (name) in genitive case.

**Diagnosis.** *Leptonetela arvanitidisi* sp. n. is similar to *L. kanellisi* but can be separated by the basal tibial spine with bifurcated tip (Fig. 1D) (not bifurcated in *L. kanellisi*), the wave-shaped anterior margin of the atrium and the tightly twisted spermathecae (Fig. 2C); *L. kanellisi* has an arc-shaped anterior margin of the atrium and the spermathecae are loosely twisted (see Wang and Li 2011: figs 16–19).


**Description.** Male (holotype). Total length 1.75 (Fig. 1A). Carapace 0.83 long, 0.72 wide. Opisthosoma 1.00 long, 0.65 wide. Prosoma yellowish, with one seta in the middle of the carapace. Ocular area with a pair of setae, eyes absent. Median groove, cervical grooves and radial furrows indistinct. Clypeus 0.10 high. Sternum and legs yellowish. Opisthosoma pale brown, ovoid, lacking distinctive pattern. Leg measurements: I 8.39 (2.25, 0.38, 2.45, 2.08, 1.23); II 7.14 (1.88, 0.38, 2.03, 1.72, 1.13); III 5.93 (1.73, 0.35, 1.62, 1.55, 0.68); IV 7.83 (2.13, 0.38, 2.15, 2.00, 1.17). Male palp (Fig. 1C–D): tibia with 5 spines retrolaterally, the basal one strong, conspicuous, with bifurcated tip. Bulb with triangular embolus; prolateral lobe oval. Median apophysis (Fig. 1B) distal edge round, with six small teeth. Conductor membranous, triangular in ventral view.

Female (one of the paratypes). Similar to male in color and general features but larger and with shorter legs. Total length 2.03 (Fig. 2A–B). Carapace 0.85 long, 0.73 wide. Opisthosoma 1.23 long, 0.90 wide. Clypeus 0.10 high. Leg measurements: I 7.24 (1.90, 0.38, 2.08, 1.75, 1.13); II 5.92 (1.68, 0.33, 1.63, 1.38, 0.90); III 5.32 (1.50, 0.32, 1.42, 1.30, 0.78); IV 6.63 (1.75, 0.35, 1.80, 1.70, 1.03). Vulva (Fig. 2C): spermathecae coiled, atrium fusiform, anterior margin of the atrium wave shaped.

**Distribution.** Known only from the type locality.



**Figure 1.** *Leptonetela arvanitidisi* sp. n., holotype male. **A** Habitus, dorsal view **B** Palpal bulb, ventral view **C** Palp, prolateral view **D** Palp, retrolateral view.



**Figure 2.** *Leptonetela arvanitidisi* sp. n., one of the paratype female. **A** Habitus, dorsal view **B** Habitus, ventral view **C** Vulva, dorsal view.

***Leptonetela paragamiani* Wang & Li, sp. n.**

<http://zoobank.org/28C5914F-58AB-409B-8110-61F5B1D5004D>

Figs 3–4, 7

**Types.** Holotype ♂ (IZCAS): GREECE, near Athens, Pan Cave, 38°08'48.54"N, 23°40'06.04"E, elevation 660 m, 7 April, 2013, S. Li leg. Paratypes 2 ♀ (IZCAS), same data as holotype.

**Etymology.** The specific name is dedicated to Mr. Kaloust Paragamian of the Hellenic Institute of Speleological Research in Crete, a leading speleologist in Greece; noun (name) in genitive case.

**Diagnosis.** *Leptonetela paragamiani* is similar to *L. kanellisi* and *L. arvanitidisi* sp. n. but can be separated by the second tibial spine, which is longest in *L. paragamiani* sp. n., whereas in *L. kanellisi* and *L. arvanitidisi* sp. n. (Fig. 3D) the basal spine is longest; the median apophysis has 3 small teeth (Fig. 3B) in *L. paragamiani* sp. n., whereas it has 6 teeth in *L. kanellisi* and *L. arvanitidisi* sp. n.; and the spermathecae are tightly twisted (Fig. 4C) compared to the spermathecae of *L. kanellisi* and *L. arvanitidisi* sp. n.

**Description.** Male (holotype). Total length 1.63 (Fig. 3A). Carapace 0.75 long, 0.62 wide. Opisthosoma 0.88 long, 0.62 wide. Prosoma yellowish, with one seta on the median part. Ocular area with a pair of setae, two eyes, reduced to white spots. Median groove, cervical groove and radial furrows indistinct. Clypeus 0.10 high. Sternum and legs yellowish. Opisthosoma pale brown, ovoid, lacking distinctive pattern. Leg measurements: I 5.53 (1.50, 0.28, 1.60, 1.27, 0.88); II 4.78 (1.38, 0.25, 1.27, 1.13, 0.75); III 4.01 (1.13, 0.25, 1.03, 1.00, 0.60); IV 5.25 (1.45, 0.28, 1.47, 1.25, 0.80). Male palp (Fig. 3C–D): tibia with 5 retrolateral spines, the basal one strong, conspicuous, and the second one longer than others. Bulb with spoon-shaped embolus, prolateral lobe oval. Distal edge of median apophysis round (Fig. 3B), with three small teeth, conductor membranous, shield shaped in ventral view.

Female (one of the paratypes). Similar to male in color and general features but larger and with longer legs. Total length 1.88 (Fig. 4A–B). Carapace 0.75 long, 0.68 wide. Opisthosoma 1.00 long, 0.88 wide. Clypeus 0.10 high. Leg measurements: I 6.26 (1.75, 0.28, 1.80, 1.53, 0.90); II 5.36 (1.58, 0.28, 1.50, 1.25, 0.75); III 4.69 (1.38, 0.25, 1.25, 1.13, 0.68); IV 6.19 (1.78, 0.28, 1.75, 1.50, 0.88). Vulva (Fig. 4C): spermathecae twisted, atrium oval.


**Distribution.** Known only from the type locality.

***Leptonetela penevi* Wang & Li, sp. n.**


<http://zoobank.org/C819DE91-5E16-4B18-B336-26AE51FFD15F>

Figs 5–6, 7


**Types.** Holotype ♂ (IZCAS): GREECE, Thiva, Kakalitsa, Skoteini Cave, 38°29'59.81"N, 23°59'01.06"E, elevation 443 m, 29 March, 2013, S. Li leg. Paratypes 2 ♀, same data as holotype.




**Figure 3.** *Leptonetela paragamiani* sp. n., holotype male. **A** Habitus, dorsal view **B** Palpal bulb, ventral view **C** Palp, prolateral view **D** Palp, retrolateral view.




**Figure 4.** *Leptonetela paragamiani* sp. n., one of the paratype female. **A** Habitus, dorsal view **B** Habitus, ventral view **C** Vulva, dorsal view.



**Figure 5.** *Leptonetela penevi* sp. n., holotype male. **A** Habitus, dorsal view **B** Palpal bulb, ventral view **C** Palp, prolateral view **D** Palp, retrolateral view.



**Figure 6.** *Leptonetela penevi* sp. n., one of the paratype female. **A** Habitus, dorsal view **B** Habitus, ventral view **C** Vulva, dorsal view.



**Figure 7.** Locality records for three new species of *Leptonetela* in Greece: ① *L. arvanitidisi* sp. n. (Athens) ② *L. paragamiani* sp. n. (near Athens) ③ *L. penevi* sp. n. (Thiva).

**Etymology.** The specific name is dedicated to Prof. Dr. Lyubomir Penev, zoologist and founder of Pensoft Publishers; noun (name) in genitive case. Pensoft Publishers is a leading company in publishing taxonomic works.

**Diagnosis.** *Leptonetela penevi* sp. n. is similar to *L. kanellisi* and *L. paragamiani* sp. n. but can be separated by having the basal tibial spine longer than others, and slender (Fig. 5D) compared to the basal spines of *L. kanellisi* and *L. paragamiani* sp. n.; median apophysis distally without teeth (Fig. 5D) and spermathecae strongly twisted and longer than those of *L. kanellisi* and *L. paragamiani* sp. n. (Fig. 6C).

**Description.** Male (holotype). Total length 1.83 (Fig. 6A). Carapace 0.77 long, 0.65 wide. Opisthosoma 1.15 long, 0.80 wide. Prosoma yellowish, with one seta on the median part. Ocular area with a pair of setae, eyes absent. Median groove, cervical groove and radial furrows indistinct. Clypeus 0.10 high. Sternum and legs yellowish. Opisthosoma pale brown, ovoid, lacking distinctive pattern. Leg measurements: I 6.76 (1.88, 0.38, 1.87, 1.55, 1.08); II 5.44 (1.38, 0.33, 1.58, 1.27, 0.88); III 4.87 (1.37, 0.30, 1.25, 1.20, 0.75); IV 6.32 (1.82, 0.35, 1.73, 1.50, 0.92). Male palp (Fig. 5C–D): tibia with 5 spines retrolaterally, with the basal one strong, conspicuous, and longest. Bulb oval, with spoon-shaped embolus, prolateral lobe oval. Median apophysis (Fig. 5B) without teeth distally, conductor membranous, rugose and shield shaped in ventral view.

Female (one of the paratypes). Similar to male in color and general features but larger and with shorter legs. Total length 2.03 (Fig. 6A–B). Carapace 0.75 long, 0.72 wide. Opisthosoma 1.38 long, 0.85 wide. Clypeus 0.10 high. Leg measurements: I 6.51 (1.88, 0.38, 1.83, 1.50, 0.92); II 5.54 (1.63, 0.33, 1.55, 1.25, 0.78); III 4.91 (1.42, 0.33, 1.28, 1.13, 0.75); IV 6.31 (1.80, 0.35, 1.80, 1.48, 0.88). Vulva (Fig. 6C): spermathecae strongly twisted, atrium oval.

**Distribution.** Known only from the type locality.

## Acknowledgements

The manuscript benefitted greatly from comments by Yuri M. Marusik (Magadan, Russia), Christo Deltshv (Sofia, Bulgaria) and Joel Ledford (San Francisco, USA). English of the final draft was kindly checked by Sarah Crews. This study was financially supported by the National Natural Sciences Foundation of China to Guo Zheng (NSFC-31172121, 31372224) and Shuqiang Li (NSFC-31272280, 31471960, 31530067). Part of the laboratory work was financially supported by Eco-Environmental Research Center Foundation of Shenyang Normal University (EERC-T-201502), Liaoning Excellent Talents in University (LJQ2012094).

## Reference

- Brignoli PM (1979) Spiders from Turkey, VI. Four new species from the coast of the Black Sea (Araneae). *Bulletin of the British Arachnological Society* 4: 310–313.
- Deltshv CD (1985) New data concerning cave spiders (Araneae) in Greece with description of a new *Leptonetela* (Araneae, Leptonetidae). *Acta Zoologica Bulgarica* 27: 41–45.
- Dunin PM (1990) *Leptonetela caucasica* sp. n. - a first finding of spiders of the family Leptonetidae (Aranei, Haplogynae) in the USSR. *Zoologicheskii Zhurnal* 69(1): 147–149.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3(5): 294–299.
- Gasparo F (2005) Una nuova *Leptonetela cavernicola* di Grecia (Araneae, Leptonetidae). *Bollettino del Museo Regionale di Scienze Naturali di Torino* 22: 517–524.
- Gertsch WJ (1974) The spider family Leptonetidae in North America. *The Journal of Arachnology*, 45–203.
- Kratochvíl J (1978) Araignées cavernicoles des îles Dalmates. *Přírodovědné práce ústavů Československé akademie věd v Brně (N. S.)* 12(4): 1–59.
- Ledford J, Paquin P, Cokendolpher J, Campbell J, Griswold C (2011) Systematics of the spider genus *Neoleptoneta* Brignoli, 1972 (Araneae: Leptonetidae) with a discussion of the morphology and relationships for the North American Leptonetidae. *Invertebrate Systematics* 25: 334–388. doi: 10.1071/IS11014

- Lin Y, Li S (2010) Leptonetid spiders from caves of the Yunnan-Guizhou plateau, China (Araneae: Leptonetidae). *Zootaxa* 2587: 1–93.
- Miller JA, Carmichael A, Ramirez MJ, Spagna JC, Haddad CR, Řezáč M, Johannesen J, Král J, Wang XP, Griswold CE (2010) Phylogeny of entelegyne spiders: Affinities of the family Penestomidae (NEW RANK), generic phylogeny of Eresidae, and asymmetric rates of change in spinning organ evolution (Araneae, Araneoidea, Entelegynae). *Molecular Phylogenetics and Evolution* 55: 786–804. doi: 10.1016/j.ympev.2010.02.021
- Wang C, Li S (2011) A further study on the species of the spider genus *Leptonetela* (Araneae: Leptonetidae). *Zootaxa* 2841: 1–90.
- World Spider Catalog (2016) World Spider Catalog. Natural History Museum Bern. Version 17.0. <http://wsc.nmbe.ch> [accessed 29 January 2016]